
NUTRITION AND ATHEROSCLEROSIS

LOUIS N KATZ M D

*Director Cardiovascular Department Medical Research Institute Michael Reese Hospital
and Professorial Lecturer in Physiology University of Chicago Chicago Illinois*

JEREMIAH STAMLER M D

*Previously Assistant Director Cardiovascular Department Medical Research Institute
Michael Reese Hospital Chicago Illinois and Established Investigator
of the American Heart Association Presently Director
Heart Disease Control Program Chicago
Board of Health Chicago Illinois*

RUTH PICK M D

*Assistant Director Cardiovascular Department Medical Research Institute
Michael Reese Hospital Chicago Illinois and Established Investigator
of the American Heart Association*

67 Illustrations



Lea & Febiger

Philadelphia

1958

© by Lea & Febiger
November 1958

Library of Congress Card Catalog Number 58-14185

Printed in the United States of America

Preface

THIS book is based on the 1956 Henry Jackson Lecture of the New England Cardiovascular Society, given by one of us (L N K.) in Boston on May 14, 1956. In reviewing the original manuscript, it was decided to undertake its expansion into a comprehensive review and appraisal of the rapidly developing field of atherosclerosis research. This review of the literature was undertaken primarily by one of us (J S). In addition, newer data from our own studies have been incorporated. For the documentation of this evaluation, attention was given principally to the newer data, since previous reviews had covered the earlier literature in detail (1, 8, 9, 12, 13, 645). For this reason, the bibliography is in the main made up of references since 1953.

As a result of this review and of our own work to date, we have presented some positive practical suggestions for prophylaxis and therapy in this book. This is an active, positive approach to this disease problem and is in marked contrast to the therapeutic nihilism of only a few years ago. It is an approach fully consonant with the recent advances and the present day status of our knowledge in this field. It is, in our opinion, the only sound conclusion to a review of progress in this research area—for the progress has indeed been formidable—although the work is as yet unfinished.

LOUIS N. KATZ
JEREMIAH STAMLER
RUTH PICK

Chicago, Illinois

Acknowledgments

The long term program of animal experimental clinical and epidemiologic research of the Cardiovascular Department has been made possible by sustained grant support particularly from the Chicago Heart Association the Albert D and Mary Lasker Foundation, and the National Heart Institute National Institutes of Health United States Public Health Service The continued support of the Michael Reese Research Foundation and of Mr Neison Harris and other local private benefactors is gratefully acknowledged Grant support has also been received from Ayerst Laboratories the Lilly Laboratory for Clinical Research Pfizer Laboratories and the Pitman Moore Company The program was also aided by cooperation from several pharmaceutical houses, including Abbott Laboratories Armour Laboratories Corn Products Refining Company Hoffman LaRoche Inc, Lederle Laboratories Merck and Company Parke Davis and Company, Proctor and Gamble Company, Schering Corporation G D Searle Company Upjohn Company, VioBin Corporation and Winthrop Stearns Inc

In the investigation of the possible efficacy of estrogens in the long term therapy of coronary heart disease in middle aged males acknowledgment is gratefully expressed to Dr L A Baker Chief of Medicine, Hines Veterans Administration Hospital Hines Illinois who made a decisive contribution in enlisting the cooperation of the several participating Veterans Administration Hospitals in the Midwest Dr B Kaplan has also made a major contribution throughout latterly he and Dr A Pick have served as clinical coordinators of the entire study This research endeavor has proceeded only by virtue of the conjoint efforts of members of the medical staffs at several Veterans Administration Hospitals including Drs J Berry D T Foxworthy E Kaplan and W R O Connor Hines Hospital Hines Illinois, Drs R Eckhart C Pilz and H Zummerman West Side Hospital Chicago Illinois Drs R E Reichert Jr and J A Wood, Ann Arbor Michigan Drs A P Klotz and W E Larsen Kansas City Missouri Drs M Armstrong W E Connor and W M Kirkendall Iowa City Iowa Drs L J Baer and M R Weed, Dearborn Michigan Drs M W Garry and G Spencer Wood Hospital Milwaukee Wisconsin Drs G Fischer and R Gieselman St Louis

Missouri, Drs W R Cook and A S Glushein, Pittsburgh, Pennsylvania
Drs R J Kinney and J C Parsons, Des Moines, Iowa

Appreciation is also expressed to Drs W S Middleton, J B Barnwell, M Cummings, B Nagler and G Beebe of the Veterans Administration, Washington D C for their continued cooperation

It is also a pleasure to acknowledge the fine cooperation of Drs L A Lewis and I H Page of the Cleveland Clinic, who accomplished the ultracentrifugal lipoprotein analyses on sera of patients in the estrogen evaluation

The following fellows and residents of the Cardiovascular Department made an essential contribution to this clinical investigation through their work with the patients at Michael Reese Hospital Drs D Berkson, M Buxbaum B W Carnow S Cogan D Cohn, H Danneman L Doctor, G Desir, J Frankel, J Glassberg H Gold, M Goldstein, W Harnischfeger, S A Kabins, S Kimmelblat A Lichtman, C Molina, and J Voltman The cooperation of the attending Medical Staff of the Michael Reese Hospital is also gratefully acknowledged

The epidemiologic analyses of the Chicago, the Illinois and the United States mortality data were accomplished through the cooperation of Drs H N Bundesen President Chicago Board of Health O K Sagen, Office of Vital Statistics, Illinois Department of Health and I Moriyama National Office of Vital Statistics United States Public Health Service Excellent cooperation was extended by the staff of the Chicago Board of Health Messrs F Bauer E D Elia C Kolometz M Templeton, Miss M Stahnke and others It is also a pleasure to acknowledge the encouragement and guidance in these studies of Dr J Watt, Director, National Heart Institute National Institutes of Health United States Public Health Service The cooperation in these endeavors of Miss M Bellows, American Heart Association Messrs L deBoer and D Clough, Chicago Heart Association and Dr A E Rikli Chicago Regional Office United States Public Health Service is also gratefully acknowledged

An excellent contribution was also made by the members of the Cardiovascular Department's epidemiologic research team Mrs Y F Hall, Mr M Kjelsberg and Miss W Muller research assistants

In both the clinical investigation on estrogen therapy and the animal experimental research on atherosclerosis outstanding service has been rendered over the years by the staff of the Cardiovascular Department's atherosclerosis team particularly Mrs D Century Mr A Lillis and Mr P Johnson research assistants and Mr D Bryant Mr G Crowley Miss M Michael Mrs F Miller, Mrs C Thompson and Mrs M Van Linscott

It is also a pleasure to express appreciation for the excellent work of the Cardiovascular Department's secretaries especially Mrs M F Cannamore and Miss B Petzold who unstintingly contributed their painstaking

and meticulous efforts in the preparation of the manuscript legends and bibliography

It is also a pleasure to express appreciation to Mr W Sharer, Miss M Hortatsos and the staff of Wesley Sharer and Associates for their fine cooperation in the preparation of the figures

Acknowledgment is gratefully extended to the many authors and publishers who graciously granted permission to reproduce data and figures from reports in print

Contents

Chapter 1	Introduction	13
Chapter 2	Nutrition and Atherosclerosis	
A	The Nutritional Metabolic Cholesterol Lipid Lipoprotein Theory of Atherogenesis	15
B	The Epidemiologic Method	16
C	Human Nutrition—Historical and Ecological Background	16
Chapter 3	Nutrition and Atherosclerosis (continued)	
D	The Epidemiologic Findings	19
1	Economically Underdeveloped Countries vs the United States—Habitual Diet and Atherogenesis	19
2	Economically Underdeveloped Countries—Social Class Differences in Diet and Atherogenesis	20
3	Economically Underdeveloped Countries vs the United States—Correlations among Diet Serum Cholesterol Lipids-Lipoproteins and Atherogenesis	20
4	Italy and Japan vs the United States	24
5	Effects of Emigration—Yemenite Jews Italians and Japanese	26
6	The Low Countries the Scandinavian Countries <i>et al</i> and the United States	26
7	World Wars I and II	30
8	The United States—1910 to 1955	30
9	The United States—Morbidity and Mortality Findings in Different Population Groups and Geographic Areas	35
10	The United States—Serum Cholesterol Findings in Different Population Groups and Geographic Areas	40
11	The United States—Findings in Prospective Studies on the Living Population	41
Chapter 4	Nutrition and Atherosclerosis (continued)	
E	The Clinical Pathologic Findings	44
1	Historical Development	44

2	Morphologic Atherosclerosis in Cases with and without Clinical Coronary Heart Disease	44
3	Atherosclerosis in Nephrosis, Hypothyroidism, Diabetes, Essential Familial Xanthomatosis	45
4	Serum Cholesterol Lipid Lipoprotein Levels in Clinical Coronary Heart Disease	46
5	Dietary Effects on Cholesterolemia—Weight Gain and Loss, Low Fat Low Cholesterol Diets	48
6	Dietary Effects on Cholesterolemia—Unsaturated Vegetable and Marine Oils—Complete Substitution, Partial Substitution and Supplementation Studies	50
7	Essential Fatty Acids and the Mechanism of the Effects of Unsaturated Oils	56

Chapter 5 Nutrition and Atherosclerosis (continued)

I	The Animal Experimental Findings	61
1	Historical Development	61
2	Diet induced Hypercholesterolemia and Atherogenesis in Different Species	61
3	Dietary Cholesterol and Lipid in Experimental Atherogenesis	63
4	Diet induced Reversibility of Experimental Atherosclerosis	64
5	Unsaturated Vegetable Oils—Essential Fatty Acids	64
6	Corn Germ Sitosterols Dihydrocholesterol Brain Lx tracts	69
7	Nutritional Interrelationships in Experimental Hypercholesterolemia and Atherogenesis	70

Chapter 6 Hormonal and Other Endogenous Factors and Atherosclerosis

A	The Key but Not Exclusive Role of Diet in Atherogenesis—Interrelationships between Diet and Other Factors (Heredity, Physical Activity Stress, Endocrine Function Smoking and the Like)	76
B	Blood Clotting and Fibrinolytic Mechanisms	82
C	Thyroid Hormone	83
D	Estrogens	83
1	Background	83
2	Prophylactic Experiments with Estrogens	83
3	Therapeutic Experiment with Estrogens—Reversibility of Lesions	86
4	Effects of Endogenous Estrogen Secretion—Sex Differential in Chickens	87

5 Effects of Estrogens Combined with Other Hormones	88
6 Effects of Estrogens in Mammalian Species	90
7 Estrogens, Plasma Lipids-Lipoproteins and Atherogenesis in Man	91

Chapter 7 Definitive Approaches to Prophylaxis and Therapy of Atherosclerotic Disease in Man

A The Meaning of <i>Definitive Approaches</i>	93
B Criteria for Evaluating Prophylactic and Therapeutic Regimens	93
C Precautions in Evaluating Prophylactic and Therapeutic Regimens	94
D Specific Proposed Prophylactic and/or Therapeutic Regimens	94
1 Obesity Correction	94
2 Low Fat, Low Cholesterol Diets	95
3 Unsaturated Oils	97
4 Sitosterols	97
5 Lipotropic Factors	98
6 Anticoagulants	98
7 Antihypertensives	99
8 Diabetes Control	100
9 Thyroid Hormone	101
10 Estrogens	101
E Overall Evaluation	106
F An Interim Approach to Prophylaxis for High Risk Persons, and to Therapy	106
G Specific Recommendations	108

Chapter 1

Introduction

ATHEROSCLEROSIS together with hypertension is the most important pathologic process producing morbidity and mortality in the United States today. It is present in the coronary arteries of a majority of American males over 25 years of age (290-292, 294, 300). It produces

US DEATH RATES 1950 --AGE 45-54




FIG. 1. Stamler J. ('53). Death rates (at top of column) in this and subsequent figures are per 100,000 population. Note that arteriosclerotic heart disease (A S H D — category 470 in the International Statistical Classification of Diseases and Causes of Death [I S C]) and arteriosclerotic and degenerative heart disease (A S and D H D — categories 470 + 472 in the I S C) account for approximately one third of all deaths among white males age 45-54. Note the marked sex differential in A S H D mortality rates of the Negro mortality patterns (see text). The category C V R diseases is the broad grouping of the cardiovascular renal diseases.

disease at epidemic rates, thus, the annual incidence rate for coronary heart disease is approximately 1,000 cases per 100,000 males age 45 to 64 (2). It is the number one killer, not only of the elderly, but also of the middle aged (particularly men) in the prime of life (Fig. 1) (23). The current research emphasis on this scourge is, therefore, highly in order.

The intensive investigations of recent years have shed light particularly on the relationships of dietary patterns and endogenous (especially hormonal) factors to the etiology and pathogenesis of morphologic atherosclerosis and clinical atherosclerotic disease (1, 3-9, 12-23, 71, 171-645, 646). Their findings have led to the elaboration of important theoretical concepts which integrate present knowledge, point the way to further study, and presage effective prophylaxis and therapy. This monograph is a review of recent research in these two major areas, including a summation of our group's work during this period. It attempts to delineate the interrelationships between dietary and endogenous factors in the etiology and pathogenesis of atherosclerosis. On this basis it arrives at an evaluation of the current status of the problem of nutrition and atherosclerosis.

Chapter 2

Nutrition and Atherosclerosis

A The Nutritional Metabolic Cholesterol-Lipid-Lipoprotein Theory of Atherogenesis

IN previous reports detailed factual and conceptual elaboration was presented of the nutritional metabolic cholesterol lipid lipoprotein theory of atherogenesis (1, 12-23, 645). The basic ideas may be briefly summarized as background for examining the specific problem of the relationship between diet and atherosclerosis. Atherosclerosis is a distinct entity. It is one among the arterioscleroses by far the most important (1). It is a process separate and different from—but intertwined with—hypertension. Atherosclerosis is the chief pathologic lesion in coronary heart disease (290-292, 294, 297-299). It is a disease. Therefore, it is not inevitable nor is it irreversible. Rather it is preventable and (at least up to a point) curable.

Further, atherosclerosis is a metabolic disease in which altered cholesterol lipid lipoprotein metabolism plays a critical and decisive (but not exclusive) role. Elevated levels of circulating plasma cholesterol lipids-lipoproteins are cardinal signs of this metabolic abnormality and constitute the metabolic prerequisites for atherogenesis in most persons afflicted during middle-age. Without such lipid metabolic derangements clinically significant atherosclerosis would rarely occur in middle age regardless of the functional status of the cardiovascular system and the organism as a whole.

Finally—and this is the decisive point in terms of the problem of the relationship between diet and atherosclerosis—the atherogenic alterations in cholesterol lipid lipoprotein metabolism are frequently brought about by the life span pattern of diet. They are byproducts of an habitually unbalanced diet excessive in total calories, empty calories, total fats, saturated fats, cholesterol, refined carbohydrates, salt, and inadequate (relatively and/or absolutely) in certain essential nutrients (vitamins, minerals, essential amino acids and essential fatty acids) and in bulk. Empty calories are calories derived from processed, refined foods, high in energy value and low in essential nutrients (minerals, vitamins, essential amino acids and essential fatty acids)—foods such as sugar, white

flour breads and pastries, solid cooking and table fats, and many others (16 64)

These concepts concerning the interrelationships among diet, cholesterol lipid lipoprotein metabolism and atherogenesis are documented by abundant data accumulated by the three major investigative approaches to the problem—the epidemiologic, the clinical pathologic and the animal experimental methods of research

Let us examine the evidence from each

B The Epidemiologic Method

The epidemiologic approach (23–32) is essentially demographic and ecologic in orientation focusing on populations and occurrence of disease in them—in contrast to the clinical approach which concentrates on individual patients and groups of sick people irrespective of the populations-at-large from which they originate. Epidemiologic research is based on the premise that important leads to the etiology of disease may be obtained by getting answers to such questions as: Does the incidence of coronary heart disease vary in time and space—and if so, why? Is it greater today than 50 years ago—and if so, why? Is it greater in some countries than in others—and if so, why? Is it greater among males or females, urban or rural dwellers, whites or Negroes, rich or poor, manual (blue collar) or white collar workers—and if there are differences, or similarities in incidence between such groups, why?

The significance of the *why* is that it may give leads to the etiology of disease—it is not enough merely to delineate group trends in incidence of disease. Once patterns of incidence are defined, it becomes essential to proceed to the decisive questions: What accounts for observed differences or similarities? What is the role of diet, occupation, physical activity, smoking, climate, income, housing, mores, tensions, stresses, race, ethnic origin, heredity, genetics, and other factors? And what are the mechanisms whereby specific causative factors operate?

Obviously the complex questions posed by the epidemiologic method can be fruitfully clarified only by extensive cross fertilization among many disciplines (sociology, anthropology, nutrition, psychology, genetics, and others) and by close coordination among medical investigators utilizing the clinical pathologic, animal experimental and epidemiologic approaches.

C Human Nutrition—Historical and Ecological Background

The relationship between diet and atherosclerosis is best understood against the background of certain cardinal facts concerning human nutrition. In terms of man's phylogenesis, biological and sociological, contemporary American patterns of diet represent relatively recent innovations in nutrition. They are diets markedly different in composition

qualitatively and quantitatively, from any ever consumed by wild animals or most pre literate peoples (12, 13, 33) These nutritional patterns are relatively recent products of a long and complex development In terms of the time scale of human evolution their origins go back only about 8,000 years Prior to that time man had been exclusively a food gatherer He had not yet learned to be a food producer Inevitably therefore his eating habits differed markedly from ours (33) Composed of a variety of natural unprocessed foods yielding a high ratio of essential nutrients to calories his diets were invariably very nourishing and well balanced (as long as nature's bounty was adequate) This was true irrespective of their specific patterns, reflective of given local conditions (32-34)

In the Fertile Crescent of the Middle East man made the decisive historic leap from food gathering to food producing and became a farmer and a herder This Neolithic transition created the economic pre-conditions for urban life (35-37) It also made available for the first time many of the foods commonly consumed by most Americans today, *e.g.* dairy products, eggs cereals breads, and other foods (33)

The historic record clearly reveals that the privileged classes in the ancient empires soon learned to enjoy rich "luxus" diets very much like our own This was true of the Egyptian nobility and priesthood—whose mummies reveal bits of aorta with gross atherosclerosis (32) In the words of the Biblical scribes they lived 'off the fat of the land' (38 39)

The bulk of the population however subsisted on very different far simpler fare in Egypt of the Pharaohs it was bread and beer (32) From ancient times right down to the present bread—with rice and potatoes—has been the 'staff of life' For the majority of mankind—particularly in economically underdeveloped Africa Asia Latin America eastern and southern Europe—grains have remained the chief staples supplying 60 to 90 per cent of daily calories (Fig 2) (40-63)

Unquestionably people on such diets are not well provided for nutritionally (66) Many of them suffer from permanent malnutrition—more precisely *undernutrition*

In the economically developed countries a marked change in nutritional patterns has occurred during the last century In conjunction with industrialization urbanization and increase in per capita national income richer diets have become commonplace—diets containing sizable quantities of the more expensive high lipid foods of animal origin (well fattened meats eggs butter and other dairy products) plus 'elegant' white bread and refined sugar These foods are now consumed *en masse* in countries like the United States As a result intake *en masse* of total calories empty calories total fats cholesterol saturated fats has tended to increase significantly in these economically developed countries It should be stressed that the rise in total fat intake is almost exclusively

attributable to the increased ingestion of *saturated* fats of animal origin (*not* unsaturated vegetable and marine oils). It is quite exceptional to find a sizable population consuming a high level of fat, *e.g.* 30 to 40 per cent of total calories, derived chiefly from vegetable sources, *i.e.* mainly unsaturated fat (*cf.* 67, 139, 167, 251). This evolutionary change in dietary habits has been repeatedly observed, as virtually an invariable concomitant of improved economic conditions (40-45, 63, 65-67).

DIET IN MORE DEVELOPED AND LESS DEVELOPED COUNTRIES-1950




FIG 2 Stamler J (22-23) (for source of data *cf.* ref (41)) Fats and oils are the amounts of these foodstuffs available for consumption as such not the total amount of fat derived from all dietary constituents

Hence, longstanding gross undernutrition has ceased to be a significant mass phenomenon in the economically developed countries. Most people ingest a diet providing them with enough calories except under conditions of war or depression. In fact the caloric intake is frequently in excess of the total required to balance energy expenditure with consequent widespread obesity. Moreover, these calories are frequently supplied by a diet of inadequate composition—a diet unbalanced in terms of the proportions among foodstuffs. The nature of the aberration is (1) an excess of total calories empty calories total fat saturated fat cholesterol refined sugar salt and (2) an inadequacy (relative and/or absolute) of essential nutrients (vitamins minerals essential amino acids and essential fatty acids) and bulk. Thus malnutrition of a different type is actually widespread in the economically developed countries—malnutrition in the literal meaning of the word, *bad* nutrition.

Hence, the provision of a *well balanced optimal diet*—along lines demonstrated by recent broad advances in nutritional science—is the final goal, as yet unfulfilled for large sections of the population (66).

Chapter 3

Nutrition and Atherosclerosis (*Continued*)

D The Epidemiologic Findings

1 *Economically Underdeveloped Countries vs the United States—Habitual Diet and Atherogenesis*—Conditions of life in general—and patterns of diet and disease in particular—differ markedly in the economically underdeveloped countries and the United States (Fig 2) (40–63, 65–68) Most people in the underdeveloped countries usually consume a diet significantly lower than ours in total calories total protein and animal protein, total fat and saturated fat Unlike ourselves, they rely overwhelmingly upon grains and potatoes—relatively inexpensive high calorie high bulk foods—for their subsistence Hence, their fare is also low in cholesterol

The dietary differences between the economically underdeveloped countries and the United States are multiple and complex Moreover, there are many other areas of differences—racial, ethnic climatic geographic socioeconomic cultural public health and medical It is essential to be concerned with all these variables and their possible bearing upon observed patterns of disease

Clinical pathologic surveys on prevalence of atherosclerosis in the peoples of the economically less developed countries are extensive and consistent (69–70) Clinical atherosclerotic disease (particularly coronary heart disease) and severe morphologic atherosclerosis with myocardial infarction at postmortem examination are rarities at all ages

The focus on coronary disease has a twofold basis First the most extensive data are available on this entity Second this clinical disease is a resultant of the pathologic process of atherosclerosis in a very high percentage (over 90%) of cases (290–292 294–299) Therefore, the prevalence and incidence rates of coronary heart disease apparently are a close reflection of the occurrence of severe coronary atherosclerosis in a given population This correlation does not necessarily hold to a high degree for disease at other sites With clinical cerebrovascular disease for example it is difficult to distinguish among cerebral embolism, cerebral hemorrhage and cerebral thrombosis These are processes having different etio-pathologic bases Thus recent pathologic studies indicate that cerebral hemorrhage in hypertensives may be a complication—not

chiefly of atherosclerosis (as has been widely assumed)—but rather of a hypertensive military aneurysmal lesion in the smaller cerebral vessels. Unlike with coronary disease therefore, it is probably not correct to assume that prevalence and incidence rates for cerebrovascular disease reflect the occurrence of severe cerebral atherosclerosis in a given population (21). This invalid assumption has tended to confuse analysis of international epidemiologic trends (139). For valid comparisons cerebral thrombosis (usually atherosclerotic in its pathologic basis) would have to be separated out from the heterogeneous cerebrovascular disease grouping.

As long ago as 1934, the relative freedom from atherogenesis was clearly correlated with life span dietary patterns. The following generalization was formulated based on an analysis of 28 papers in the literature (69):

“ In no race for which a high cholesterol intake (in the form of eggs, butter and milk) and fat intake are recorded is atherosclerosis absent. Where a high protein diet is consumed which naturally contains small quantities of cholesterol, but where the neutral fat intake is low, atherosclerosis is not prevalent.”

This deduction was based on an analysis of reported clinical and pathological surveys made in China, East Africa, Egypt, India, and Malaya, compared with studies in Austria, Germany, the United States, and elsewhere.

2 *Economically Underdeveloped Countries—Social Class Differences in Diet and Atherogenesis*—This study yielded another highly relevant finding. Data from Egypt and India indicated that within these economically less developed countries significant social class differences were demonstrable in prevalence of atherosclerosis. Thus in both the countries atherosclerosis was apparently rare in the poorer classes subsisting largely on cereals, whereas it was relatively frequent in the more well-to-do classes enjoying a diet high in saturated fat and cholesterol (69). These and other findings indicated that racial, ethnic and climatic influences could not be primarily responsible for the observed differences in atherogenesis.

During the last two decades surveys in Ceylon, Costa Rica, East Africa, Guatemala, India, Indonesia, Iraq, Israel, Malaya, Mexico, Nigeria, Okinawa, South Africa, Uganda, all reinforce the same basic conclusion. Significant atherosclerosis is rare in peoples whose diet over the life span is predominantly vegetarian and low in calories, total lipids, saturated lipids and cholesterol (1, 4, 5, 12, 13, 21, 23, 68–102). It is rare clinically and morphologically in both males and females—in marked contrast to the United States where among middle aged males particularly atherosclerotic coronary heart disease is epidemic and a high percentage manifests severe gross lesions in the coronary arteries (2, 23, 290–292, 294–300).

3 *Economically Underdeveloped Countries vs. the United States—Cor*

relations among Diet, Serum Cholesterol Lipids Lipoproteins and Atherogenesis—Recent studies have also demonstrated that the correlation may be extended from a twofold to a threefold one—among diet, plasma cholesterol lipid lipoprotein levels and atherogenesis. Thus the significantly lower prevalence and incidence of atherosclerotic disease in economically less developed countries correlate also with differences (apparently diet induced) in levels of plasma cholesterol lipids lipoproteins (Figs 3 to 5) (1 4-6, 12-14 22, 23, 67, 71, 72 77 81, 91-99, 103-109)

FAT INTAKE, SERUM CHOLESTEROL AND FATAL CORONARY HEART DISEASE—AMERICANS VS BANTUS




FIG 3 Walker A R P and Arvidsson U B (92) Higginson J and Pepler W J (93) Note the differences in fat intake serum cholesterol level and percent of all autopsied deaths attributed to coronary heart disease. Note also the marked tendency for the serum cholesterol of Americans to rise from age 30 to age 45—a tendency virtually absent in the South African Bantu. Note also the sex differential in the occurrence of coronary disease in the Americans and its absence in the Bantus (cf text and ref 23)

Groups of clinically normal people in these economically less developed nations ingesting a predominantly vegetarian diet low in calories lipids-cholesterol * have mean plasma cholesterol lipid lipoprotein levels signi

*Throughout this book this characterization—diet low or high in calories-lipids-cholesterol is used as a convenient abbreviation for the longer term total calories empty calories total fats saturated fats cholesterol

chiefly of atherosclerosis (as has been widely assumed)—but rather of a hypertensive military aneurysmal lesion in the smaller cerebral vessels. Unlike with coronary disease, therefore it is probably not correct to assume that prevalence and incidence rates for cerebrovascular disease reflect the occurrence of severe cerebral atherosclerosis in a given population (21). This invalid assumption has tended to confuse analysis of international epidemiologic trends (139). For valid comparisons, cerebral thrombosis (usually atherosclerotic in its pathologic basis) would have to be separated out from the heterogeneous cerebrovascular disease grouping.

As long ago as 1934 the relative freedom from atherogenesis was clearly correlated with life span dietary patterns. The following generalization was formulated based on an analysis of 28 papers in the literature (69):

' In no race for which a high cholesterol intake (in the form of eggs, butter and milk) and fat intake are recorded is atherosclerosis absent. Where a high protein diet is consumed which naturally contains small quantities of cholesterol but where the neutral fat intake is low atherosclerosis is not prevalent.'

This deduction was based on an analysis of reported clinical and pathological surveys made in China, East Africa, Egypt, India, and Malaya compared with studies in Austria, Germany, the United States, and elsewhere.

2 *Economically Underdeveloped Countries—Social Class Differences in Diet and Atherogenesis*—This study yielded another highly relevant finding. Data from Egypt and India indicated that within these economically less developed countries significant social class differences were demonstrable in prevalence of atherosclerosis. Thus in both these countries atherosclerosis was apparently rare in the poorer classes subsisting largely on cereals, whereas it was relatively frequent in the more well to do classes enjoying a diet high in saturated fat and cholesterol (69). These and other findings indicated that racial, ethnic and climatic influences could not be primarily responsible for the observed differences in atherogenesis.

During the last two decades surveys in Ceylon, Costa Rica, East Africa, Guatemala, India, Indonesia, Iraq, Israel, Malaya, Mexico, Nigeria, Okinawa, South Africa, Uganda, all reinforce the same basic conclusion. Significant atherosclerosis is rare in peoples whose diet over the life span is predominantly vegetarian and low in calories, total lipids, saturated lipids and cholesterol (1, 4, 5, 12, 13, 21–23, 68, 102). It is rare clinically and morphologically in both males and females—in marked contrast to the United States where among middle aged males particularly atherosclerotic coronary heart disease is epidemic and a high percentage manifests severe gross lesions in the coronary arteries (2, 23, 290–292, 294, 300).

3 *Economically Underdeveloped Countries vs the United States—Cor*

factors (1, 5, 22, 23, 71, 72, 91-100, 103-109, 111-115) (Guatemala data are lacking on upper class patterns of disease)

These findings cast serious doubt upon the validity of the standards commonly accepted in the United States for normal plasma cholesterol concentration. They pose the questions: Is the life span pattern of diet inducing a chronic low grade hypercholesterolemia in millions of Americans, leading to widespread atherogenesis? What is an optimal serum cholesterol level for optimal freedom from atherosclerotic disease over an optimal life span?

Diet Serum Cholesterol and Atherosclerosis in Middle Aged Guatemalans and North American Males




FIG 5 Data from Scrimshaw N S Trulsson M Tejada C Hegsted D M and Stare F J (108) Mann G V Munoz J A and Scrimshaw N S (109) Tejada C and Gore I (100). Caloric intake is calories per person per day. Relative weight is an index: it is the ratio ($\times 100$) of an individual's observed weight to the median weight for his sex height group. Protein intake is expressed in relation to the level of daily intake recommended by the U S National Research Council. In the fat intake columns the upper values are for vegetable fat, the lower for animal fat.

Two approaches are identifiable to standard of normalcy. The first—and usual—involves measurement of a given index *e.g.* serum cholesterol concentration, in an adequate sample of individuals who at the given moment are apparently normal clinically. The values obtained are then accepted as indicative of the mean and range of normal. A second—and in our opinion more valid—approach involves consideration not only of the momentary but also of the long term medical significance of a given value. In arriving at values for normal cholesterol, for example, it asks the question: What values (mean and range) are consonant with optimal

significantly lower than groups of clinically normal persons in the United States. Moreover, they exhibit a different pattern of change in plasma cholesterol level with age. Thus, in middle class Americans plasma cholesterol levels tend to rise with each decade of adult life. This pattern is absent in the South African Bantu, but present in the economically more privileged South African Europeans, whose diet (like that of the Americans) is "rich" in calories-lipids-cholesterol (Figs 3 and 4). It is absent in manual laborers in Guatemala and India (largely vegetarians), but present in better off physicians, business men, officials, army officers ingesting a high calorie, high fat, high saturated fat diet (Fig 5). The several population groups in India generally exhibit corresponding differences in incidence of coronary heart disease—again pointing to the inconsequential role of ethnic and climatic, in contrast to nutritional




FIG 4 Brock J F and Bronte Stewart B (96) Keys A Anderson J T Aresu M Björck G Brock J F Bronte Stewart B Fidanza F Keys M H Malmros H Poppi A Postelli T Swahn B and del Vecchio A (159) Relative average fat intake (2nd graph) is the ratio ($\times 100$) of the fat intake of the given group to the mean fat intake for all 9 groups. Data on fat intake were obtained by semiquantitative weekly dietary surveys which assumed that each individual consumed an average helping. By this method, fat intakes were apparently estimated in grams per person per day and apparently converted into fat units (3rd graph) by dividing by 3.15. It was estimated that the Bantu consumed 17 per cent of total calories in the form of fats, the Coloured 25 per cent and the Europeans 35 per cent. The greater total fat intakes with greater incomes were due to greater intakes of animal (chiefly saturated) fat, not vegetable fat. Note the correlation between cholesterolemia and race, income, total fat intake, animal fat intake. Note that for any given ethnic group, cholesterolemia correlates with income, total fat intake, animal fat intake—i.e., race *per se* is apparently not a decisive influence. Note the lack of correlation between cholesterolemia and grade of physical work.

The Bantu group is a predominantly Negro group modified historically by Hamitic (Southern Arabian) influence. The Bantu migrated from central equatorial to southern Africa. The term Bantu is mainly linguistic in significance. The Cape Coloured group is a mixture of the original Hottentot inhabitants of southern Africa, Indonesians, Malaysians and whites. The European group is made up of whites of European stock born and reared in the Union of South Africa (96).

Serum Cholesterol in several countries and percent of calories from fat




FIG 7 Keys A (122)

Diet & Cholesterolemia in 284 Clinically Healthy Japanese Men Age 40-49 in seven groups




FIG 8 Keys A (107) Note the similar low levels of serum cholesterol in Japanese farmers miners mechanics clerks—values correlating with habitual diets low in fats and not with occupation and physical activity of work. Note the higher serum cholesterol concentrations in Japanese physicians and in Japanese Hawaiians and Japanese Americans—again correlating with dietary patterns

freedom from atherosclerotic disease over an optimal life span? Newer research knowledge has brought us to the point where it seems essential to re define standards of normalcy for serum cholesterol, based on this latter approach

4 *Italy and Japan vs the United States*—Significant findings concerning the relationship between diet and atherosclerosis have been forthcoming recently also from countries intermediate in the scale of economic development, e g, Italy and Japan. Thus, vital statistics data reveal that these countries manifest markedly lower heart disease mortality rates than our own, due principally to the relative rarity of coronary heart

DIET IN THE UNITED STATES COMPARED WITH ITALY AND JAPAN-1950




FIG 6 Stamler J (22 23) (for sources of data cf refs 41 and 42)

disease. At the same time the people of Italy and Japan *en masse* consume a diet much lower than our own in total calories, empty calories, total fats, saturated fats and cholesterol (Fig 6) (5, 6, 22, 23, 40-42, 61, 62, 65, 68, 71, 72, 81, 91, 99, 105-107, 116-121, 123).

Further recent studies of the clinical pathologic survey type indicate that the threefold correlation—among diet, plasma cholesterol, lipid lipoprotein levels and atherogenesis—is also demonstrable for large sectors of the populations of Japan and Italy (and Spain as well) (Figs 7 to 9) (5, 6, 22, 23, 71, 72, 91, 99, 105-107, 118-121, 123, 124). Here again plasma cholesterol levels fail to increase grossly with age. Moreover, as in South Africa, Guatemala, Egypt and India, more well-to-do Italians, Japanese, Spaniards—consuming diets different from those of their poorer fellow citizens but similar to our own—have patterns of plasma lipids,

Death rates in Italian and Japanese migrants to U.S. compared with
Italians in Italy and Japanese in Japan
Males, age 45-54 1950




FIG 10 Stamler J (22 23) Stamler J Kjelsberg M Hall Y and Scotch N (126)
Note that the Italian Americans have higher death rates than the Italians in Italy
The difference is conspicuous for cardiovascular renal diseases heart diseases arterio-
sclerotic heart diseases The same phenomenon obtains for the Japanese Americans
compared with the Japanese in Japan Note also that for these three cause-of-death
categories the Italian Americans and the Japanese Americans have lower mortality
rates than all Americans Finally note the different epidemiologic patterns for mortality
due to vascular lesions of the central nervous system—patterns probably reflecting
chiefly hypertensive (not atherosclerotic) vascular disease in this age group (21 23
196 230)

Occurrence of Myocardial Infarction in Yemenite
Jews compared with other Jews in Israel




FIG 11 Dreyfuss F (132) Dreyfuss F Toor M Agmon J and Zlotnick A (128)
Note that the Yemenite Jews constituted 25 per cent of the Jewish population of Israel
and 22 per cent of all Jewish hospital admissions but only about 2 per cent of all Jewish
admissions for myocardial infarction Contrast this with the Ashkenazi (Eastern and
Central European) Jews (Sefardic Jews are of Spanish origin)

and morbidity and mortality due to coronary heart disease, approaching those prevailing in the United States

These findings again emphasize that ethnic and racial factors are not primarily responsible for the international epidemiologic pattern of coronary heart disease. Moreover, they decisively reinforce the general thesis that life span pattern of diet is a critical factor in the causation of the variable incidence of atherosclerosis among different peoples

Incidence of high grade Coronary Sclerosis in consecutive autopsies US vs Japan




FIG 9 Kimura N (116) White N K Edwards J I and Dry T J (793) Note the markedly higher incidence of severe coronary sclerosis in American males at all ages. Note also the presence of a gross sex difference for Americans and its virtual absence for the Japanese (cf Fig 3)

5 *Effects of Emigration—Yemenite Jews, Italians and Japanese*—This conclusion is fortified by recent work showing that changes in the mode of life particularly the diet consequent upon emigration are associated with increases in plasma cholesterol lipid lipoprotein levels and in coronary heart disease incidence rates. This has been indicated in studies on Yemenite Jews in Israel on Italians in the United States and on Japanese residing for many years in Hawaii or the United States (Figs 10 and 11) (6, 22, 23, 71, 72, 81, 99, 107, 125-138)

6 *The Low Countries the Scandinavian Countries et al, and the United States*—Significant data are also available on European populations usually ingesting diets higher than most Italians in total calories total

fluence of relatively high intakes of unsaturated marine oils on cholesterolemia and atherogenesis in the populations of these countries (167)

A negative correlation was noted between intake of vegetable lipid and death rates attributed to arteriosclerotic heart disease (or arteriosclerotic plus degenerative heart disease). This finding is particularly intriguing in view of recent data demonstrating low serum cholesterol levels in American vegetarians and in Yugoslavs subsisting on diets high in vegetable oils (67, 251). It is also of interest in connection with observations on the reduction of cholesterolemia by incorporation of unsaturated vegetable oils in American type diets.

The cited study also noted a positive correlation between protein intake and arteriosclerotic heart disease death rates (139). The question therefore becomes: Does the statistical correlation between death rates and levels of total protein/animal protein consumption reflect a *cause-and-effect* relationship between protein intake and atherosclerotic disease? These data pose this question but cannot answer it. The answer must be sought from other available evidence.

Extensive experimental data indicate that this is *not* the case (1, 451). On the contrary, they suggest that high protein intake may afford a partial protection against the hypercholesterolemic and atherogenic effects of high fat/high cholesterol diets (454, 455, 458, 482, 504-516). On the other hand, extensive clinico-pathologic and animal-experimental data indicate that the correlation between death rates and intakes of total calories, total fats/saturated (animal) fats is significant cause-and-effect-wise.

A considerable spread was noted in the data on diet and death rates as analyzed in this study (139). For several economically developed countries with total fat intakes in the range 30 to 40 per cent of total calories, age-specific death rates varied from 500 to 600 or more per 100,000 population (e.g., Austria, West Germany, Sweden, Norway, Denmark, the Netherlands, Australia, Canada, Finland). A few of these countries manifested middle-aged death rates considerably lower than those for the United States (739 per 100,000).

Based on findings of this nature, several recent reports critically reviewed the problem of the relationship between intake of lipids (total and saturated) and occurrence of atherosclerotic coronary disease (139-143). These critiques have in turn been recently subjected to penetrating analysis (101). In evaluating these critiques, it is appropriate to note that the broad statistical data—despite their recognized limitations—*verified* significant overall correlations between death rates and intakes of total calories, total fats and saturated fats (Fig. 12) (139). These findings therefore did *not* refute the validity of the correlation. A valid conclusion—considering the scatter in the data (as well as other evidence)—is that the correlation among diet, serum cholesterol and coronary disease is not reflective of a one-to-one cause and effect relationship.

fats, saturated fats and cholesterol (Figs 7 and 12) (1, 5, 6, 22, 23, 25, 40-42, 65, 67-72, 90, 91, 97, 107, 139-160). Thus, serum cholesterol levels were found to be consistently higher in Swedish population groups compared with Italians (Fig 7). The former had values almost as high as those typically recorded in the United States. Serum cholesterol levels in middle aged males (firemen and Rotarians) in Helsinki, Finland were 260 mg %, exceeding those for the United States (107). By 1950, the

Rank correlation coefficients (r) between various dietary components and death rates for 22 countries -- Males age 55-59

Dietary Component	Atherosclerotic and Degenerative Heart Disease	Atherosclerotic and Degenerative Heart Disease and the Diseases of the Heart
total calories	0.723	0.593
calories from fat	0.659	0.470
calories from animal fat	0.684	0.562
calories from vegetable fat*	-0.236	0.282
calories from protein	0.709	0.694
calories from animal protein	0.756	0.695
calories from vegetable protein	-0.430	-0.153
calories from carbohydrate	0.305	0.423
% calories from fat	0.587	0.390
% calories from animal fat	0.677	0.557
% calories from vegetable fat*	-0.468	0.509
% calories from protein	0.172	0.465
% calories from animal protein	0.643	0.608
% calories from vegetable protein	-0.651	0.483
% calories from carbohydrate	-0.562	-0.386

* Number of countries (N) 21 (data not available for France) otherwise N=22

Critical Values of r		
N	$\alpha = 0.05$	$\alpha = 0.02$
21	± 0.438	± 0.521
22	± 0.428	± 0.508

FIG 12 Yerushalmy J and Hilleboe H E (139)

effects of World War II had receded and prevalence and incidence of atherosclerotic coronary heart disease were again sizable in the Scandinavian and Low Countries (1, 5, 68, 72, 107, 139-142, 157-166).

Calculation of rank coefficients of correlation for 1950 data for 22 countries demonstrated statistically significant correlations between death rates and intakes of total calories, total fats and animal fats (Fig 12) (139). One of the problems needing further investigation is the in

In any attempt to arrive at a tentative judgment in this matter, the following facts stand out. Cardiovascular renal death rates for middle-aged white males have been high throughout the last 30 to 40 years. However, the rate did not stay constant during this period in males but rose noticeably. In contrast, middle aged white female rates declined markedly (Fig 13) (184). Problems inherent in the available statistics render difficult any evaluation of the trend of death rates over the last 30 to 40 years for arteriosclerotic heart disease *per se*. Hence, an inter

Trend of Cardiovascular-Renal Mortality in middle aged Americans by sex and race—1920-1955




FIG 13 Moriyma I Woolsey T and Stamler J (184). The trends for middle aged whites are discussed in the text. Note that Negro rates are higher than white throughout the 35 year period due principally to the several fold higher mortality rates from hypertensive and related diseases in Negroes. Note also that the sex differential in mortality rates for Negroes is slight compared with whites due chiefly to the high death rates in Negro females (*cf* Fig 1). This pattern is different than in the economically less developed countries where the low order of the sex differential is due chiefly to low male death rates (*cf* Figs 3 and 9).

pretation must be made of the data for a broad grouping of several diseases—the cardiovascular renal diseases (172, 173, 180, 183, 184, 195). The question therefore arises: if the rise in white male rates is chiefly an apparent one due merely to changes over the years in diagnostic acumen and death certification procedures (as some investigators have suggested)—why then have these changes produced a rising trend only for males?

As is well known, atherosclerotic coronary heart disease is common in middle aged American male whereas it is remarkably rare in premenopausal females (Fig 1) (1-5, 12-23, 71, 90, 134-137, 143, 155, 172-187,

Further, a number of the countries—particularly the Scandinavian and Low Countries—were in 1950 only a few years away from the privations of World War II, with its significant restrictions in diet and concomitant declines in arteriosclerotic heart disease death rates (1, 5, 6, 23, 71, 72, 101, 143-146, 155, 160-165, 168, 169). It is highly possible that these effects of World War II still found expression in 1950.

Moreover, the observed patterns of mortality for these European countries and the United States ranged *in toto* on the *high* side of a distribution curve for middle aged population groups—particularly in comparison with the economically underdeveloped countries. Irrespective of the observed differences in rates, therefore, coronary heart disease is unquestionably a major public health problem in these European countries, in contrast to the underdeveloped nations.

Finally these differences indicate the need for further detailed field studies of the hospital clinical pathological survey type on diet, serum cholesterol and coronary disease. The several such investigations accomplished in recent years have in fact yielded impressive data supporting the nutritional metabolic cholesterol lipid lipoprotein theory of atherogenesis (5, 6, 71, 107, 157-159).

7 *World Wars I and II*—This theory is also amply supported by the findings on time trends of diet and disease in the Scandinavian and Low Countries during World War II. This subject was reviewed previously (1), and is therefore not presented *in extenso* here. The marked reduction in intake of total calories, total fats, saturated fats and cholesterol imposed by the German occupation was associated with a gross decline in arteriosclerotic heart disease mortality rates (1, 5, 6, 23, 71, 72, 143-146, 155, 160-165). In view of recent clinical research findings on dietary lipids and blood coagulability, it is noteworthy that the occurrence of thrombo embolic episodes also declined in Scandinavia during the war years. Hence it has been suggested that the decrease in coronary heart disease rate reflected a diet induced decline in both atherogenesis and thrombogenesis (161). Similar findings were reported from Leningrad during the World War II siege and from Central Europe during the famine years following World War I (10, 168, 170).

8 *The United States—1910 to 1955*—This whole problem of the trend of atherosclerotic disease over the years—and its relationship, if any to evolving dietary patterns—has recently received the attention of several investigators. American and British data suggest a marked increase in middle age mortality rates in these two countries in recent decades (Fig 13) (5, 22, 23, 143-146, 155, 171-190). The evaluation of these trends however is complicated by a host of statistical and methodological problems—some of them permanently beyond precise quantification because of the nature of the data (23, 139, 140, 142-146, 172-174, 191-194). Hence it is not surprising that disagreements arise concerning the actual degree and extent of the apparent increases in coronary disease mortality (23, 139, 140, 142-146, 172-174, 178, 180, 183, 184, 195).

and has remained so—much higher even than in most other economically developed countries (Fig 14) (210–212). Another unique American dietary characteristic—the large intake of butter, lard and other predominantly saturated fats—has apparently been further accentuated, partly as a result of the introduction in 1910 of the catalytic hydrogenation process for converting vegetable oils into plastic shortenings and margarines. As a consequence of the hydrogenation process, which saturates double bonds, these fats have a high percentage of *trans* fatty acid isomers

CONSUMPTION OF FOOD IN THE UNITED STATES -1909 & 1952




FIG 14 Stamler J (22, 23) (for source of the data cf ref 212). Data indicate foods available for consumption, not foods actually consumed. Unlike Figs 2 and 6 based on United Nations sources, the data on fats and oils from the U.S. Department of Agriculture include bacon and salt pork, hence the higher values. The data on fats and oils are subdivided into specific components; from above down they are: butter, lard, margarine, bacon and salt side, shortening, other edible oils. Values for eggs are number of eggs per person per year.

(23 to 42 per cent) and a low content of essential fatty acids (2 to 8 per cent) (217, 218). Unless enriched they are practically devoid of other essential nutrients, e.g. fat soluble vitamins, hence they are an 'excellent' source of empty calories.

Intakes of milk, dairy products, eggs have also exhibited significant increases. A marked rise in ingestion of refined sugars (100 per cent empty calories) is another important long term development in the American diet—from 8 pounds/person/year in 1820 to 30 in 1860, 66 in 1900 and 95 today (210–212). All these upward trends have been as-

196-206, 208, 209, 294, 461, 645, 646) This sex differential in susceptibility for middle aged whites is in the order of 5-25 to 1

The 1920-1955 trend of middle aged white male vs middle aged white female cardiovascular renal deaths in the United States therefore, reflects the following For males, a sizable rise in coronary disease leading to a net increase in cardiovascular renal disease mortality, despite declining death rates for infectious diseases involving the cardiovascular renal system For females, continued low coronary disease rates, and a marked decline in death rates due to puerperial and infectious diseases of the cardiovascular renal system resulting in a gross fall in overall cardiovascular renal death rates This tentative conclusion is borne out by other data Thus, while life expectancy at birth for Americans has risen by about 20 years since 1900 life expectancy at age 50 has increased only 2.2 years In view of the great advances during these decades in the prevention of death in middle age due to pneumonia and other infectious diseases, tuberculosis, luetic and rheumatic heart diseases, subacute bacterial endocarditis diseases amenable to surgical treatment, and others, a greater increase in life expectancy at age 50 might have been anticipated Almost certainly this absence of a marked improvement is attributable chiefly to an increasing mortality from coronary heart disease in middle age (167)

If this foregoing analysis be correct the qualitative judgment must be accepted that a true increase has actually occurred in mortality among middle aged white American males due to atherosclerotic coronary heart disease This conclusion is a widely but not unanimously held one (167 171-190 195)

In terms of the basic problem of the relationship among diet, atherosclerosis and atherosclerotic disease it therefore becomes essential to attempt an evaluation of the evolution of dietary patterns in the United States over the last several decades Here it is important to note that people aged 50 dying of coronary disease in 1950 lived their life span over the years 1900-1950 Those aged 50 dying in 1920 lived from 1870-1920 If life span patterns of diet exert a key influence in the etiology of atherosclerosis then any review of dietary patterns must seek out data on conditions prevailing in the third and fourth quarters of the nineteenth century as well as in the twentieth century

Here again serious methodological limitations exist due not only to a relative paucity of data but to the particular scarcity of information concerning diets in different regions of the country and among different racial, ethnic and socioeconomic strata at different times (cf refs 210-213 216) A further difficulty stems from the fact that much of the data on dietary patterns in Americans deals with foods available for consumption, rather than foods actually consumed (102 195) With these shortcomings in mind, it is possible to delineate the following broad trends Meat consumption has generally tended to be relatively high in the United States,

the need for a multifactorial analysis of atherosclerotic disease causation. Because of these complexities—and the limitations of both the mortality and the nutritional data, plus the insurmountable obstacles to their further delineation—extreme caution is essential in interpreting them. However, it is not inappropriate to note that the concepts of the nutritional metabolic cholesterol lipid lipoprotein theory are certainly not contradicted by these data.

9 *The United States—Morbidity and Mortality Findings in Different Population Groups and Geographic Areas*—It has come to be widely recognized that this country—with its large population stratified by region, occupation, income, ethnic and racial background—constitutes an excellent epidemiologic “laboratory.” Based on this approach, one of us (J S) initiated a long term investigation on the epidemiology of the major cardiovascular renal diseases in the city of Chicago and the state of Illinois (19, 21–23, 125, 126, 196, 197, 228–230).

First an analysis of 1951 and 1953 mortality data was undertaken for specific groups in the population stratified by age, sex, race, nationality, occupation, place of residence—with a particular focus on the middle decades of life (Figs. 1, 13 and 15). The following findings are of particular significance. The death rates for arteriosclerotic heart disease were several fold higher in middle aged white men than women (Fig. 1)—an observation fully in accord with extensive data from multiple sources (1–5, 12–23, 71, 90, 134–137, 143, 155, 172–187, 196–206, 208, 209, 294, 461, 645, 646). It is quite unlikely that diet is a major factor responsible for this gross sex differential, since diets of American males and females are probably not markedly disparate (22, 23, 102, 231). This is a problem meriting further study in view of the paucity of extant data and evidence suggesting that obesity is more prevalent in white men than women (254). This sex differential in coronary disease morbidity and mortality in a population habitually ingesting a diet high in calories-fats-cholesterol is either absent or much less marked in economically less developed countries—chiefly because rates for middle aged males are much lower in the latter countries (*cf* Figs. 3 and 9). This finding illustrates the interplay between diet and other factors in the etiology of atherosclerotic disease. When the diet is non atherogenic, coronary rates are low in both sexes. When the diet is atherogenic, the susceptible male sex manifests a high rate in middle age. The resistant female sex continues to exhibit a relatively low rate (although a rate somewhat higher than her counterparts in the economically less developed countries).

With respect to this problem, Negroes in the United States appear to present a unique pattern. The sex differential is apparently much less marked than for whites, chiefly because Negro female rates are relatively high (Figs. 1 and 13). Further work is needed to delineate the parameters and bases for this phenomenon (23, 196, 228–230).

These complex facts concerning the sex differential again serve to em

sociated with a steady decline in grain consumption, of such marked proportions that for tens of millions of Americans today bread can hardly be designated the "staff of life" (Figs 2, 6 and 14) (210-212). A qualitative metamorphosis in flour has also occurred over the decades, i.e., a decline in percentage utilization of whole meal. The resultant nutritional losses have been only partially compensated by the limited bread enrichment measures of recent years (224-226). The increased consumption of highly processed, refined foods has also involved a decline in intake of bulk—a decline only partially compensated by the increased ingestion of fruit and vegetables. Little work has been done on the possible implications of this trend for health and disease (*cf.* refs 92, 93, 227).

Because of the limited data on food actually consumed (as distinct from food available for consumption), and the related difficult problem of estimating waste of fats in cooking, some investigators doubt the validity of data indicating an increase in fat consumption by Americans during the last 40 to 50 years (102, 195). It is noteworthy that these writers are also skeptical about the data indicating an increase in coronary disease incidence in middle aged males over these years (195). Thus, these authors imply that both fat intake and coronary disease incidence have been high since the turn of the century. If these are actually the facts, and the two phenomena are so correlated, this certainly is not inconsistent with the nutritional metabolic theory.

In this connection, one other problem merits attention. Even if fat intake has increased since the turn of the century, it was—as already noted—high even then. If coronary disease incidence in middle aged males was much lower then, as some data suggest, why *this* lack of correlation? This possible disparity serves to re-emphasize that the relationship between diet and disease is not one-to-one; that multiple factors interact with diet to influence occurrence of coronary disease. With respect to the trends over the last 40 to 60 years, attention needs to be focussed on the interplay between diet and psychological stress, and particularly between diet and large muscle activity.

The slight decline in mean total calorie ingestion between 1910 and 1950 must be assessed in terms of another related phenomenon, the decline in physical activity. The problem of the possible effects of sedentary life on lipid metabolism and atherogenesis is discussed below.

Viewed overall, these trends are—within the limitations noted above—highly suggestive of a definite increase (relative and absolute) in consumption of total fats, saturated fats, cholesterol, refined carbohydrates and empty calories (210-212, 219-223).

These apparent changes with time in patterns of diet and disease may or may not be significantly interrelated cause and effect wise. Thus these decades witnessed other manifold changes in American life, e.g., increase in urbanization, automobile transportation, cigarette consumption, sedentary work, and other similar changes—phenomena emphasizing

tive epidemiologic studies—corroborate the basic finding that coronary heart disease is widely prevalent in middle aged American males of most population subgroups (2, 10 22 23 69, 126 172–188 196, 197 199 206, 208, 209, 232–237, 260–278 290–292 294 300)

Several studies—in different parts of the country over the last two decades or more—have yielded disparate results on the problem of the relative incidence of coronary disease in Negroes and whites (10, 23, 172–

OCCUPATION RACE AND ARTERIOSCLEROTIC HEART DISEASE DEATH RATES--CHICAGO 1951 MALES AGE 45-54




FIG 15 Stamler J (23) Stamler J, Kjelsberg M and Hall Y (196) It is likely that the death rates for the laborers are falsely high—probably due to inaccurate listings of occupation on death certificates. The data were therefore pooled into two broad groups—white collar workers (A + B) and blue collar (manual) workers (C + D + E).

180 196 199 206 208 228–230 260–271 274–277) In contrast it is generally agreed that hypertensive disease is several times more frequent in Negroes than whites (23 196 261)

Data from Norway indicate that a situation similar to the United States prevails for that country (160). The data concerning Great Britain are more complex. There a social class difference in arteriosclerotic heart disease mortality has been observed for several decades—a finding complicated by the fact that an opposite social class distribution has been recorded for so-called degenerative heart disease (150–153). Conspicuous in these data is the trend for coronary disease death rate to

phasize that no simple one to one relationship exists between diet and atherosclerotic disease. The full significance of this conclusion is discussed subsequently at length.

High arteriosclerotic heart disease death rates were recorded for the city of Chicago, for other Illinois municipalities and for rural Illinois (2, 23, 197). However, a certain differential was noted, i.e., relatively higher rates for Chicago, lower for rural Illinois. These observations are in accord with data from other states (175, 183, 232-235) (*cf* also refs 234-237). Here again, the possibility must be investigated of multiple causative factors operating to produce this apparent result, since dietary patterns in urban and rural areas, at least of the northern United States, are not markedly different (22, 23, 107, 197, 210-214, 219-223, 238-253). (Rural southern Negroes may be—or may have been—to a degree exceptional in this regard [238-239].) This fact suggests that the high calorie lipid cholesterol diet of most Americans, urban and rural, (22, 23, 107, 197, 210-214, 219-223, 238-253) as compared to less developed countries, does in fact result in extensive atherosclerotic disease, irrespective of other factors possibly operating to produce differential effects in urban vs. rural areas.

Analysis of the foreign born by nationality leads to a similar conclusion. Thus, death rates were high for all nationality groups—considerably higher than those for the given nationalities in their countries of origin (Fig. 10) (22, 23, 126) (*cf* refs 6, 68, 71, 72, 81, 99, 102, 107, 116-121, 123-127, 132-137, 139-148, 150-160, 165). However, a moderate range in rates was recorded with the distribution tending to parallel that for the countries of origin. The foreign born nationalities therefore bear the stamp of their tenure in both their countries of birth and in the United States. This is probably true not only for the death rates but also for the dietary patterns of these first generation Americans. Their nutritional habits represent an amalgam of old country and American dietetics (22, 23, 107, 126, 133, 135, 216). One subgroup of the American Indian ethnic grouping—the Navajo—has recently been the subject of several reports differing in their conclusions (107, 256-259). The much discussed, but inadequately studied Eskimo was analyzed in detail previously (1, *cf* also ref 107). Therefore, without discounting the possible influence of other factors, these findings are consistent with the concept that diet plays a key (although not exclusive) role in the causation of atherosclerotic disease.

The analysis of Chicago mortality data also revealed that coronary heart disease death rates were high for middle aged males of both Negro and white races and of all occupation income groups (Fig. 15) (19, 22, 23, 125, 196, 228-230) (*cf* refs 10, 69, 172-180, 183-188, 199, 206, 208, 209, 232, 233, 260-273). The toll from atherosclerotic disease was at least as great in the poorer strata as in the more well to do. Considerable data from other sources—e.g., life insurance statistics and findings of prospec-

lower economic groups in the United States have—at least since the end of the depression in 1939-40—been consuming sizable amounts of meat, milk, eggs, fats and oils, white flour products, refined sugar, salt. Their intakes of total calories and empty calories saturated fats (animal and vegetable), cholesterol tend to be virtually as high as those of upper income Americans (Fig 16) (22 23, 107, 196 197, 210-214 216, 219-223, 238-253) (As already indicated, rural Southern Negroes may

Serum Cholesterol levels of clinically healthy American males age 40-59 of different social classes in different sections of the country




FIG 17 Lewis L A Olmsted F Page I H Lawry E Y Mann G V Stare F J Hanig M J aufer M A Gordon T and Moore F E (111) The subjects are almost exclusively white. Note the similar high mean serum cholesterol levels in different groups of middle aged American males irrespective of occupation and geographic region. The single exception is the group of Federal penitentiary prisoners with a somewhat lower mean value. Dietary data on this group would be of considerable interest. The particularly high values for Californians also remain to be accounted for.

be—or may have been—exceptional in this regard [23, 238-239]). Thus most social strata in our country ingest an habitual diet that is potentially atherogenic. Therefore the finding that coronary heart disease is prevalent among most sections of the American population does not contradict, but rather lends additional strong support to the theory of a decisive interrelationship between life span pattern of diet and atherogenesis.

rise over the recent decades while the social class differential narrowed (144-155). It would seem that Great Britain may be experiencing an increase in the incidence of this disease, embracing all sectors of its population. Despite this increase, death rates in Great Britain were in 1950-1954 still considerably below those for age sex race matched groups in the United States. Nonetheless, in Great Britain—as in Norway and the United States—this disease has apparently become widespread in the middle aged male population of all social classes.

FOOD CONSUMPTION OF LOW AND HIGH INCOME URBAN FAMILIES U S A -1948




FIG 16 Stamler J (22-23) (for source of data cf ref 213 see also ref 214) (cf legend Fig 14). These data represent foods available for consumption not foods actually consumed. Data for 1955 reveal essentially similar findings among different income groups in various sections of the country (214).

These findings forcefully pose an important question. Why is it that the poor in countries like Egypt, Guatemala, Japan, India, Italy, Spain have low coronary disease death rates whereas Americans on the lower rungs of the United States economic ladder have rates at least as high as more well-to-do groups? Are these apparently contradictory trends consistent with the basic thesis of a key relationship between diet and atherosclerosis? Our present answer is: By no means—and for good reasons of fact, namely the important differences that prevail in the dietary patterns of lower income Americans and the poorer classes of all the aforementioned countries (Fig 16 cf Figs 2 and 8).

In contrast to the populations *en masse* of these other countries, the

imbalance hypothesis, suggested a few years ago by one of us (J S) (16) for contemporary Americans, particularly lower income Americans, the word malnutrition is applicable in its literal meaning *bad nutrition*

Virtually no evidence has been presented contradicting the thesis that patterns of cholesterolemia in the United States population are a result of life span habitual diets. Nor have any other hypotheses been advanced to account for these patterns of cholesterolemia

11 *The United States—Findings in Prospective Studies on the Living Population*—The basic concepts of the nutritional metabolic theory received further verification in the recently published findings of the Framingham, Albany and Los Angeles epidemiologic studies (2 209, 272, 278) (*cf* also 112, 281). In general, the results of all of these investigations point in the same direction. When sizable groups of middle aged American males—originally free of clinical coronary heart disease—were followed prospectively the risk of developing this disease was definitely found to be proportional to levels of body weight, blood pressure and cholesterolemia (Fig 18) (*cf* foregoing refs to so called retrospective studies dealing with the effects of these variables. *cf* also refs 283–285 301–318 351, 431–448). Thus from the Framingham report, in the population at risk, normal with respect to all three of these variables the 4 year incidence rate was only 10 per thousand. This is one fourth the rate for American middle aged males in general (40/1 000/4 years). In contrast in the population at risk with two or three of these variables abnormally elevated the 4 year incidence rate was 143 per thousand—a more than tenfold increase in risk compared with the Framingham normals. These data are not only of great theoretical and research significance they also have profound implications for the practice of medicine (see below)

The Framingham data also serve to re-emphasize the important role of hypertensive disease as a factor in the development of clinical coronary disease during middle age particularly in populations with the prerequisite cholesterol lipid lipoprotein metabolic derangements

At the same time it is also essential to recognize that the incidence of coronary disease was by no means insignificant in the nonhypertensive middle aged American males followed in these prospective studies. The etiology of coronary disease in normotensive persons therefore remains a key problem

Similarly with respect to obesity. Thus while the overweight group in Framingham had a coronary disease rate approximately twice that of the normal weight group the rate in the latter (28/1 000/4 years) was itself a high one

In this regard available data on obesity and cholesterolemia are worthy of attention. In at least two studies a positive correlation has been demonstrated between serum cholesterol levels and body weight but this is a *low order* correlation (112 283). Put somewhat differently,

10 *The United States—Serum Cholesterol Findings in Different Population Groups and Geographical Areas*—This concept is further reinforced by abundant data indicating that serum cholesterol levels tend to be high in all strata of the middle aged United States population compared with values observed in the people of many other countries (Figs 3, 4 5, 7, 8, 17, 18, 19, 20 and 21) (1, 2, 6, 12-14, 22, 23 71, 72, 92-99, 103-109, 111, 112, 118-121, 123, 124, 127, 130, 133-137, 156-159 195, 209, 251, 272, 278-281 645) Such findings pose anew the problem of valid standards for normal—and optimal—serum cholesterol The evidence—epidemiologic, clinical and animal experimental—indicates that values less than 200 or even 175 mg % may afford optimal freedom from atherosclerosis and atherosclerotic disease and may therefore be designated normal (in the most profound sense of the term) The animal studies show that in virtually all other species analyzed serum cholesterol levels normally are in a range considerably below that observed in healthy people in the United States (1)

From the standpoint of the nutritional metabolic cholesterol lipid lipoprotein theory of atherogenesis these serum cholesterol findings were the anticipated predicted ones They were expected based on the two foregoing observations—i.e. the finding of diets high in total calories, total fats saturated fats, cholesterol in most strata of the United States population and the finding of high death rates from coronary disease in these strata

What is responsible for the distribution of serum cholesterol levels in the middle aged American male population and particularly the high mean values, compared with those of other peoples? Extensive data support the concept that this is critically a by product of habitual life span dietary patterns

In this connection one other aspect of contemporary nutrition in the United States may be of considerable importance While many lower income Americans consume a diet about as rich as that of their upper income compatriots in total calories and empty calories total fats and saturated fats cholesterol a larger percentage of them ingest suboptimal amounts of certain essential nutrients (vitamins minerals essential amino acids and essential fatty acids) (Fig 16) (22 23 107 191 210-216, 219-223 227 238-253 282)

It is therefore misleading one sided inaccurate and oversimplified to speak of Americans as eating 'too well' Their diets tend simultaneously to deviate in two directions, away from standards for optimal nutrition towards *overnutrition* i.e. excess of total calories and empty calories total fats and saturated fats refined carbohydrate cholesterol salt and towards *undernutrition* i.e. inadequacy (relative and/or absolute primary and/or secondary—see below) in essential nutrients and bulk Therefore it is most accurate and scientific to characterize the dietary problem in the United States today as one of dietary imbalance According to this

imbalance hypothesis suggested a few years ago by one of us (J S) (16) for contemporary Americans particularly lower income Americans the word malnutrition is applicable in its literal meaning *bad nutrition*

Virtually no evidence has been presented contradicting the thesis that patterns of cholesterolemia in the United States population are a resultant of life span habitual diets Nor have any other hypotheses been advanced to account for these patterns of cholesterolemia

11 *The United States—Findings in Prospective Studies on the Living Population*—The basic concepts of the nutritional metabolic theory received further verification in the recently published findings of the Framingham Albany and Los Angeles epidemiologic studies (2 209, 272 278) (*cf* also 112 281) In general the results of all of these investigations point in the same direction When sizable groups of middle aged American males—originally free of clinical coronary heart disease—were followed prospectively the risk of developing this disease was definitely found to be proportional to levels of body weight blood pressure and cholesterolemia (Fig 18) (*cf* foregoing refs to so-called retrospective studies dealing with the effects of these variables *cf* also refs 283–285 301–318 351, 431–448) Thus from the Framingham report in the population at risk normal with respect to all three of these variables the 4 year incidence rate was only 10 per thousand This is one fourth the rate for American middle aged males in general (40/1 000/4 years) In contrast in the population at risk with two or three of these variables abnormally elevated the 4 year incidence rate was 143 per thousand—a more than tenfold increase in risk compared with the Framingham normals These data are not only of great theoretical and research significance they also have profound implications for the practice of medicine (see below)

The Framingham data also serve to re-emphasize the important role of hypertensive disease as a factor in the development of clinical coronary disease during middle age particularly in populations with the prerequisite cholesterol lipid lipoprotein metabolic derangements

At the same time it is also essential to recognize that the incidence of coronary disease was by no means insignificant in the nonhypertensive middle aged American males followed in these prospective studies The etiology of coronary disease in normotensive persons therefore remains a key problem

Similarly with respect to obesity Thus while the overweight group in Framingham had a coronary disease rate approximately twice that of the normal weight group the rate in the latter (28/1 000/4 years) was itself a high one

In this regard available data on obesity and cholesterolemia are worthy of attention In at least two studies a positive correlation has been demonstrated between serum cholesterol levels and body weight but this is a *low order* correlation (112 283) Put somewhat differently

the middle aged American population ingesting its habitual diet tends generally to be at least slightly obese and to have high values for serum cholesterol, with or without concomitant gross obesity. When this high fat, high cholesterol diet is ingested in excess, with resultant development of gross obesity, serum cholesterol levels tend to be slightly—but not

EFFECTS OF OBESITY, HYPERCHOLESTEROLEMIA AND HYPERTENSION ON FOUR-YEAR INCIDENCE OF ARTERIOSCLEROTIC HEART DISEASE IN MEN AGED 45-62




FIG 18 Dawber T R Moore F E and Mann G V (209) Normal blood pressure—consistently below 140/90 elevated blood pressure—consistently 160 or over systolic and/or 95 or over diastolic Normal weight—relative weight under 100 elevated weight—relative weight 113 or over Normal cholesterol—under 225 mg % high cholesterol—260 mg % or above

markedly—higher. Given this nutritional metabolic situation it is to be expected that incidence of coronary disease would be substantial in non grossly overweight middle aged American males, and even greater (but not several fold greater) in the obese. (Life insurance data indicate that healthy thin persons have a significantly lower risk of coronary disease [284, 285]. No systematic serum lipid data are available on such persons.)

In brief, then, non hypertensive, non overweight middle aged American males—better off though they may be—nonetheless still have a by no means insignificant incidence of elevated serum cholesterol levels and coronary disease. The relationship of life-span diet to these phenomena cannot be gainsaid.

Chapter 4

Nutrition and Atherosclerosis (*Continued*)

E The Clinical Pathological Findings

EXTENSIVE data from clinical pathological research also lend strong support to the dietary metabolic cholesterol lipid lipoprotein theory of atherogenesis

1 *Historical Development*—Before proceeding to recent work it is valuable to review the long term contribution of clinical pathological research in clarifying the interrelationship between diet and atherogenesis. For the fact is that historically the initial posing of the problem came primarily as an outgrowth of pathologic studies. In the middle of the last century, cellular pathology identified atherosclerosis as a separate specific lesion among the arterioscleroses—distinguishable particularly by its large content of cholesterol and other lipids. Laterally, the intimal foam cell cushion made up of one or more layers of cholesterol lipid laden lipophages, was identified as the earliest histologically recognizable stage of plaque formation in human arteries (1-3).

Data were also collected indicating that the lipids of the atherosclerotic plaque originated chiefly from the plasma (1-286). Recent findings demonstrating arterial wall synthesis of cholesterol and phospholipids do not constitute a refutation of the conclusion that the accumulated lipids in atherosclerotic plaques are derived primarily and chiefly from the plasma (287-289).

A few workers, particularly in Great Britain, adhere to an alternative concept concerning the basic pathogenesis of the atherosclerotic plaque—the thrombogenic theory (449-450).

All these observations by pathology brought to the fore—but could not answer—the critical question: Is there any etiologically significant interrelationship among cholesterol and fat in the diet, the plasma and the lesion (*cf.* refs. 1-3, 4, 10, 11, 69, 70, 170)? In assessing this contribution of pathology it is worth emphasizing that the proper formulation of a scientific question is often the greatest impetus to its solution.

2 *No pathologic Atherosclerosis in Cases with and without Clinical Coronary Heart Disease*—More recently, careful pathologic investigations in the United States showed that atherosclerosis of the coronary arteries is generally of greater severity in cases of clinical coronary heart dis-

case compared with cases clinically free of this disease ante mortem (290-292 294-300) This demonstration is of key practical and theoretical importance for research in this field (see below)

3 *Atherosclerosis in Nephrosis Hypothyroidism Diabetes, Essential Familial Xanthomatosis*—Clinical research long ago noted that athero-

SERUM LIPIDS IN MIDDLE-AGED NORMAL MALES AND MALES WITH CORONARY HEART DISEASE




FIG 19 Data on normal Central Americans from Mann G V Munoz J A and Scrimshaw N S (109) on normal South African Bantus from Walker A R P and Arvidsson U B (97) and Higginson J and Pepler W J (93) on normal Navajo Indians from Page I H Lewis L A and Gilbert J (257) on Bostonians with and without clinical coronary heart disease (all men under age 40) from Gertler M M and White P D (202) on Chicagoans with coronary heart disease (all men under age 50) from Stamler J Pick R and Katz L N (645) The data from analyses done in different laboratories are not amenable to strict quantitative comparison

sclerosis was more frequent premature and severe in a number of disease states characterized by a sustained period of hypercholesterolemic hyperlipemia—nephrosis hypothyroidism essential familial xanthomatosis diabetes (1) This observation pointed up the relationship between hypercholesterolemia and atherogenesis in a group of diseases

in which hypercholesterolemia is mainly endogenous (and not diet induced) It therefore lent further support to the concept that atherosclerosis is a metabolic disease However, it did not fundamentally illuminate the problem of the relationship of nutrition to this disease beyond indicating that it was not exclusively diet induced

Serum Cholesterol and Lipoprotein levels in Clinically healthy Americans and in males with Coronary Heart Disease




FIG 20 Lawry E Y Mann G V Peterson A Wysocki A F O'Connell R and Stare F J (112) Note the differences in serum cholesterol levels between the clinically normal and the myocardial infarction groups Note also that these differences are greatest in the youngest age group and decline with age

4 *Serum Cholesterol Lipid Lipoprotein Levels in Clinical Coronary Heart Disease* — Clinical investigation made a critically important contribution through studies on cholesterol lipid lipoprotein metabolism in patients with coronary heart disease It conclusively demonstrated that any group of middle aged males with coronary heart disease had significantly higher levels of circulating cholesterol lipids β lipoproteins than a

comparable clinically normal group* (Figs 19 to 21) (1, 3-5, 12-14, 21, 22, 69, 71, 112, 171, 202, 279-281, 283 301-318, 645)

It should be emphasized that these observations refer to *group* differences. Within any group of middle aged males with coronary heart disease, a small percentage of *individuals* will invariably exhibit low plasma cholesterol levels. Correspondingly, epidemiologic and clinical

New Episodes of Coronary Heart Disease in Relation to levels of Serum Cholesterol in the Base Population of Men Age 40-59 Cooperative study on Lipoproteins & Atherosclerosis




FIG 21 Gofman J W *et al* and Andrus E C *et al* (281) Each circle represents an episode of clinical coronary heart disease in an individual with the indicated value for serum cholesterol prior to illness. Note that the 40 per cent of the population with the lower cholesterol levels yielded 23 per cent of the cases (13 of 57) whereas the 40 per cent with the higher cholesterol levels yielded 66 per cent of the cases (38 of 57) (compare Fig 18)

studies demonstrate that a small percentage of cases of clinical coronary disease do in fact develop from people with low normal cholesterol levels (Fig 21) (2 23 72, 93-98 112 116 117 134-137, 209 272 278, 645) It is of decisive theoretical importance that the percentage of such cases is *low*. These "exceptions to the rule" certainly merit intensive study.

*Recent reports have dealt extensively with the comparative merits of serum cholesterol vs lipoprotein fractions in segregating coronary patients from clinically normal persons (see particularly refs 112 279 281 and other cited refs) In view of these comprehensive discussions this problem is not dealt with in this review.

since elucidation of the etiology and pathogenesis of disease in the individuals should contribute significantly to clarifying the overall problem.

It was further observed that the differences in serum cholesterol between men with and without clinical coronary disease were greater, the younger the two groups (Fig. 20). This difference emerged despite the fact that the "control" group was not really a normal group, but only a *clinically* normal group. It was not free of atherosclerosis morphologically, since most American males by middle age have gross coronary atherosclerosis (290-292, 294-298, 300). Therefore the two groups represented more disease vs. less-disease, not disease vs. no-disease.

The latter comparison could be made only by analyzing Americans vs. Bantus or Guatemalans, for example. Then the disparity in levels of serum cholesterol lipids & lipoproteins would be even greater (Fig. 19).

This comparison—or more precisely contrast—brings once more to the fore several critical questions. What are proper standards for normal—and optimal—serum cholesterol? Why do extensive segments of our clinically normal healthy middle aged population have serum cholesterol levels in a range inordinately predisposing them to coronary heart disease? What is the origin of this lipid metabolic derangement? Is life span pattern of diet a decisive causative factor—and if so what in the diet?

5. *Dietary Effects on Cholesterolemia—Weight Gain and Loss, Low Fat, Low Cholesterol Diets*—The last few years have witnessed a vigorous attack on these problems by clinical investigation. As a result, it has been conclusively demonstrated—contrary to opinions widely held only a decade ago (1, 352)—that serum cholesterol lipid lipoprotein levels of middle aged Americans are highly amenable to alteration by dietary means. Thus simple overeating with rapid weight gain was found to be associated with a marked increase in serum cholesterol (283, 319-323). Further increased intake of saturated fats (e.g. pemmican) in an iso-caloric diet also was correlated with a rise in cholesterol (324, 325). These data from human experiments reinforced the conclusion that diets high in calories, fats, saturated fats, cholesterol lead to elevated plasma cholesterol levels.

Several clinical studies also indicated that ingestion of a high fat meal is followed postprandially by an increased blood coagulability *in vitro*. Clotting time was reported to be decreased irrespective of the type of fat used, i.e. animal or vegetable, saturated or unsaturated (326-345). It was not altered after low fat meals. It should be noted that several negative reports challenge the validity of this observation. It has also been reported that ingestion of certain saturated fats, particularly butter fat, results in a significant retardation of fibrinolysis, i.e., of the ability of plasma to lyse formed clots. No data are available concerning the *in vivo* effects of fat vs. non fat meals on the clotting mechanism. Furthermore the relationship—if any—of this phenomenon to atherogenesis, thrombogenesis, and development of clinical atherosclerotic disease re-

means to be elucidated. This is a relatively new area of investigation. Undoubtedly it will be thoroughly explored in the next few years. At this juncture estimates of the clinical significance of these observations on diet and clotting can at best be only conjectural. Should the initial findings receive firm verification, they would constitute a further support for the nutritional metabolic theory, with obvious practical implications for clinical medicine in its efforts to prevent and treat atherosclerotic disease (see below).

EFFECT OF WEIGHT LOSS ON SERUM CHOLESTEROL AND LIPOPROTEIN LEVELS




FIG 22 Walker W J Lawry E Y Love D E Mann G V Levine S A and Stare F J (319) cf also refs 320 346 347. Note that during a period of active weight loss serum cholesterol lipoprotein levels fall. Note also that they remain at the reduced concentrations with maintenance of weight loss.

Clinical investigation also demonstrated that plasma cholesterol concentration may be readily lowered by dietary means. It tends to fall during periods of weight reduction (Fig 22) (319 320 346-349). Low fat, low-cholesterol diets also effect such a decline in most patients (Figs 23 and 24) (1, 3-6 12-14 21 22 71 279 319 346-349 351, 353-375). The response apparently occurs irrespective of gross weight loss. It tends to supervene rapidly being manifest usually during the initial weeks on diet. However on occasion it develops more slowly and gradually. The fall in serum cholesterol is often substantial particularly in hypercholesterolemic patients. It apparently persists in most persons with continued adherence to the low fat, low cholesterol diet.*

*Limited data indicate that addition of cholesterol to this diet without increase in fat intake has little influence on serum cholesterol levels at least in short term experiments (71 107 375 3 6).

6 *Dietary Effects on Cholesterolemia—Unsaturated Vegetable and Marine Oils—Complete Substitution, Partial Substitution and Supplementation Studies*—Varying results were initially obtained in investigations on the effects of substituting unsaturated vegetable lipid for carbohydrates in low fat, low cholesterol regimens. Some studies reported a tendency for serum cholesterol to rise again (354, 358, 362, 365, 371, 376). Others observed an additional slight reduction in cholesterolemia (Fig 25) (372

DECREASE IN SERUM CHOLESTEROL DURING INGESTION OF THE RICE DIET

Summary of representative studies




FIG 23 Keys A, Mickelsen O, Milles E, V O and Chapman C B (354), Kempner W (353), Watkin D M, Froeb H F, Hatch F T and Gutman A B (355), Starke H (350)

377–379). Still others observed essentially no effect (380). The basis for this apparent disparity in results remains to be elucidated. It is presumably unrelated to the degree of unsaturation of the vegetable oils utilized (see below) since this was similar in the foregoing studies yielding apparently contradictory data. It may be a function of plant sterol or protein content of the respective diets (see below). It is generally agreed that incorporation of saturated animal fat *e.g.* butter fat led to a re-elevation of serum cholesterol; however, the saturated fat ethyl stearate had no such effect (380).

Several other recent studies investigated the effects of unsaturated oils on cholesterolemia. It was shown that isocaloric substitution of unsaturated vegetable or marine oils for saturated animal lipids in the diet resulted in significant falls in plasma cholesterol levels (Figs 25 and 26) (166, 365, 369, 370, 372, 373, 375-383, 385-405, 415).

EFFECTS OF REDUCED FAT ON SERUM CHOLESTEROL IN MEN ON DIET CONSTANT IN CALORIES-PROTEIN MINERALS-VITAMINS




Fig 24 Keys A Anderson J T Fidanza F Keys M H and Swahn B (364)

The response with substitution of unsaturated vegetable oils for saturated animal fats tended to occur promptly within a few days and persisted for at least several months with continued adherence to this dietary regimen. These experiments were all of the *complete substitution* type i.e. they uniformly involved the virtually complete exclusion of all animal fats from the diet. In many cases formula diets—with unsaturated vegetable oils supplying as much as 60 per cent of the total calories—were utilized to assure accomplishment of this objective.

Supplementary to this observation is the finding that Americans who are habitual pure vegetarians have significantly lower serum cholesterol levels than omnivores or ovo lacto vegetarians—despite a high intake of total (all vegetable) fat (Fig 27) (251). This is consistent with previously cited findings in Yugoslavs (67). In this connection the *negative* correlation between vegetable fat intake and arteriosclerotic heart disease death rates for 22 countries is also noteworthy (Fig 12) (139).

6 *Dietary Effects on Cholesterolemia—Unsaturated Vegetable and Marine Oils—Complete Substitution, Partial Substitution and Supplementation Studies*—Varying results were initially obtained in investigations on the effects of substituting unsaturated vegetable lipid for carbohydrates in low fat, low cholesterol regimens. Some studies reported a tendency for serum cholesterol to rise again (354, 358, 362, 365, 371, 376). Others observed an additional slight reduction in cholesterolemia (Fig. 25) (372).

DECREASE IN SERUM CHOLESTEROL DURING INGESTION OF THE RICE DIET

Summary of representative studies




FIG 23 Keys A, Mickelsen O, Milles E, V O and Chapman C B (354), Kempner W (353), Watkin D M, Froeb H F, Hatch F T and Gutman A B (355), Starke H (350).

377–379) Still others observed essentially no effect (380). The basis for this apparent disparity in results remains to be elucidated. It is presumably unrelated to the degree of unsaturation of the vegetable oils utilized (see below) since this was similar in the foregoing studies yielding apparently contradictory data. It may be a function of plant sterol or protein content of the respective diets (see below). It is generally agreed that incorporation of saturated animal fat, e.g., butter fat, led to a re-elevation of serum cholesterol; however, the saturated fat, ethyl stearate, had no such effect (380).

More recently, reports have begun to appear on the effects of only *partial substitution* (Fig 25) (369, 372, 394-398, 406, 407) Thus, in one study a 140 gram fat diet was utilized composed of 40 grams of mixed food fats plus 100 grams of butter—with experimental substitution of 100 grams of unsaturated oil for the butter (394-397) In another study, 100 gram animal fat diets were utilized—with partial substitution experimentally of 70 grams of corn oil for 70 grams of animal fat (leaving 30 grams of animal fat) (398) This partial substitution of unsaturated

DECREASE IN SERUM CHOLESTEROL FOLLOWING SUBSTITUTION OF VEGETABLE FAT FOR ANIMAL FAT IN AMERICAN- "TYPE" HABITUAL DIETS




FIG 26 Kinsell L W Iarttridge J Bohng L Margen S and Michaels J (381) (*cf also refs 382 383 385-388*) Ahrens E H Insull W Blomstrand R Hirsch J Tsaltas T T and Peterson M L (393) (*cf also refs 372 and 392*) Beveridge J M R Connell W F Mayer G A Firstbrook J B and DeWolfe M S (377) Malmros H and Wigand G (365)

FIG 25 Beveridge J M R Connell W F and Mayer G A (378) Note in the upper figure the decline in serum cholesterol with a fat free diet This decline proceeds further with a diet containing 60 per cent of calories from corn oil It is partially reversed with a diet containing 20 per cent of calories from butter completely reversed with 60 per cent of calories from butter Note in the middle figure the decline in serum cholesterol with complete substitution of a corn oil formula diet This decline is partially reversed in a few days when the corn oil is replaced by butter chickenfat lard or beef Note on the lower figure the decline in serum cholesterol with partial substitution of corn oil for butter fat

Effects of various types of formula diets on Cholesterolemia




FIG 25—Legend on opposite page

Such a supplementation regimen rigidly applied, would tend, of course, to increase total caloric ingestion. On this basis alone, a rise in serum cholesterol might have been anticipated (see above). In actuality subjects in some cases spontaneously adjusted their *ad lib* normal diets, thereby accommodating for the extra calories and avoiding weight gain. In such supplementation studies a fall in plasma cholesterol was reported by one laboratory in limited experiments on a few South Africans (Fig. 28).

Effects of a Supplement of Essential Fatty Acids on Serum Lipids in a Patient with Hypercholesterolemic Hyperlipemia




FIG 29 Nothman M M Belin L and Proger S (410) The arrow (↓) indicates the date of onset of daily supplementation with the mixture of essential fatty acids

(379) This was the case even when one of the animal fat sources (eggs) also yielded a high cholesterol intake. On the other hand studies by other groups yielded less clearcut results (Fig. 28) (383 385 408 409). Further another group reported that addition of fat—either vegetable or animal—to a mixed diet induced increased cholesterolemia (376–378 389–390). Precise experimental design and specific data from this study were not presented.

At the October 1957 meeting of the American Heart Association, im

oils for saturated fats was also effective in lowering serum cholesterol levels

Another variation in experimental design has been explored, viz unsaturated vegetable oil *supplementation* of a high calorie, high fat, high saturated fat, high cholesterol diet. This involved neither withdrawal of fat, nor any other departure from the customary fare

NUTRITIONAL AND BIOCHEMICAL FINDINGS IN OMNIVOROUS VS HERBIVOROUS AMERICANS




Fig 27 Hardinge M G and Stare F J (251) Note the markedly lower serum cholesterol level in the pure vegetarians on a diet of 130 grams of vegetable fat per day (35 per cent of calories)

EFFECTS ON SERUM CHOLESTEROL OF VEGETABLE FAT SUPPLEMENTATION OF AN ANIMAL FAT CONTAINING DIET




Fig 28 Bronte Stewart B, Antonis A, Eales L and Brock J F (379), Kinsell L W, Michaels G D, Cochrane J C, Partridge J W, Jahn J J and Balch H E (383) Note the tendency for cholesterol levels to fall with vegetable fat supplementation of an animal fat containing diet

volved (382, 383, 385-388, 393, 403) On the other hand, one group of investigators has presented considerable evidence to support their conclusion that plant sterols are decisively responsible for the cholesterol emia reducing effects of such products as corn oil (376-378, 389, 390, 414) Further work is apparently needed to resolve this disagreement

The serum cholesterol response with vegetable and marine oils has also been attributed to their high content of unsaturated fatty acids This

Effects of Unsaturated vs Saturated Vegetable Oil on Serum Cholesterol in Subject on a Formula Diet

NS ♂ 27 yrs




FIG 30 Abrens E H Insull W Blomstrand R Hirsch J Tsaltas T T and Peterson M L (393) Note the renewed elevation of serum cholesterol and phospholipids with transfer from corn oil to coconut oil (high in saturated fat) formula diet PFC dietary protein fat and carbohydrate as percent of total calories TC and FC total and free cholesterol respectively PL phospholipids TG triglycerides

pressive data were presented demonstrating marked, consistent falls in serum total lipids, fatty acids, cholesterol and cholesterol/phospholipid ratios in 22 hypercholesterolemic patients given 25 to 30 cc daily of a mixture of linoleic acid (49%), linolenic acid (49%) and arachidonic acid (2 per cent) (Fig 29) (410). There were no dietary restrictions. The results, therefore, represent the effects of unsaturated oil *supplementation*—in this case unsaturated oils of the polyethenoic, essential fatty acid (EFA) type (see below).

Positive results with safflower oil (300 cc daily) "supplementation" (isocaloric) were also reported to the 1957 meeting of the American Society for the Study of Arteriosclerosis by another investigator in studies on diabetics (411). It was further noted that spontaneous retinal hemorrhages ceased to occur during the period of safflower oil ingestion and that tourniquet tests revealed decreased capillary fragility (411)—a finding apparently confirming a previous report that essential fatty acids enhance capillary integrity (412). This finding is particularly intriguing in relation to the problem of intramural hemorrhage in atherosclerotic plaques as a significant mechanism in the conversion of morphologic coronary lesions into clinical coronary heart disease (413).

7 *Essential Fatty Acids and the Mechanism of the Effects of Unsaturated Oils*—Extensive studies are being reported on the factor or factors responsible for the decline in cholesterolemia in these experiments with unsaturated vegetable oils, and the mechanism of their effect. From observations to date it has been suggested that in the substitution experiments removal of dietary cholesterol was not significantly responsible for the observed results. Thus, addition of cholesterol to a vegetable oil diet induced little or no renewed rise in serum cholesterol level (372, 375-379, 382, 383, 385, 389, 390).

In the few such studies reported to date cholesterol was added to a *purified formula ration* high in unsaturated vegetable oils and low in saturated animal fats. Under these conditions a slight rise in serum cholesterol supervened over a relatively short time course (1 to 10 weeks). Very different results have been obtained in several experimental animals: i.e. the development of severe hypercholesterolemia with the incorporation of cholesterol and unsaturated vegetable oil in the diet (see below). In view of these findings, additional studies are needed before any conclusions are justified concerning the interrelated effects of cholesterol and different fats on cholesterolemia in man. It may well turn out that the findings in substitution studies result from removal of dietary fats and cholesterol.

In other studies evidence was obtained indicating that the cholesterolemia reducing influence of unsaturated vegetable oils could not be attributed to their content of phospholipids (382, 383, 385-388). It was also tentatively concluded by some workers—based on their research findings—that the plant sterols in vegetable oils are not critically in

a basal low fat diet of natural foodstuffs (380) Feeding the saturated fat, ethyl stearate with this diet was also associated with a fall in serum cholesterol—a finding possibly related to incomplete absorption of this lipid Two other groups of investigators also observed that feeding a corn or safflower oil formula diet resulted in a marked rise in serum EFA, specifically a more than two-fold increase in the linoleic acid content of serum cholesterol esters and phospholipids and an elevenfold elevation in the amount of this FFA in triglycerides (402, 423)

These observations with pure lipids seem to indicate that neither phospholipids nor plant sterols nor oleic acid are essential factors in the serum cholesterol response with vegetable oils in either substitution experiments with formula diets or supplementation experiments with *ad lib* mixed diets They are consistent with the hypothesis that the essential fatty acids in unsaturated vegetable and marine oils are decisive factors in the mechanism of their effect on serum cholesterol However they do not represent definitive proof of the validity of this hypothesis

Recent reports have also served to clarify the mechanism of action of unsaturated oils on cholesterol metabolism Thus it was shown in tracer studies using cholesterol-4 C¹⁴ that changing a hypercholesterolemic patient's diet from *ad lib* feeding to a 40 per cent butter formula—with a resultant rise in plasma cholesterol from 478 to 720 mg %—was associated with a marked decrease in fecal sterol excretion When corn oil replaced the butter plasma cholesterol decreased to 350 mg % Concomitantly fecal excretion of cholesterol and its end products rose markedly (424) The data strongly indicated that fecal cholesterol is derived from the plasma and that bile acid excretion is not a major pathway for disposal of cholesterol by the human organism Moreover they pointed to the conclusion that the character of dietary lipids influences cholesterolemia by increasing or decreasing the fecal excretion of cholesterol and its transformation products (424-425)

One further observation of a general theoretical nature may be relevant to this whole problem It seems particularly premature to conclude that the basic lipid metabolic derangement leading to atherogenesis is a resultant of *primary* EFA deficiency in human diets The fact is that usual American diets for example are not low in essential fatty acids—quite the contrary Thus butter, margarine and other hydrogenated vegetable fats contain 2 to 10 per cent EFA Milk and milk products have a significant EFA content Animal fats in general are composed of about 9 per cent essential fatty acids (217-218 420 422 427-429) Most Americans therefore ingest large amounts of EFA every day irrespective of their vegetable and marine oil intakes They actually consume much more EFA than most other peoples—much more than any estimated minimum daily requirement Consistent with this fact is the observation that American and British patients with coronary disease do not exhibit ab

interpretation is supported by observations indicating that the pattern of change in serum cholesterol levels correlated closely—but not completely—with the degree of unsaturation (iodine number) of the vegetable oils (372, 379, 393–397). It was reinforced by data showing that serum cholesterol rose anew when saturated vegetable fats (margarine or coconut oil or hydrogenated peanut fat or hydrogenated corn oil) were substituted for unsaturated vegetable oils (Fig. 30) (372, 378, 379, 393–397, 402). On the other hand, other workers failed to observe this phenomenon (380, 414), hence this aspect of the mechanism problem must also be regarded as *sub judice* at present. The apparent disparities in findings in man compared with other species have further complicated the problem (see below).

The unsaturated vegetable oils contain large amounts of both mono- and poly ethenoic fatty acids. Oleic acid typifies the former; linoleic acid the latter. Linoleic acid is one of the essential fatty acids (EFA) (393, 416–420). It is not surprising therefore—in view of the high EFA content of unsaturated vegetable oils—that speculation has been rife on the role of essential fatty acids in the mechanism of the cholesterolemia-reducing effect (426; cf. discussion in refs. 6, 7, 20–23, 71, 142, 166, 369–373, 376–379, 381–383, 385–400, 421, 422, 426, 461).

The achievement of a definitive experimental resolution of this scientific problem is rendered difficult by the lack of available chemically pure lipid compounds. Since most unsaturated vegetable oils contain both oleic and linoleic acids in quantity as well as plant sterols and phospholipids, results of feeding experiments with them do not readily yield a clearcut differential answer (393).

Recently, reports have begun to appear on the effects of feeding pure preparations. Mention has already been made of the impressive reductions in serum cholesterol achieved in 22 patients in a supplementation study utilizing 25 to 30 cc. of a mixture of three essential fatty acids (Fig. 29) (410). Another group initially reported a marked reduction in serum cholesterol in one patient with ingestion of a formula diet containing a synthetic lipid mixture made up of oleic acid (75 per cent), linoleic acid (2 per cent), palmitic acid (19 per cent) and stearic acid (5 per cent) (415). At the November 1957 meeting of the American Society for the Study of Arteriosclerosis, these workers reported their results in experiments with ethyl oleate, triolein, ethyl linoleate (96 per cent pure), trilinolein, and a phosphatide of animal origin containing approximately 12.5 per cent tetraenoic (presumably arachidonic) acid. Each of the latter three poly ethenoic EFA preparations produced a significant sustained fall in plasma cholesterol associated with a rise in EFA esterified with cholesterol (405). The two mono ethenoic preparations, ethyl oleate and triolein, were apparently without effect—a finding in accord with the conclusions of others (393–397). Similar results were obtained by another investigator working with an ethyl linoleate concentrate incorporated in

Chapter 5

Nutrition and Atherosclerosis (*Continued*)

I The Animal-Experimental Findings

THE animal experimental laboratory is the third broad source of data concerning the interrelationship among diet cholesterol lipid lipoprotein metabolism and atherogenesis.

1 *Historical Development*—In evaluating the contribution made by this methodology, it is important to appreciate the evolution of research in this field. Experimental atherosclerosis has a pre-history and a history. The dividing line is roughly the year 1910. Up until then every effort to reproduce atherosclerosis in laboratory animals had met with failure. Different types of vascular sclerosis had been produced by a variety of means but never atherosclerosis (1 10 11 170 461). In 1908-1912 success was first achieved in producing atherosclerosis by feeding animal tissues to rabbits. It was soon demonstrated that lipid constituents (*i.e.* cholesterol and neutral fats)—and not proteins—were the decisive atherogenic factors in this diet (1 451). Thus hypercholesterolemia and atherosclerosis resulted when cholesterol and fat *per se* were added to the feed.

2 *Diet induced Hypercholesterolemia and Atherogenesis in Different Species*—In the last decade hypercholesterolemia and atherosclerosis have been induced in all the major laboratory species herbivorous omnivorous and carnivorous (1 3 4 6 12-14 461). In the chick like the rabbit this was achieved by feeding a diet supplemented with cholesterol and oil. In these two susceptible species it was further demonstrated that atherosclerosis could be induced without massive hypercholesterolemia and organ lipidosis (Fig 32) (1 451 452). This was accomplished by feeding small amounts of cholesterol lipid over long periods of time. In cockerels minimal hypercholesterolemia developed with diets of mash plus $\frac{1}{4}$ per cent cholesterol and 5 per cent cottonseed oil. Cross atherosclerosis of the aorta was demonstrable after 35 weeks on this regimen—a time period representing roughly 20 to 25 per cent of the chick's life span (1 452). This approaches the dietary metabolic and temporal prerequisites for atherogenesis in man.

In species more resistant than the chick and the rabbit it proved possible to induce hypercholesterolemia and atherosclerosis by using more complex forms of the same fundamental nutritional metabolic alteration.

normal, low levels of polyethenoic essential fatty acids in their sera (Fig 31) (104, 430)

Therefore—even if a metabolic deficiency of EFA were actually present in the American population—it could not be the resultant of a primary dietary lack of EFA *per se*. It could only be a *relative* deficiency, *secondary* perhaps to excessive intake of saturated fats, cholesterol, refined carbohydrates and/or total calories. This would be a derivative imbalance, with inadequacy of EFA (if actually present) due to an inordinate need stemming from other dietary aberrations and excesses.

Unsaturated Fatty Acids in the blood of patients with Coronary Artery Disease




FIG 31 James A T Lovelock J E Webb J and Trotter W R (430) Note that the clinically normal and coronary disease patients have similar patterns of blood fatty acids including essential fatty acids

Undoubtedly, work in the years immediately ahead will effectively clarify this group of problems concerning the unsaturated vegetable oils and essential fatty acids. Irrespective of the ultimate resolution of these questions the demonstration of an effect of dietary unsaturated oils on serum cholesterol levels represents one of the important advances of recent years. By deepening and refining our knowledge it has significantly reinforced the nutritional metabolic cholesterol lipid lipoprotein theory of atherogenesis. It is also of great practical meaning (see below).

In summary then the recent findings of clinical pathological research have amply demonstrated a close relationship between level of cholesterolemia and development of atherosclerotic disease on the one hand, and between pattern of diet and level of cholesterolemia on the other. The clinico-pathologic data parallel and complement the epidemiologic findings.

Chapter 5

Nutrition and Atherosclerosis (*Continued*)

F The Animal-Experimental Findings

THE animal experimental laboratory is the third broad source of data concerning the interrelationship among diet, cholesterol lipid lipoprotein metabolism and atherogenesis

1 *Historical Development*—In evaluating the contribution made by this methodology, it is important to appreciate the evolution of research in this field. Experimental atherosclerosis has a pre-history and a history. The dividing line is roughly the year 1910. Up until then every effort to reproduce atherosclerosis in laboratory animals had met with failure. Different types of vascular sclerosis had been produced by a variety of means but never atherosclerosis (1, 10, 11, 170, 461). In 1908–1912 success was first achieved in producing atherosclerosis by feeding animal tissues to rabbits. It was soon demonstrated that lipid constituents (*i.e.*, cholesterol and neutral fats)—and not proteins—were the decisive atherogenic factors in this diet (1, 451). Thus hypercholesterolemia and atherosclerosis resulted when cholesterol and fat *per se* were added to the feed.

2 *Diet induced Hypercholesterolemia and Atherogenesis in Different Species*—In the last decade hypercholesterolemia and atherosclerosis have been induced in all the major laboratory species: herbivorous, omnivorous and carnivorous (1, 3, 4, 6, 12–14, 461). In the chick, like the rabbit, this was achieved by feeding a diet supplemented with cholesterol and oil. In these two susceptible species it was further demonstrated that atherosclerosis could be induced without massive hypercholesterolemia and organ lipodosis (Fig. 32) (1, 451, 452). This was accomplished by feeding small amounts of cholesterol lipid over long periods of time. In cockerels minimal hypercholesterolemia developed with diets of mash plus $\frac{1}{4}$ per cent cholesterol and 5 per cent cottonseed oil. Gross atherosclerosis of the aorta was demonstrable after 35 weeks on this regimen—a time period representing roughly 20 to 25 per cent of the chick's life span (1, 452). This approaches the dietary, metabolic and temporal prerequisites for atherogenesis in man.

In species more resistant than the chick and the rabbit it proved possible to induce hypercholesterolemia and atherosclerosis by using more complex forms of the same fundamental nutritional/metabolic alteration

Thus, in the dog positive results were obtained by feeding a high-cholesterol, high lipid diet combined with hypothyroidism, in the monkey and the rat by feeding a high-cholesterol high lipid diet combined with a nutritional deficiency of the sulfhydryl containing amino acid methionine (1 453-458) More recently hypercholesterolemia and atherosclerosis were produced in monkeys and rats with high fat high cholesterol diets alone (457 459 460)

**Atherosclerosis in Cockerels with Minimal
Hypercholesterolemia due to feeding mash supplemented
with $\frac{1}{4}\%$ Cholesterol + 5% Cottonseed Oil
--Age Period Weeks 5-40**




FIG 32 Stamler J and Katz L N (452) Here and in subsequent figures the symbol C is for cholesterol O for oil $\frac{1}{4}\text{CO}$ is $\frac{1}{4}$ per cent cholesterol + 5 per cent cotton seed oil in chick starter mash the symbol O without a numerical prefix is 5 per cent cottonseed oil throughout Unless otherwise indicated all groups had similar patterns of feed-caloric intake growth and development Here and in subsequent figures plasma cholesterol levels are in mg per cent aorta lesions are gross plaques in the thoracic aorta incidence is the percent of birds per group with lesions grade is the average grade of lesions in birds with lesions grading is on a scale of 0 to 4 (cf ref 1)

Moreover coronary thrombosis and myocardial infarction were recently induced in rats by dietary means This was accomplished by giving rats thiouracil and a diet containing 5 per cent cholesterol 2 per cent sodium cholate and 40 per cent butter fat (or 20 per cent Crisco + 20 per cent cod liver oil) for 14 weeks Coronary thrombosis and myocardial infarction occurred in 20 per cent of the animals Marked lipid accumulation was observed in the coronary arteries but not atherosclerosis When 40

per cent fat was given in the form of 20 per cent corn oil plus 20 per cent Crisco coronary thrombosis and myocardial infarction did not supervene (see previous and subsequent discussion on unsaturated oils) (462) An other investigator induced calcified atheromas and coronary occlusion by a combination of nutritionally induced hyperlipemia and vascular injury (463 464) These accomplishments go a long way toward meeting a longstanding major challenge to atherosclerosis research

3 *Dietary Cholesterol and Lipid in Experimental Atherogenesis*—In viewing this research development in perspective it is essential to appreciate the common thread running through all these achievements Invariably the *sine qua non* for experimental atherogenesis was a nutritional

EFFECTS OF DEFATTED MASH IN CHOLESTEROL-FED COCKERELS




FIG 33 Stamler J Pick R and Katz L N (471) Note the minimal hypercholesterolemia and atherogenesis in chicks fed 1 per cent cholesterol in fat extracted mash

alteration the critical change being to a high fat high cholesterol intake maintained over a sustained period and leading to a hypercholesterolemic hyperlipemia With a few exceptions (see below) dietary excess of both cholesterol and neutral fat played a key role in all species studied This combination of the two appeared to be decisively important Thus excess cholesterol feeding in the complete absence of neutral fat and excess neutral fat feeding in the complete absence of cholesterol tended to be minimally atherogenic (Fig 33) (1 461 465 471) Further at a given level of excess dietary cholesterol atherogenicity tended to rise as neutral fats—either unsaturated vegetable oils or saturated animal fats—were added in increasing amounts Without this combined cholesterol lipid nutritional aberration other changes in diet were by themselves generally not atherogenic

Several partial exceptions to the foregoing statements have emerged from recent studies. Thus, moderate hypercholesterolemia and atherosclerosis may develop in animals on a cholesterol free diet—in ground squirrels (hibernators) after 3 to 18 months on an *ad lib* fattening diet high in vegetable lipid from peanuts (466), in geese after 3 months on a forced fed, high carbohydrate, fattening regimen—a diet rendered far more atherogenic with addition of cholesterol (467), in rabbits after 12 months on an *ad lib* diet high in vegetable fat from peanuts (468), in rabbits after 100 days on a purified ration containing 20 per cent highly saturated hydrogenated coconut oil (469). The published reports, which are available on the first three of the above studies indicate that atherogenesis was relatively slight to moderate on these non-cholesterol containing regimens, in marked contrast to the extensive, severe lesions developing rapidly in cholesterol lipid fed rabbits or fowl. These data suggest that hypercholesterolemia and atherosclerosis may be induced—to a limited extent in some species at least and after relatively long time periods—by excessive intake of calories *per se*, even when derived from carbohydrates and/or unsaturated vegetable oils, without cholesterol or animal fat*. They further suggest that highly saturated fat without cholesterol may induce hypercholesterolemia and atherogenesis, at least in rabbits even without concomitant development of obesity. Finally, they support the significant conclusion that *prolonged ingestion of a diet combining excess of cholesterol lipids† and calories has particularly pernicious hypercholesterolemic and atherogenic effects. This is the typical contemporary American diet.*

4 *Diet induced Reversibility of Experimental Atherosclerosis*—Another major early contribution of experimental atherosclerosis was the demonstration that the atherosclerotic plaque is, within limits a reversible lesion. This was shown in rabbits chicks and dogs (1, 451, 470, 472–474). Hypercholesterolemia and atherosclerosis were first produced by feeding an atherogenic diet upon withdrawal of this diet and substitution of a low fat low cholesterol ration hypercholesterolemia and atherosclerosis underwent gradual regression.

Earlier studies of this type were confined to analyses of lesions in the aorta. Recently it was clearly demonstrated that coronary lesions were also reversible by dietary and other means (see below) (1, 472–473).

5 *Unsaturated Vegetable Oils—Essential Fatty Acids*—The recent work on the effects of unsaturated vegetable oils on cholesterolmia in man led to exploration of this problem in experimental animals. Before consideration of the newer findings it is worth noting that for decades experimental atherosclerosis was routinely produced—originally in rab

*The interpretation of these animal-experimental findings in relation particularly to the recent observations on unsaturated oils vs saturated fats in man must be regarded as *sub judice* pending additional data. This problem is discussed further below.

†See preceding footnote

bits and latterly in chicks and other animals—utilizing diets containing cholesterol plus unsaturated vegetable oils. In apparent contradiction to the limited findings of supplementation studies in man (see above), the incorporation of large amounts of unsaturated vegetable oils in the high cholesterol diets of experimental animals did not inhibit hypercholesterolemia and atherogenesis (1, 451). In fact it is an old observation in rabbits and chicks that ingestion of mash supplemented with cholesterol plus vegetable oil resulted in more marked hypercholesterolemia than feeding

EFFECTS OF COTTONSEED OIL VS HYDROGENATED VEGETABLE FAT IN CHOLESTEROL FED COCKERELS

— Series 47 13-23 weeks




FIG 34 Stamler J Pick R and Katz L N (476) Note the essentially similar findings in the two groups despite the marked differences in amounts of saturated fat and essential fatty acids in the two rations

mash plus cholesterol alone (1). Thus, hypercholesterolemia and atherogenesis were minimal when cholesterol was fed with defatted mash *vs* mash presumably deficient in essential fatty acids (Fig 33) (471). Further as already noted above rations high in unsaturated vegetable fats (*e.g.* from peanuts) may induce hypercholesterolemia and atherosclerosis in rabbits and squirrels in the complete absence of dietary cholesterol (468). Moreover, feeding mash plus 20 per cent cottonseed oil alone (without cholesterol) apparently induced slight hypercholesterolemia (1). Again these are results in apparent contradiction to those in man.

Recent experiments showed that feeding chicks 1 per cent cholesterol mashes supplemented with varying lipids at the 5 to 10 per cent level—*e.g.* relatively saturated lard or hydrogenated vegetable fat *vs* unsaturated vegetable oils (linseed cottonseed corn oil or a linoleic acid con

EFFECTS OF SATURATED ANIMAL FAT VS UNSATURATED VEGETABLE FAT IN CHOLESTEROL-FED COCKERELS

S-37 12-17 Weeks




FIG 35 Stamler J Pick R and Katz L N (476) Note again the essentially similar findings in the two groups

Effects of Various High Oleic Acid Preparation in Cholesterol fed Cockerels S-51 - Weeks 9-14




FIG 36 - Stamler J Pick R and Katz L N (477) All groups received 1 per cent cholesterol + 10 per cent fat. In the control group the fat was cottonseed oil. The oleic acid concentrate was 94 per cent oleic 17 per cent linoleic acid the olive oil was 86 per cent oleic 6 per cent linoleic the Neo fat was 73 per cent oleic 12 per cent linoleic the chickenfat was 52 per cent oleic + vaccenic 17 per cent linoleic acids. The results of this experiment were somewhat unusual in that the group receiving oleic acid had significantly less aorta and coronary atherogenesis than the other groups despite similar marked hypercholesterolemia. It is not clear why this one group on a high oleic acid diet exhibited these findings different from the other groups receiving like amounts of other preparations high in oleic acid. Feed intake and rate of weight gain were similar in all groups.

concentrate)—yielded little or no differences in hypercholesterolemia and atherogenesis (Figs 34 to 36) (461, 476, 477). Similarly addition of a linoleic acid concentrate to a mash containing powdered egg yolk also failed to suppress hypercholesterolemia and atherogenesis. Comparable experiments of other investigators in rabbits and chicks generally—with few exceptions—yielded similar results (482–494 *cf* also refs 478–481). Further with cottonseed or corn oil at the 20 per cent level hypercholesterolemia and atherogenesis were markedly enhanced in cholesterol fed chicks (461, 476–477).

Similar extensive experiments with several oils and combinations of oils have recently been carried out by a research group utilizing rats fed cholesterol + cholic acid (495–496). Their data led them to the following interpretation considerably different from the trend of thinking extant concerning the findings in man. The non essential unsaturated fatty acids (oleic *et al*) promote hypercholesterolemia in rats on this regimen. In contrast hypercholesterolemia is counteracted by both saturated and essential fatty acids. The product dietary EFA (linoleic and arachidonic acids) times dietary saturated fatty acids has a high degree of negative correlation with the serum cholesterol values—and EFA and saturated fatty acids are equally active substituting for each other in inducing this anti hypercholesterolemic effect. It is suggested that these relationships may be valid for man. This hypothesis remains to be explored in further studies.

On the other hand in a rabbit experiment of different design utilizing a purified cholesterol free ration it was observed that highly saturated hydrogenated coconut oil (20%) induced hypercholesterolemia and atherogenesis whereas unsaturated safflower oil had no such effect (469 *cf* ref 477). Further as already noted 40 per cent fat in the form of 20 per cent corn oil plus 20 per cent Crisco failed to induce coronary thrombosis and myocardial infarction in cholesterol thiouracil cholate fed rats—in contrast to 40 per cent butter fat (462).

In calling attention to the apparently divergent results of experiments with unsaturated vegetable oils in man vs chick rat and rabbit it is essential to note the differences in experimental design. Except for a few studies in South African Bantus the human subjects investigated were individuals on American type high calorie high fat high saturated fat high cholesterol diets. In association with these habitual diets most subjects exhibited moderate or marked hypercholesterolemia. Most of the studies in such people were of the complete substitution type involving withdrawal of saturated animal fat and cholesterol and incorporation of unsaturated vegetable oil. These investigations were patently not analogous in design to the aforementioned animal experiments. The latter involved addition of cholesterol and various fats to the low fat low cholesterol rations of rabbits rats and chicks originally exhibiting low normal serum cholesterol levels.

It would seem that further experiments are essential in animals having a design similar to the human studies—including complete substitution, partial substitution, and supplementation experiments. Until these are accomplished, it would be premature to attribute the seemingly disparate findings in animals and man to “species differences.” It may yet be that in this new area of work, further experiments and deliberations concerning them will yield an intellectual advance effectively accounting for all the apparent discrepancies. In any case, should species differences actually be demonstrated, their mechanism becomes an important problem *per se* (1).

In an earlier section of this book, a discussion was presented of the current hypothesis that atherosclerotic disease may be a resultant of essential fatty acid deficiency. At this juncture, it is appropriate to deal with animal experimental evidence bearing upon this problem. Obviously, from the foregoing experiments, atherosclerosis was readily produced in chicks and rabbits by cholesterol feeding—even in the presence of large amounts of high EFA unsaturated vegetable oils. In fact, their presence in the diet was associated with enhanced hypercholesterolemia and atherogenesis. These results supervened, despite the fact that the experimental animals exhibited no tissue deficiency of polyethenoic essential fatty acids (Fig. 37) (493–494). Thus, they cast doubt upon the hypothesis that atherogenesis is a resultant of EFA deficiency.

Here it is important to recall the experimental conditions required for the production of true essential fatty acid deficiency in animals. As the original work over two decades ago demonstrated signs of FFA deficiency can be elicited only by accomplishing virtually complete removal of all fat from the diet (416–420, 497). The incorporation of only slight amounts of any natural fat (animal or vegetable) in the experimental ration rendered impossible the production of signs of EFA deficiency.

Recently it was shown that EFA deficiency was more readily and rapidly produced when a diet rigidly devoid of lipid was combined with other nutritional metabolic abnormalities *e.g.*, high cholesterol intake induced diabetes mellitus, or other stresses (498–503). These findings suggested that FFA need may be a function of the overall nutritional metabolic situation of the organism. This phenomenon of variability in the need for dietary essentials—dependent upon other nutritional metabolic factors—has also been demonstrated in experimental animals for such nutrients as methionine, vitamin B₁₂, folic acid, choline, tocopherols, and pyridoxine. This may be of considerable practical and theoretical importance particularly in view of data suggesting a role for the *c* metabolites in the regulation of cholesterol metabolism and atherogenesis (see below) (454, 455, 458, 461, 504–511, 513–539). These experimental findings lend in direct support to the imbalance hypothesis of atherogenesis (see above) (16, 19, 21, 22, 461). Moreover, they constitute animal experimental support for the conclusion—already presented above in the section on

clinical investigation—that if an EFA deficiency does in fact exist with ingestion of diets high in calories, total fats saturated fats, cholesterol, then it can only be a relative, secondary, derivative deficiency

6 *Corn Germ, Sitosterols, Dihydrocholesterol, Brain Extracts*—In one recent study on the effects of various corn oil preparations in cholesterol fed chicks it was demonstrated that incorporation of whole corn germ yielded highly effective inhibition of hypercholesterolemia and atherogenesis (486) This finding was confirmed in subsequent studies in our

Polyenoic fatty acids (EFA) in tissues of chicks fed cholesterol and peanut oil for 4 weeks




FIG 37—Dam H Kristensen G Kolod N Prange I and Sondergaard E (494) Note that the two groups receiving cholesterol + peanut oil had tissue levels of essential fatty acids greater than those of the control group Note also that the group receiving 1 per cent cholesterol without peanut oil had levels of tissue EFA similar to the controls Thus no evidence of reduction in tissue EFA or of EFA deficiency was noted in these cholesterol fed birds

laboratory (540) The mechanism of this phenomenon is apparently related to the lack of fat absorption of the whole germ Whole corn germ has its oil so enclosed that it is not available for absorption It therefore acts like a low fat diet

Several years ago it was demonstrated that the incorporation of sitosterols in the diets of cholesterol fed chicks and rabbits resulted in marked suppression of hypercholesterolemia and atherogenesis (Fig 38) (1, 541-552) Large amounts of plant sterols were necessary to achieve this effect

the ratio to dietary cholesterol being 1-7 1 This action was attributed to inhibition of intestinal absorption of cholesterol possibly due to formation of a poorly soluble mixed crystal of cholesterol and sitosterol (553-559) This conclusion was supported by the observation that sitosterols failed to reduce hormone induced endogenous hypercholesterolemia, resulting from administration of estrogens or hydrocortisone to chicks (552) Limited data indicated that sitosterols were also ineffective in speeding up the mobilization of previously accumulated tissue cholesterol *ie*, they were effective only prophylactically, but not therapeutically (560, 561)

EFFECTS OF SOY STEROLS IN CHOLESTEROL FED COCKERELS




FIG 38 Stamler J, Pick R and Katz L N (552) Note the virtually complete inhibition of hypercholesterolemia and atherogenesis in the group receiving plant sterols

Dihydrocholesterol was also shown to block diet induced hypercholesterolemia and atherogenesis in chicks rats and mice However when fed for long periods of time, this sterol was itself absorbed in sufficient quantities to produce elevated plasma levels and lesions This was also true for lanosterol and delta 4 cholestenone With the exception of one report these phenomena were not observed to occur to any significant degree with ingestion of plant sterols (562-571)

Results similar to those obtained with plant sterols were noted following the administration of defatted brain extract preparations (572-576)

7 *Nutritional Interrelationships in Experimental Hypercholesterolemia and Atherogenesis*—A highly significant development in experimental atherosclerosis research in recent years has been the undertaking of studies on the interrelated effects among cholesterol lipid and other nutrients (454, 455, 458, 461 504-511, 513-529, 537-539 577-597) For example it was demonstrated that hypercholesterolemia and atherogenesis were ap

parently aggravated in cholesterol oil fed chicks permitted to ingest only 60 per cent of their usual total feed intake, or subjected to intermittent starvation (Fig 39) (1, 577-578 also *cf* ref 510, 579)

Further, as already noted it was shown that hypercholesterolemia and atherogenesis could be successfully produced in monkeys and rats by utilizing a ration high in cholesterol lipid and inadequate in the sulfhydryl containing amino acid methionine (1, 453-458, *cf* refs 252-511, 513-516-521, 527, 537-580-582) This phenomenon was amply confirmed in experiments in chicks (Fig 40) (461, 504-509)

EFFECTS OF DIETARY RESTRICTION IN CHOLESTEROL-FED COCKERELS




FIG 39 Rodbard S Bolene C and Katz L N (577) Note the more marked hypercholesterolemia and the earlier emergence (19-24 weeks) of atherosclerotic lesions in the group receiving a diet of similar composition to the control but restricted in amount to 60 per cent of the control group's feed intake

Other studies showed that feeding a ration low in choline also aggravated the hypercholesterolemic and atherogenic effects of a high cholesterol high fat diet in rats (286-463-524-526-537-539) (In contrast to these findings with deficiency of methionine and choline experiments in recent years tended generally—with few exceptions—to confirm previous observations that supplementation of an adequate ration with lipotropic factors failed to suppress hypercholesterolemia and atherogenesis [1, 12-14])

In subsequent experiments it was shown that hyperlipemia hypercholesterolemia and atherogenesis were aggravated in cholesterol oil fed chicks ingesting a diet low in total protein. In one such study sucrose was incorporated in the diet at the expense of protein, so that total pro-


EFFECTS OF A METHIONINE-DEFICIENT DIET ON
CHOLESTEROL-FED COCKERELS

FIG 40 Stamler J Pick R. and Katz L N (505) The chicks on synthetic diet had a ration of purified constituents adequate in known nutrients. The experimental group (1 CO methionine deficient) had a similar ration except that a 50% protein of low methionine content was fed and supplementary methionine was excluded from the diet. Here and in subsequent figures body weight is the terminal weight in grams feed intake is in grams per bird per day. Note the more marked hypercholesterolemia and atherogenesis in the methionine deficient group compared with the group on an adequate synthetic ration. Note also the significantly less marked hypercholesterolemia and atherogenesis in the 1 CO synthetic group compared with the 1 CO control on commercial mash. This may be due to the higher protein content in the diet of the former (35 per cent vs 20 per cent).


Effects of addition of refined Carbohydrates to High Fat,
High-Cholesterol Chick Mash with and without reduction of Dietary Protein

FIG 41 Stamler J Pick R. and Katz L N (506) The third group—1 CO 45 per cent sucrose 20 per cent protein—received purified soy protein in order to increase the dietary protein to 20 per cent. Note the intensified hypercholesterolemia and atherogenesis in the group with reduced protein intake due to incorporation of 45 per cent sucrose in the atherogenic mash. Note that this was prevented by addition of protein.

tein was reduced from the usual 20-25 per cent to 10-15 per cent *. Another group of chicks received a similar high level of sucrose ("empty calories" from carbohydrate), but with purified soy protein added to maintain total protein at the 20 per cent level. Unlike the high sucrose low protein group, this high sucrose, adequate protein group behaved similarly to the control group (Fig 41) (504-506). These data indicated

EFFECTS OF A HIGH VITAMIN-HIGH PROTEIN SUPPLEMENT TO A CHOLESTEROL-OIL-REGULAR MASH DIET

Series 46-Weeks 9-16




FIG 42—Stamler J, Pick R, and Katz L N (505). The protein supplement was at the 33 per cent level by weight and included 7 per cent of each of the following: casein, defatted fish meal, soy protein, defatted liver, Brewer's yeast. In addition to the vitamins of the Brewer's yeast, individual vitamins were added to the mash at a level five times that used routinely in a synthetic ration. Note that the protein vitamin supplement inhibited hypercholesterolemia and atherogenesis.

that the protein content of the diet—not its sucrose content—was the decisive factor influencing the cholesterolemic and atherogenic response to the high fat, high-cholesterol intake. Recently, other workers reported that the quantity and type (glucose vs. sucrose) of refined carbohydrate in an experimental or human ration may influence the cholesterolemic response (516, cf. also ref. 372).

In other related experiments it was demonstrated that high protein, high vitamin supplementation suppressed hypercholesterolemia and atherogenesis in cockerels ingesting either a purified or commercial ration high in cholesterol and fat (Figs 42 to 44) (504-505-507). In studies on the mechanism of this phenomenon it was shown that vitamin supple-

*In the diets of most Americans, proteins supply 12 to 14 per cent of calories (22-23, 107, 197, 210-214, 238).

EFFECTS OF HIGH VITAMIN-HIGH PROTEIN VS HIGH PROTEIN ALONE IN CHOLESTEROL-FED COCKERELS S-47 Weeks 13-23




FIG 43 Stamler J Pick R and Katz L N (505) The vitamin and protein supplements were the same as those in the experiment illustrated by Fig 42. Note that combined high protein high vitamin supplementation was most effective in inhibiting hypercholesterolemia and atherogenesis. Note that high vitamin alone was ineffective and that high protein alone was partially effective.

Effect of adding Casein and Egg Albumen to Mash containing Butter + Cholesterol and Egg Yolk respectively -S-50 12 to 18 weeks




FIG 44 Stamler J Pick, R and Katz L N (507) Note that supplementation of both atherogenic rations with casein and egg albumen respectively had a significant inhibitory effect on hypercholesterolemia and atherogenesis. Note also that hypercholesterolemia and atherogenesis tended to be more marked in the group on 1 per cent cholesterol + 5 per cent cottonseed oil than in the cholesterol + butter and the egg yolk groups again demonstrating lack of inhibitory effects of unsaturated vegetable oils in chicks.

mentation alone was without influence (Fig 43) * On the other hand, high protein alone apparently suppressed coronary atherogenesis in a manner similar to the high protein high vitamin combination (Figs 43 and 44) However, high protein alone apparently had less definitive and clearcut effects than combined high protein, high vitamin supplementation These findings suggested that the high protein diet was the decisive factor responsible for the results but that high vitamin intake may have had an adjuvant, synergistic influence These results constituted the first demonstration of an anti atherogenic influence of dietary protein They were consistent with several concurrent independent reports on the effects of protein supplementation on hypercholesterolemia in cholesterol lipid fed animals (511, 513, *cf* also refs 510 514-516 520 580 582) †

In addition to the foregoing experiments on nutritional interrelationships, the last year or two witnessed the development of studies on the effects of minerals in animals on a high fat, high cholesterol diet Thus, evidence was obtained indicating that cobalt magnesium and vanadium may inhibit hypercholesterolemia and atherogenesis (518-520 584 585)

It should be emphasized that all these experiments on nutritional interrelationships represent analyses of the effects of specific nutrients (proteins vitamins carbohydrates minerals and other nutrients) in animals on high fat, high cholesterol diets By demonstrating significant influences of some of these nutrients (*e g* proteins) the experiments lend support to the imbalance hypothesis elaborated above At the same time their results reinforce the basic tenet of the nutritional metabolic cholesterol lipid lipoprotein theory of atherogenesis—*i e* that high fat, high cholesterol intake is the decisive nutritional aberration in the causation of hypercholesterolemia and atherogenesis

*Previous studies by our group had shown that addition of large amounts of vitamins B₁ and E had no effect on cholesterolemia and atherogenesis in chicks fed high fat high cholesterol diets Imbalance studies on the combined influences of various vitamin deficiencies in birds on similar atherogenic diets are currently in progress Other workers have reported that large doses of nicotinic acid (538 586-594) pyridoxine (595) or tocopherols (596) may lower serum cholesterol levels in man Work has also been done in guinea pigs on combined vitamin C deficiency and cholesterol feeding (528 583 597) and on B₁₂ B and pantothenate deficiency (517 521 523 529)

†A recent study indicated that varying the level of dietary protein had no effect on cholesterolemia in persons on a high fat diet (252)

Chapter 6

Hormonal and Other Endogenous Factors and Atherosclerosis

A The Key, But Not Exclusive Role, of Diet in Atherogenesis, Interrelationships Between Diet and Other Factors (Heredity, Physical Activity, Stress, Endocrine Function, Smoking, and the Like)

THE nutritional metabolic cholesterol lipid lipoprotein theory of atherogenesis therefore emphasizes the decisive—but not exclusive—role of diet in the causation of this disease. Since the development of disease in individuals is in fact influenced by an interplay between diet and other factors including endogenous factors, then the delineation of these factors becomes one of the decisive tasks of research in this field.

Extensive evidence has been available for many years—particularly from clinical research—concerning the atherosclerosis potentiating effects of diabetes mellitus, hypothyroidism, renal disease and essential familial hypercholesterolemic xanthomatosis (1). Similarly, it has been abundantly demonstrated—by epidemiologic, clinical, pathological and animal experimental work—that hypertension markedly increases the risk of atherosclerotic disease, particularly when the prerequisite derangement in cholesterol lipid lipoprotein metabolism is also operative (Figs. 18 and 48) (1-5, 12-14, 21-23, 69, 71, 90, 96, 107, 112, 125, 126, 134-137, 143, 196-203, 208, 209, 228-230, 260-272, 278, 294, 314, 315, 436, 444-448, 461, 632, 647-650).

It is essential to re-emphasize that we attribute a *key*—but not exclusive—role to diet. We in no way imply that atherosclerosis is purely and simply a dietary disease. Such a viewpoint is definitely at variance with well established facts and therefore quite untenable. Thus the fact that many individuals eating a typical American diet for decades reach old age without clinical evidence of atherosclerotic disease—this fact alone refutes any attempt to imply a one to one cause and effect relationship between diet and disease. Further, the marked sex differential—the high incidence of coronary disease in middle aged American males in contrast to females—almost certainly contravenes any pure and simple dietary concept. Similarly, the greater incidence of atherosclerotic dis-

ease in persons with diabetes, nephrosis, hypothyroidism or hypertension contradicts any exclusive dietary concept. Actually, no serious scientist is making any such futile attempt to "fit" all the phenomena of atherosclerotic disease into the confines of a pure and simple dietary "explanation."

Obviously, as extensive clinical and experimental data demonstrate, the nature of the organism plays an important role in determining the effects of life span ingestion of a potentially atherogenic diet. Hence,

Relative mortality among insured persons reporting two or more cases of early Cardiovascular Renal diseases in their families




FIG 45 Lew E A (183) These data indicate that familial factors possibly hereditary and genetic influence the occurrence of atherosclerotic disease in individuals from population groups ingesting an habitual diet high in total calories, total fats, saturated fats, cholesterol.

different individuals ingesting similar habitual diets exhibit different levels of cholesterolemia. In these individuals many factors may interact significantly with diet—genetics and heredity, metabolism and endocrinology, clotting mechanisms, previous medical history, psychological make-up and physical condition (fatigue, stress, tension, frustration and other conditions), environmental factors (smoking, habitual physical activity, work, urbanization and other factors) (Figs 45 and 46) (1-6, 10-14, 23, 71, 113, 148, 149, 159, 202, 275, 300, 322, 446, 451, 598-623, 625-645, 785). Antecedent allergic or infectious disease, for example—possibly by damaging vessels and setting up sites of predilection for atherogenesis—may be of great importance in some cases. Hypertension,

hypothyroidism, renal dysfunction, diabetes, certainly tend to potentiate atherogenesis (1) The evidence is considerable that estrogenic secretion in women has an opposite effect (see below) (645, 646)

In recent years, great interest has been aroused concerning the possible influences of psychological stress and physical activity on the development of atherosclerotic disease, particularly coronary heart disease Up

*Distribution of excess deaths by disease
among men with a history of regular
cigarette smoking*




FIG 46 Hammond E C and Horn D (623-624) A total of 187,183 men aged 50-69 in 1952 were followed with almost 12,000 deaths occurring in 5 years The age adjusted death rate for all causes of the regular cigarette smokers was 68 per cent higher than that of the group who never smoked with a total of 2,665 excess deaths in the smokers Of these 2,665 excess deaths 1,388—52.1 per cent—were due to coronary artery disease The coronary disease mortality rate of these regular cigarette smokers was 70 per cent higher than that of those who never smoked These data suggest that smoking increases the risk of coronary disease in middle aged males habitually ingesting a diet high in calories total fats saturated fats cholesterol (Compare the findings of other recent epidemiologic studies—ref 2 also data on smoking and lipids-lipoproteins—ref 626 and a discussion of the whole problem of the interrelationship among diet smoking and coronary artery disease—ref 113)

to the present few data have been reported concerning the first of these (1-6 11-14 23 71 113 300 451 607-615 617-622 636-638 785)—an understandable fact in view of the complex and difficult problems of objective mensuration

Data were reported from Great Britain concerning physical activity Comparisons were made of sub groups within relatively homogeneous

socio-economic strata of the British population. The results indicated that the incidence of coronary heart disease tended to be lower in the subgroups habitually engaged in greater physical activity at work. Data along these lines were obtained in comparisons of sedentary bus drivers vs active bus conductors, sedentary telephone operators vs active postmen and other

Incidence of Coronary disease in active vs inactive workers Great Britain




FIG 47 Morris J N Heady J H Raffle P H B Roberts C G and Parks J W (148) These data indicate that the incidence of coronary disease (nonfatal and fatal) is lower in active than inactive workers in Great Britain i.e. in a population ingesting an habitual diet moderately high in total calories total fats saturated fats cholesterol

similar comparisons (Fig 47) (25 148 598). However further studies by the same British group tended to cast doubt upon the significance of these observations (149). Moreover all strata of the population in Britain as in the United States and Norway—irrespective of socio-economic status and habitual physical activity of work—were found

hypothyroidism, renal dysfunction, diabetes, certainly tend to potentiate atherogenesis (1) The evidence is considerable that estrogenic secretion in women has an opposite effect (see below) (645, 646)

In recent years great interest has been aroused concerning the possible influences of psychological stress and physical activity on the development of atherosclerotic disease, particularly coronary heart disease Up

*Distribution of excess deaths by disease
among men with a history of regular
cigarette smoking*




FIG 46 Hammond E C and Horn D (623-674) A total of 187,183 men aged 50-69 in 1952 were followed with almost 12,000 deaths occurring in 5 years The age adjusted death rate for all causes of the regular cigarette smokers was 68 per cent higher than that of the group who never smoked with a total of 2,665 excess deaths in the smokers Of these 2,665 excess deaths 1,388—52.1 per cent—were due to coronary artery disease The coronary disease mortality rate of these regular cigarette smokers was 70 per cent higher than that of those who never smoked These data suggest that smoking increases the risk of coronary disease in middle aged males habitually ingesting a diet high in calories total fats saturated fats cholesterol (Compare the findings of other recent epidemiologic studies—ref 2 also data on smoking and lipids-lipoproteins—ref 626 and a discussion of the whole problem of the interrelationship among diet smoking and coronary artery disease—ref 113)

to the present few data have been reported concerning the first of these (1-6 11-14, 23 71 113 300 451 607-615 617-622 636-638 785)—an understandable fact in view of the complex and difficult problems of objective mensuration

Data were reported from Great Britain concerning physical activity Comparisons were made of sub groups within relatively homogeneous

atherogenesis diet is the decisive factor. Thus, it would seem valid to designate diet as virtually a *sine qua non* for atherosclerosis and atherosclerotic disease even as the tubercle bacillus is a *sine qua non* for tuberculosis—although patently a multiplicity of host and environmental factors (nutrition and other socio economic variables) enter into complex interrelationship with the bacillus to influence the developmental patterns of the disease. For without the habitual ingestion of a potentially atherogenic diet, clinical atherosclerotic disease would be rare among the middle aged members of any population, irrespective of the operation of other potentially offensive factors.

INTERRELATIONSHIPS BETWEEN HYPERTENSION AND ATHEROGENESIS IN COCKERELS




FIG 48 Stamler J, Pick R and Katz L N (647) Stamler J and Katz L N (648). Note that hypertension failed to induce atherosclerosis in normocholesterolemic cockerels on a non atherogenic diet. Note also that in this experiment a diet of $\frac{1}{4}$ CO for only 15 weeks (cf Figs 32 and 39) leading to moderate hypercholesterolemia failed to induce gross lesions in the thoracic aorta. However the combination of blood pressure elevation plus the $\frac{1}{4}$ CO diet resulted in frank atherosclerosis.

It should be noted that this conclusion has as its cornerstone the concept that the pathological lesion atherosclerosis is the prerequisite for clinical coronary heart disease in at least 90 per cent of cases (1, 290, 291, 292, 294, 299, 300). As is well known however morphologic atherosclerosis frequently exists without clinical disease. What factors are responsible for the qualitative change from occult (pre clinical) to clinical disease? What are the causes of thromboses on plaques or hemorrhages into plaques leading to infarction? Unquestionably, part of the answer is quantity—the severity and extent of atherosclerosis (1, 290, 291, 292, 294, 299, 300). But it is unlikely that this is the whole answer. Thus as already indicated diet may play a role here as well e.g. the fat content of the diet via its possible tendency to increase blood coagulability.

to manifest high coronary heart disease incidence rates (2, 6, 10, 19, 21-23, 71, 125, 126, 134-137, 143, 150-154, 160, 172-180, 196, 197, 199, 208, 209, 228-230, 232-237, 260-273, 278)

Correspondingly, recent observations indicate that epidemiologic patterns of cholesterolemia and atherosclerotic disease were more decisively influenced by diet than by level of physical activity at work (see foregoing figures). Thus in South African Bantu—with a habitual diet low in total fats, saturated fats, cholesterol—the level of cholesterolemia and the incidence of coronary disease were found to be low in both sedentary and active workers (Fig. 4) (4-6, 21-23, 71, 72, 91, 107, 119-121, 123). Similar observations were also reported from Japan and Italy (Fig. 8) (72, 91, 107, 116-118, 133). Contrariwise, in Sweden and Finland—with a habitual diet high in total calories, total fats, saturated fats, cholesterol—level of cholesterolemia and incidence of coronary disease were observed to be high in both sedentary and active workers (72, 107, 157-160). Similarly, recent studies in the United States consistently revealed a high level of cholesterolemia and a high prevalence and incidence of coronary disease in all sectors of the middle aged male labor force, irrespective of nationality, socio economic status, income, occupation, physical activity of work, urban or rural residency (1, 2, 6, 10, 12-14, 21-23, 71, 72, 105-109, 111, 112, 125, 126, 133-137, 143, 172-180, 196, 197, 199, 208, 209, 228-230, 232-237, 251, 260-273, 278-281, 645).

These data impressively support the concept that diet is the decisive variable determining the pattern of atherosclerotic disease in a *population*. Nonetheless, studies on the interrelationships among nutrition, physical activity, atherosclerosis and atherosclerotic disease—particularly in countries where large segments of the population ingest high calorie, high fat, high cholesterol diets and lead a relatively sedentary life—are still limited. Recent studies indicate that physical activity may prevent an increase in serum cholesterol, under circumstances when increased fat is ingested to meet greater caloric need (322, 599, 639). It has also been suggested that continued physical activity through middle age may be a possible factor in the prevention of atherosclerotic disease (640). Further work is therefore certainly indicated on the effects of large muscle activity in persons and groups ingesting potentially atherogenic diets.

With respect to the individual, therefore, the relationship between diet and other factors may be properly formulated as follows, based on our present knowledge. In the individual member of a population group habitually ingesting a potentially atherogenic diet, the development of morphologic atherosclerosis and atherosclerotic disease is influenced by multiple factors. There is a complex interplay between diet and other factors which operate to accelerate or to retard atherogenesis. Thus, it is not a matter of one factor to the exclusion of others; it is not a matter of either or. But, among these multiple factors interacting to influence

—also stimulated concern with this field of research. This entire area is at present in process of active development including exploration of its therapeutic implications (see below)

C Thyroid Hormone

Of the endocrine glands attention has been focussed for years on the thyroid. The original basis for this concern was the clinical pathological observation that cholesterolemia and atherogenesis were increased in hypothyroidism and decreased in hyperthyroidism (1, 688). These findings were supplemented with extensive experimental evidence in cholesterol fed rabbits, chicks and dogs of an interrelationship among thyroid hormone, cholesterol metabolism and atherogenesis (1, 3, 687). Recent years have witnessed only limited activity in this area of research on atherosclerosis (see below).

D Estrogens

The influence of ovarian estrogenic secretion has been a particular center of contemporary research attention (1, 3-5, 12-22, 461, 645, 646, 689). This problem has been one of the major foci of our group during the past 8 years.

1 *Background*—Two key sets of facts concerning human coronary heart disease served as a basis for this work. The first was the finding that groups of middle aged males with manifest coronary disease had significantly raised levels of both plasma cholesterol and cholesterol/phospholipid (C/P) ratio (Fig. 19) (1, 3-5, 12-14, 21, 22, 69, 71, 171, 202, 219-281, 283, 301-318, 645, 689). The question arose whether one or the other of these changes played a more important role in atherogenesis.

A second major stimulus was the demonstration of a marked human sex differential in susceptibility to coronary heart disease in the middle aged populations of the economically more developed countries (1-5, 12-23, 71, 90, 134-137, 143, 155, 172-187, 196-206, 208, 209, 294, 461, 645, 646, 689). Numerous clinical and epidemiologic studies demonstrated a remarkable resistance of young and middle aged women to coronary heart disease—an immunity tending to disappear in the decades following the menopause. At the time these investigations began, no animal experimental data were extant accounting for this highly significant phenomenon. It was a scientific enigma.

2 *Prophylactic Experiments with Estrogens*—Attention was first focussed on the differences in patterns of hyperlipemia in chicks resulting from cholesterol feeding versus estrogen administration. The former invariably produced a predominant disproportionate rise in plasma cholesterol, a smaller increment of plasma phospholipids, resulting in markedly elevated C/P ratio. Estrogens on the other hand caused a manifold increase of phospholipids, a considerably less marked elevation of choles-

and thereby perhaps enhance risk of thrombogenesis (assuming that blood coagulability is related to pathologic thrombosis) (326-344) Nervous and hormonal factors—related to stress and strain in the individual—may exert an influence, possibly by affecting lipid metabolism, and/or coagulability possibly by increasing the risk of sudden arrhythmic (non thrombotic) death in cases where coronary atherosclerotic disease has led to ischemia and increased myocardial irritability (607-622)

Undoubtedly, this problem of factors precipitating acute episodes is highly complex and important, *but a problem secondary to the decisive one, atherogenesis*

B Blood Clotting and Fibrinolytic Mechanisms

Another important development of recent years has been the extended exploration of the relationships among anticoagulants blood clotting mechanisms, lipid metabolism atherogenesis and thrombus formation It is, of course a well established fact that thrombotic episodes are in a major way responsible for the conversion of subclinical atherosclerotic pathology to clinical disease *e.g.* coronary or cerebral thrombosis with infarction The importance of this problem resulted in extensive studies with anticoagulants soon after their biochemical identification This line of research received further impetus with the demonstration of the profound effects of heparin on cholesterol lipid lipoprotein metabolism (1 651-661 674 675) It was shown that this biological product has a remarkable ability to clear lipemic serum Heparin apparently activates a plasma enzyme system (a lipoprotein lipase) capable of markedly altering the spectrum of plasma lipoproteins Evidence has been presented suggesting that endogenous heparin may play a role physiologically in lipid metabolism and that this mechanism may be abnormal in atherosclerotic subjects (662-673)

Based on these findings experimental work was done on the effects of heparin on cholesterol induced hyperlipemia and atherogenesis in chicks and rabbits Limited inconsistent data suggested that heparin may suppress these phenomena presumably by virtue of its effects on lipid metabolism (676-684) Similar findings with dicumarol were also reported (1, 685) On the other hand an extended series of experiments in our laboratory consistently yielded negative results In fact the evidence indicated that both heparin and dicumarol may potentiate atherogenesis under certain experimental conditions (686)

Recently the additional observation was reported that ingestion of high fat meals induced a decreased fibrinolysis and an increased blood coagulability, measured *in vitro* (326-345) This finding further enhanced research interest in interrelationships among anticoagulants lipid metabolism blood coagulability fibrinolysis atherogenesis and thrombogenesis Correspondingly, the thrombogenic theory of atherogenesis (499, 450)—alternative to the dietary nutritional cholesterol lipid lipoprotein theory

that estrogen induced inhibition of coronary lesions may be closely interlinked in individual birds with maintenance of C/P ratios at or near normal levels (Fig 50) (1 645, 689-691)

Finally this initial study—by demonstrating that estrogens were capable of preventing experimental coronary atherosclerosis in cockerels—for the first time brought forward experimental evidence supporting the hypothesis that the sex difference in human susceptibility to coronary disease may be at least in part a resultant of ovarian hormonal secretion

EFFECTS OF ESTROGENS ON PLASMA LIPIDS AND CORONARY ATHEROGENESIS IN CHOLESTEROL FED COCKERELS




FIG 50 Pick R Stamler J Rodbard S and Katz L N (691) Note that all birds free of lesions had cholesterol/phospholipid (C/P) ratios less than 0.80 \pm or near normal values

Subsequently it was demonstrated that oral administration of mixed conjugated equine estrogens was equally effective in producing feminization altered plasma lipid patterns and prevention of coronary atherogenesis. Estrogen effects on plasma lipids and lipoproteins were also further delineated (1 645 689). Curiously estrogens apparently effected a significant reduction of α lipoproteins in cockerels a change precisely opposite to that induced in man (see below). Estrogen treated cholesterol oil fed chicks also exhibited apparent decreases in plasma low density β lipoproteins (of the S_{120} to 400 class) and in plasma esterified/total cholesterol ratios.

Several synthetic estrogens given at oral or parenteral dosage levels adequate to accomplish feminization also maintained normalization of C/P ratios and prevented coronary atherogenesis in cholesterol fed

terol, so that C/P ratio was significantly *depressed* below normal (Fig 49) (1, 645 689-691)

In the initial experiments, which were of the "prophylactic" type, estrogens were administered to cockerels simultaneously transferred to an atherogenic diet. Estrogens in these cholesterol fed cockerels did induce a marked enhancement of phospholipemia with consequent maintenance of C/P ratios at or near normal levels (Fig 50). The hormone also fem

Effects of Cholesterol Oil feeding vs Estrogen administration on plasma lipids of cockerels




FIG 49 Stamler J Pick R and Katz L N (689) Stamler J and Katz L N (690) Pick R Stamler J Rodbard S and Katz L N (691). Note the predominant hypercholesterolemia in the cholesterol fed group with increase in the cholesterol/phospholipid (C/P) ratio. Note the predominant hyperphospholipemia in the estrogen treated group with lowering of the C/P ratio.

inized the cockerels as indicated by their combs and other secondary sex characteristics. Gross examination after several weeks revealed the presence of marked atherogenesis in the *aortas* of both the estrogen treated and control birds. However histological studies showed that the estrogen treated birds in contrast to the controls were practically free of lesions in the *coronary* arterial tree (Fig 50). This was an experimental demonstration of segmental differences in atherogenesis. This observation, that different laws govern atherogenesis in different vascular beds, was supported by data on human material. The data further suggested

and fibroblasts. It was such relatively "young" plaques that estrogens completely reversed despite continued feeding of the atherogenic diet. So thoroughgoing was this estrogen "therapeutic" effect that special histological and histochemical studies revealed no deviations from normal (other than rare residual lipid infiltration) in the architecture of these coronary vessels—nor could any alterations be demonstrated in the ground substance (645-693).

This experiment proved once again that atherosclerotic plaques are, within wide limits partially and even wholly *reversible*. Finally, this experiment—and particularly the special histochemical study, accomplished in cooperation with our Argentine colleagues—yielded preliminary evidence suggesting that the estrogen anti atherogenesis may perhaps be related at least in part, to a reticulo endotheliosis.

EFFECTS OF OVIDUCT LIGATION IN MATURE EGG PRODUCING ESTROGEN-SECRETING HENS FED A CHOLESTEROL-OIL MASH




FIG. 52. Stamler J. Pick, R. and Katz L. N. (695). Note that oviduct ligation resulted in similar levels of cholesterolemia in the male and female birds. Note that both groups of hens had low C/P ratios reflecting estrogen induced hyperphospholipemia. Note the minimal coronary atherogenesis in the hens. In this experiment aorta atherogenesis was slightly less marked in the hens than in the roosters but the difference was not significant.

4 *Effects of Endogenous Estrogen Secretion—Sex Differential in Chickens*—When sexually immature male and female chicks 8 to 15 weeks old were fed an atherogenic diet both sexes exhibited significant aorta and coronary atherogenesis—no sex differential in coronary atherogenesis was observed (1-645-694).

On the other hand egg laying hens in contrast to roosters of the same age were found to be remarkably resistant to cholesterol induced coronary atherogenesis, although both sexes were susceptible to aorta atherogenesis (Fig. 52) (645, 689, 695).

cockerels. On the other hand, studies with several synthetic estrogen analogues of low feminizing potency invariably yielded negative results (645).

3 *Therapeutic Experiments with Estrogens—Reversibility of Lesions*—In subsequent "therapeutic" experiments it was shown that estrogens were able to reverse cholesterol induced plaques in the coronary vessels of chicks. In fact, this anti atherogenesis supervened even when the atherogenic diet was continued during hormone administration (Fig 51).

ESTROGEN INDUCED REVERSAL OF CORONARY ATHEROSCLEROSIS IN CHOLESTEROL-FED COCKERELS




FIG 51 Pick R, Stamler J, Rodbard S and Katz L N (692). A third group not represented in this figure was sacrificed at 16 weeks of age after 8 weeks on diet and the presence of extensive coronary lesions prior to onset of estrogen therapy was thereby confirmed. Note the marked reduction in coronary atherogenesis in the estrogen treated cockerels despite continued feeding of the potentially atherogenic diet.

(1, 645, 689, 692). Estrogens reversed both the lipid and the fibroblastic components of these atherosclerotic plaques. This does not imply that estrogens were effective in removing areas of marked hyalinization, advanced fibrosis, calcification or bone formation. Such advanced lesions were not present in the control birds sacrificed after 8 weeks on the atherogenic diet. Rather, their coronary vessels showed plaques composed only of actively proliferating fibroblastic tissue with moderate collagen deposition, plus lipid and cholesterol situated extracellularly and in lipophages.

Estrogens were also combined with pancreatectomy, with administration of corticoids including desoxycorticosterone acetate (DCA) cortisone, hydrocortisone and ACTH. Dosages of the latter were adequate to produce one or another manifestation of hyperadrenocorticism. Hydrocortisone induced frank steroid diabetes (1 12-14, 645 698). Neither pancreatectomy nor corticoids had any counteracting influence on estrogen anti atherogenesis (1 12-14, 645 698).

EFFECTS OF ESTROGEN PLUS ANDROGEN IN CHOLESTEROL FED COCKERELS




FIG 54—Stamler J, Pick R and Katz L N (697). Note in the birds receiving estrogen + androgen the masculinized comb index, the maintenance of estrogen effects on C/P ratio and on coronary atherogenesis.

More recently, estrogen administration to cockerels was coupled with induction of hypothyroidism by thiouracil feeding, thyroidectomy or thyroidectomy plus I^{131} administration. Hypothyroidism was associated with significant loss of the ability of estrogens to protect the coronary vessels against atherogenesis (Fig 55) (645 687). It was associated with a tendency to higher C/P ratios. These findings are intriguing in view of the well known association among hypothyroidism, hypercholesterolemia and atherogenesis in human beings.

Finally, a combination of insulin plus estrogens in cholesterol-fed cockerels was undertaken because insulin counteracted the regression of atherosclerotic lesions usually occurring in cockerels following transfer from an atherogenic to a normal diet (645 689). The administration of the pancreatic hormone apparently counteracted the ability of estrogens to protect the coronary vessels against cholesterol-induced atherogenesis (Fig 56). Further work is proceeding on this phenomenon.* This finding

Very recently, we have found that a combination of estrogen and a low protein diet (with or without added insulin) led to ulceration (and thrombosis and dissecting aneurysm) of the aortic lesions in the male chick (786). Still more recently, such ulcerations were found to occur in the coronary vessels as well (787). Insulin addition was not essential for this ulceration to develop; only estrogen and a low protein diet. Further ulceration was found to occur during the healing stage of the lesions (787). Thus complications of the basic atherosclerotic process have been produced experimentally.

However, the role of egg laying in the disposal of excess cholesterol lipid required analysis since cholesterol fed hens had somewhat lower plasma total cholesterol levels than their rooster counterparts (Fig 52). Surgical ligation of the oviducts was therefore accomplished so that mature ova were deposited into the abdominal cavity, and subsequently reabsorbed. Such a group of oviduct ligated, egg producing cholesterol fed hens exhibited plasma cholesterol levels similar to those of an age and diet matched group of roosters. These hens—while exhibiting marked aorta atherogenesis—remained almost completely free of coronary lesions (645, 695).

**EFFECTS OF OVARIECTOMY ON PLASMA LIPIDS
AND ATHEROGENESIS IN CHOLESTEROL-FED HENS**
S 37 + 43, Age 33-44 Weeks




FIG 53 Pick R Stamler J and Katz L N (696). Note that ovariectomy resulted in aggravated hypercholesterolemia, elevation of the C/P ratio, and marked coronary atherogenesis.

Finally, surgically ovariectomized hens, completely lacking endogenous estrogen secretion, did in fact develop grossly elevated C/P ratios and extensive coronary atherosclerotic plaques in response to cholesterol feeding (Fig 53) (645, 689, 696). Ovariectomy had eliminated their immunity to cholesterol-induced coronary atherosclerosis. Therefore, the resistance of mature egg-producing hens to coronary atherogenesis is a product of ovarian function, particularly estrogen secretion.

5 Effects of Estrogens Combined with Other Hormones—The first experiment of this type utilized estrogen plus androgen in a 1:3 and 1:100 ratio (Fig 54) (1, 12-14, 645, 689, 697). With this combination, masculinization of the major secondary sex characteristics of cockerels was readily maintained. Concomitantly, these birds exhibited full estrogenic effects on plasma lipid patterns and coronary atherogenesis. Thus, one of the triad of estrogen effects, namely feminization, was counteracted while the other two continued to operate.

an enhancement of hyperphospholipemia so that C/P ratios rose less markedly than in non-estrogen treated animals (645, 700 *cf* also refs 701, 702) These estrogen treated rats exhibited significantly less coronary atherogenesis compared with the controls

In our laboratory, studies of similar design were carried out in the herbivorous rabbit These experiments consistently yielded negative results irrespective of the variations in the level of cholesterol feeding and the route and amount of estrogen administration (Fig 57) (645, 689, *cf* refs 321 703) Estrogens in these rabbits failed to enhance hyperphospholipemia, so that C/P ratios were markedly elevated in contrast

EFFECTS OF ESTROGEN IN CHOLESTEROL-FED RABBITS - SERIES 1-4




FIG 57 Stamler J Pick, R and Katz L N (689) Note in the estrogen treated rabbits the absence of any definitive hormonal effect on C/P ratio or atherogenesis

to the estrogen effects in chicks and rats Is there any cause-and effect relationship between estrogen failure to enhance hyperphospholipemia and failure to render the coronary vessels of rabbits immune to atherosclerosis?

7 *Estrogens Plasma Lipids Lipoproteins and Atherogenesis in Man* — During the years our department was unfolding the foregoing program of research on estrogens and cholesterol induced coronary atherosclerosis in chicks significant findings were published by other investigators on possible interrelationships among estrogens plasma lipids-lipoproteins and coronary atherogenesis in man (1 4 5 12-14 645, 646, 704-719) Thus it was established that estrogens exert a profound effect on human plasma

provoked the questions Is insulin one factor in the pathogenesis of the frequent, premature, severe atherosclerosis of diabetic patients? Is the diabetic woman's loss of "immunity" to coronary atherosclerosis a resultant, to any degree, of the effects of exogenous insulin?

6 *Effect of Estrogens in Mammalian Species*—Recently experiments were reported on rats maintained for 18 weeks on a regimen of estrogens plus an atherogenic diet Like chicks rats responded to estrogens with

THIOURACIL COUNTERACTION OF ESTROGEN ANTI ATHEROGENESIS IN CHOLESTEROL-FED COCKERELS




FIG 55 Pick R Stamler J and Katz L N (687) Note in the group receiving estrogens + thiouracil the tendency for C/E ratio to be higher and the more marked coronary atherosclerosis compared with the group receiving estrogen alone

INSULIN COUNTERACTION OF ESTROGEN ANTI ATHEROGENESIS IN CHOLESTEROL FED COCKERELS




FIG 56 Stamler J Pick R and Katz I N (689) Note in the group receiving estrogens + long acting insulin (10-15 units/bird/day) the hypoglycemia and the more severe coronary atherosclerosis compared with the group receiving estrogens only

Chapter 7

Definitive Approaches to Prophylaxis and Therapy of Atherosclerotic Disease in Man

A The Meaning of *Definitive Approaches*

By definitive approaches to prophylaxis and therapy are meant those which will retard, arrest and even reverse atherosclerosis and its complications. They are therefore, approaches for successful specific prevention or treatment of the clinical disease (21, 645, 720, 721). These approaches are inevitably long term in nature.

B Criteria for Evaluating Prophylactic and Therapeutic Regimens

What are the criteria for evaluating proposed prophylactic and therapeutic regimens? The critical and decisive criteria are not the effects of a given regimen on symptoms, weight, plasma cholesterol or lipoprotein levels, angina pectoris and the like. These unfortunately are not adequate, reliable criteria to judge efficacy of prophylaxis or therapy. The essential problem is one of halting or at least retarding the progression of the basic atherosclerotic disease—or even reversing the process in whole or in part—with a resultant postponement or elimination of clinical episodes. The only reliable criteria, therefore, are effects on occurrence and recurrence of clinical episodes of atherosclerotic disease (coronary, cerebral and others) and above all effects on survival, as measured by the duration of life.

If these are the decisive, meaningful criteria, then many of the currently proposed approaches to prophylaxis and therapy are *at this juncture* based on inference, e.g. the inference that lowering serum cholesterol may be effective in mitigating occurrence and recurrence of atherosclerotic disease. For practically all of the proposed prophylactic and therapeutic regimens the validity of this inference has not yet been unequivocally demonstrated. They are still experimental. They are not yet components of the armamentarium of medical practice with proved and established status.

One must not be misled by the influence of a proposed regimen on plasma cholesterol. This is far more easily determined than effects on atherosclerotic disease. The former involves studies taking only weeks

lipid and lipoprotein patterns. In particular, estrogens were found to increase plasma phospholipids, occasionally lower cholesterol, reduce C/P ratios, elevate α lipoprotein, and increase the ratios α/β lipoprotein and α/β lipoprotein cholesterol. Comparisons of the plasma lipid lipoprotein patterns in normal males, normal females, and males with clinical coronary disease lent further support to the concept that estrogens may play a role in the 'immunity' of premenopausal women to coronary atherosclerosis.

During these years, four relevant studies were also published from clinico-pathologic laboratories. The first reported that young ovariectomized women, examined at autopsy several years after surgery, exhibited a significant loss of 'immunity' to coronary atherogenesis normally present in intact women of their age (713). The second study yielded similar findings. In addition, it reported that elderly males receiving prolonged high dose estrogen treatment for prostatic carcinoma, exhibited significantly less coronary atherosclerosis than a matched control group receiving little or no hormone. Finally, it presented data indicating that women with breast carcinoma, who presumably were hyperestrogenic, also had less coronary disease than their controls. It suggested that these changes might be attributable to specific anti-atherogenic effects of estrogens (714). The third study presented data indicating that men and women with myocardial infarctions tended to excrete less biologically active estrogens and 17 ketosteroids than their controls (715). The fourth investigation confirmed that eunuchs manifested less coronary atherosclerosis than non-castrate men (717-718). Correspondingly, castrate men had higher serum α lipoprotein levels, higher α/β lipoprotein ratios, lower β lipoprotein and cholesterol concentrations. These patterns resembled those seen in women or following estrogen administration or androgen withdrawal. Again, it was suggested that these estrogen-induced alterations in serum cholesterol, lipid, lipoprotein metabolism might be responsible for the relative freedom of the estrogen-dominated eunuch from coronary atherosclerosis.

In concluding this section on hormonal and other endogenous factors and atherosclerosis, one general point merits re-emphasis. These studies dealt with the effects on the organism of an *interplay between potentially atherogenic diets and endogenous factors*. This was true for investigations in both human beings and experimental animals. Without a nutritionally induced potential for atherogenesis, significant lesions were not observed irrespective of the endogenous factors operating in the organism.

keeping it reduced is therefore, clearly established as beneficial to all persons subject to atherosclerosis. It is something every physician should advocate. The diet prescribed obviously should have adequate amounts of the essential nutrients.

Life Insurance Co data on effects of Overweight and of weight reduction on Mortality experience




FIG 58 Shepard W P and Marks H H (285). Note that in the groups undergoing weight reduction relative death rates tended to be similar to those of the normal weight standard risk groups in contrast to the 47 per cent higher rates in the obese groups.

2 Low Fat Low-Cholesterol Diets—Three reports have been published on the therapeutic effects of low fat low-cholesterol diets in patients with established coronary heart disease (366-368, 372, 373, 722, 723, cf also ref 724)*. In the first of these two groups of patients with proved myocardial infarction were followed for 8 years (366). A low fat low cholesterol diet—containing approximately 1 600 calories daily, 20 to 25 grams fat, 50 to 70 mg cholesterol, 225 grams carbohydrate, 120 grams protein plus supplemental nutrients (Brewer's yeast and whole wheat germ)—was prescribed for one group of 50 subjects. The other 50 continued to ingest their habitual diets. The treated group experienced a significant weight loss occurring during the initial 3 years thereafter.

*Several recent publications present detailed practical information on diversified pleasant low fat low-cholesterol diets (369, 372-374, 722, 723).

or months the latter requires at least five years of study. The former is an essential first step, but only a first step. The latter is massive, expensive, difficult, and a long term undertaking, hence its comparative rarity. More such are needed.

C Precautions in Evaluating Prophylactic and Therapeutic Regimens

What are essential precautions in the evaluation of possible prophylactic and therapeutic regimens? The critical problem is one of control. The number of subjects under investigation must be adequate to yield a statistically 'clean' answer at the medically anticipated level and in an anticipated time (e.g. a 50 to 75 to 90 per cent reduction in 4 year mortality rate). A matched placebo group—set up on a "double blind" basis—is virtually an inevitable and vital necessity. 'Matched' should mean by age sex race ethnic origin occupation income socioeconomic status, place of residence by height weight body build. In therapeutic studies, "matched" should mean by number time location severity of episodes of clinical atherosclerotic disease diagnosed according to uniform accepted criteria. 'Matched' should further mean by cholesterol lipid lipoprotein levels by other diseases related or unrelated particularly hypertension other heart or cerebral diseases renal or thyroid or hepatic diseases diabetes mellitus. 'Matched' should mean optimal uniformity in procedures for both treated and placebo groups e.g. apparently similar approaches to dietary advice to prescription of medication visits to the physician adjuvant therapy and other procedures.

A most important additional precaution is the institution of adequate measures to assess adherence. Investigators should have accurate objective criteria for evaluating whether subjects are taking pills sticking to diets and so forth. Finally satisfactory methods are essential for follow up of cases who drop out of the study (lost patients) and for determination of cause of death (with autopsy verification insofar as possible). These complex precautions are not easily effectuated. Several studies recently published fall short of the mark in one or another of these aspects.

D Specific Proposed Prophylactic and/or Therapeutic Regimens

Let us evaluate specific prophylactic and therapeutic regimens currently under investigation on the foregoing basis. With but a single exception no adequate long term prophylactic studies of any kind have been published.

1 *Obesity Correction*—The exception is a recent report from life insurance sources on the prophylactic effects of correcting obesity. The data indicate that grossly obese individuals—with much higher risks of death due to coronary disease—experience a definite lowering of risk following weight reduction (Fig 58) (284-285). Reducing weight and

infarction recurrence rate and fatality rate (2.6 per cent vs. 10 per cent mortality rate due to recurrent myocardial infarction). This difference certainly appears significant. However, this paper presumably a preliminary report, is brief and sketchy. Final judgment must await publication of the detailed data.

The third report involved a follow up of 175 patients for a period of from 36 to 72 months (368). All patients were originally put on a low fat, low cholesterol diet. Of these 26 failed to respond with a reduction of cholesterolemia, and 39 others after a brief period failed to adhere to the diet. These two groups were used as the controls. In the 110 patients responding to the diet, the serum cholesterol level fell from a mean control value of 329 mg % to a mean treatment concentration of 270. In every one cholesterol fell more than 10 per cent, and the cholesterol/phospholipid ratio fell at least 20 per cent. In this treated group only 10 deaths occurred during the mean treatment period of 50 months (mortality rate 11 per cent); in the last 11 months of therapy only one patient of 98 died. In contrast the mortality rates in the other two groups were 31 per cent (8 of 26 patients) and 33 per cent (12 of 39 patients) during 51 and 53 months of observation respectively. The design of this study, particularly the manner of establishing the control group is open to criticism, yet the results are suggestive.

Obviously, the question of the value of such low fat diets is unsettled. Additional work along these lines is needed in view of certain possible limitations in the design of these studies judging from the published reports. However the practicing physician would do well to consider this type of diet in dealing with the prevention and retardation of atherosclerosis. This is discussed in more detail below.

3 *Unsaturated Oils*—As already indicated particular attention has focussed during the last year or two on the effects of unsaturated vegetable and marine oils on cholesterolemia and β lipoproteinemia. As yet no reports have been published on the long term prophylactic and/or therapeutic effects of these oils. However the practicing physician will be well advised to consider the use of unsaturated fats in lieu of saturated ones as part of his armamentarium in the handling of atherosclerosis. The substitution of unsaturated for saturated fats in diets kept moderate or low in total fats is warranted in our present state of knowledge. This can be accomplished by proper selection of foods (see below). It is not warranted today merely to add unsaturated fats as supplements—in the form of corn oil, safflower oil and their like. *The total fat and saturated fat content should be lowered* and overweight avoided. Specific dietary recommendations are given below.

4 *Sitosterols*—Another specific approach to therapy in recent years involves plant sterols. Extensive studies were undertaken on the effects of sitosterols in man. With few exceptions ingestion of large doses of plant sterols induced significant sustained reductions in serum choles-

weight remained stable. Serum total lipids, cholesterol and neutral fats declined markedly. Survival rates were significantly better in the treated group, compared with the controls, particularly at the 8 year interval (Fig 59). However, examination of this paper gives no clearcut indication that the two groups of "unselected patients" were in fact matched by age and sex for example. Hence the validity of these apparently impressive findings cannot be unequivocally accepted.

Effects of low-fat, low cholesterol diet on patients with Coronary Heart Disease




FIG 59 Morrison L M (366). The two groups were made up of unselected patients with proved myocardial infarction who survived their initial attack and whose condition was uncomplicated by other diseases such as hypertension, diabetes mellitus, nephritis, thyroid disease, xanthomatosis. (366). No data are given which permit assessment of the comparability of the two groups on an age-sex basis. Note the lower mortality rates at both 3 and 8 years in the group on low fat, low cholesterol dietary treatment.

The second report presented a 4 year follow up of 280 patients with previous myocardial infarction, of whom 155 presumably adhered to a low fat, low cholesterol regimen (50 grams fat, 200 mgm cholesterol daily) (367). The two groups, dieters and non dieters, were of similar age. They had comparable β lipoprotein levels at the onset of the study. The therapeutic diet effected a significant decline in these values, whereas no change occurred in the untreated group. With an average follow up period of about 4 years, the treated group had a significantly lower in

people free of clinically manifest atherosclerotic disease. It should be considered only in those who have had myocardial infarction or a cerebrovascular accident, especially when there is evidence of repeated thromboembolism. Close observation of all patients on long term anticoagulant therapy, with frequent determinations of prothrombin time, is of course absolutely essential to avoid potentially dangerous complications.

Four year mortality in cases of Myocardial Infarction treated with long-term anticoagulants




FIG 60. Keyes J W, Drake E H and Smith F J (757). Although the statistical methods for evaluation of survival time employed in this study seem predicated on comparability of control and treated groups, the published report presents no data on age, sex, previous medical history (other diseases) and so forth. Therefore comparability of the groups cannot be definitively evaluated.

7 Antihypertensives—Hypertension markedly increases the risk of atherosclerotic disease, particularly in the presence of a derangement in cholesterol lipid lipoprotein metabolism.

With recent progress in drug therapy for the hypertensive state, it is becoming possible to assess the effects of successful blood pressure reduction on development and progression of atherosclerotic disease. Thus far only limited—but encouraging—data have been presented (772). On this basis, antihypertensive drug therapy, or even surgical sympathectomy, is to be considered seriously as a means of helping retard atherosclerosis in any patient who has clinical manifestations of coronary or

terol (725-740) Recent studies indicate that marked lowering of cholesterol levels may be effected by combining plant sterols and dietary changes (362, 363, 403, 741) Preliminary data have been presented suggesting favorable long term therapeutic effects of sitosterols in patients with clinical atherosclerotic disease (742, 743) But definitive evaluation of this proposed therapeutic regimen must await more detailed reports

Results similar to those obtained with sitosterols have been reported in studies on man utilizing a brain extract (572, 573 575 576) Here again, no long term evaluations have appeared in the literature This method is therefore not yet suitable for the practicing physician

5 *Lipotropic Factors* —Negative results have been obtained in most studies in man (as well as experimental animals) with lipotropic factors—choline inositol lecithins, lipocaine and other pancreatic preparations methionine and other factors (1 12-14) With few exceptions no cholesterolemia reducing effects have been reported Further no data have been published on long term prophylactic or therapeutic effects The status of tocopherols (vitamin E) is similar

Recent reports suggest that significant results in lowering serum cholesterol may be achieved in man with large doses of nicotinic acid or pyridoxine with a dietary regimen incorporating linolenic acid pyridoxine and chelating agents and with several non biological chemicals (538 586-594, 741 744-748) However none of these has been studied in long term investigations and none of these are recommended for general clinical use

6 *Anticoagulants* —Research on the possible efficacy of long term anti coagulant therapy has advanced further (749-764) Suggestive positive results—reduction in recurrences and prolongation of life in patients with coronary heart disease—were reported in several different studies using chiefly dicumarol at dosage levels yielding sustained prolongation of prothrombin times (Fig 60) However a major cooperative British study has not yet revealed definitive results (765) Marked reduction in mortality—4 per cent cardiovascular renal mortality rate in the treated group vs 18 per cent in the control—was also reported in a series of cases of myocardial infarction treated intermittently with parenteral heparin (750) The doses utilized (200 mg twice weekly) did not produce sustained therapeutic anticoagulant levels On the other hand most studies with heparin for the treatment of angina pectoris have yielded negative results (766-771)

However detailed review of the papers on long term anticoagulant therapy raises questions as to the adequacy of controls in some of these investigations With this reservation it may be generally stated that the overall results are sufficiently suggestive particularly with dicumarol to justify its consideration for long term therapy in some cases pending the final report of the more adequately designed British study Anti coagulant therapy, at present, has no applicability prophylactically in

considerable data have been presented particularly from the Joslin Clinic indicating that effective control of diabetes mellitus results in significantly fewer vascular complications. It would seem valid that there may be a risk of aggravating athero sclerosis by attempts at very rigid control of the diabetic state, particularly with the use of too much insulin. The practicing physician would be well advised therefore, not to enforce too rigid diabetic control to avoid unnecessarily large doses of insulin with the inevitably excessive fluctuation of blood sugar level, and to use the apparently more physiologic oral insulins wherever possible.

9 *Thyroid Hormone*—Recent studies have re opened the question of the value of thyroid hormone in the prophylaxis and therapy of atherosclerotic disease (1, 8 12-14 645 688). This problem has a long history. It was recently reported that marked sustained reduction in human serum cholesterol and β lipoprotein concentrations can be effected by administration of large doses of desiccated thyroid (3 4, 5 or more grains daily) (Fig 61) (774, 775). These lowered levels were successfully maintained for many months with continuous hormone exhibition. Obviously necessary precautions are indicated in utilizing this medication in patients with manifest clinical coronary disease and resultant limited myocardial reserve. Again, the fact must be noted that no data are available on the long term prophylactic or therapeutic effects of thyroid hormone.

Possible utility of thyroid hormone as a specific definitive method of prophylaxis and therapy for atherosclerotic disease does not negate the possible value of thyroid ablation as a late palliative measure in selected cases of intractable angina pectoris (776). This applies also to the use of thiouracil and I^{131} . However the evidence available in our opinion points to a real hazard associated with all procedures which decrease thyroid function below normal—namely the potential of accelerating atherosclerosis. Proper diet control could lessen this hazard. A balance must be struck between the possible benefits of thyroidectomy, thiouracil and I^{131} on anginal symptoms—when they are debilitating—and the possible harm arising from acceleration of atherosclerosis. Obviously therefore the anginal syndrome must be sufficiently severe in the patient before these thyroid depressing procedures are contemplated.

10 *Estrogens*—The rationale for the use of estrogens in prophylaxis and therapy was summarized above. To date no data have been presented concerning the possible prophylactic utility of estrogens in preventing development of clinical atherosclerotic disease in postmenopausal women. However if uterine bleeding can be controlled there is no harm in their use—provided small doses are employed. It would appear unlikely that estrogens will find any use in the prophylaxis of atherosclerotic diseases in males unless future results should demonstrate an influence in small doses producing no side effects—or if an estrogen is found with proved antiatherogenic potency but no feminizing effect. No such estrogen is yet available.

cerebrovascular atherosclerosis. Best of all antihypertensive procedures and diet regulation should be combined. The use of antihypertensive drugs—particularly the more powerful depressors, e.g. ganglionic blocking agents in—benign hypertension, before such clinical manifestations have appeared, has not been clearly established. However their possible employment when the risk of clinically manifest atherosclerosis is great (see below) should be seriously considered.

Effects of long term administration of Desiccated Thyroid on serum cholesterol levels




FIG 61 Strisower B Gofman J W Galton E F Rubinger J H Pouteau J and Guzovich P (775) Note that the lower dosages of desiccated thyroid reduced serum cholesterol level initially but that it rose latterly. With a dosage of 325 mg /patient/day moderate sustained decrease of cholesterolemia was apparently achieved.

8 Diabetes Control The complex and controversial problem of control in diabetics and the development of vascular complications has been discussed previously (*cf* 1 12-14 773). Diabetes mellitus is a condition which indicates the use of diets restricted in total fat and cholesterol low in saturated fat adequate in essential fatty acids minerals vitamins proteins. With such a diet the marked tendency of diabetics to manifest hypercholesterolemic hyperlipemia can often be counteracted and the integrity of the vasculature maximally protected. It is also true that

PRE TREATMENT SERUM LIPOPROTEINS




FIG 63 Stamler J Pick R Katz L Kaplan B and Pick A (721) Note that the 4 groups had similar control serum lipoprotein levels

SERIAL EFFECTS OF ESTROGENS (100 MG/DAY) ON SERUM LIPIDS—SERIES PATIENTS




FIG 64 Stamler J Pick R Katz L Kaplan B and Pick A (721) Note that the effects of estrogens (100 mg/patient/day) on serum lipid levels tended to persist for as long as 2 years

Estrogen therapy in man was the area of activity which our group selected because of our work with estrogens in animals. In 1952, a long term evaluation was begun of estrogen therapy in middle aged males with previous myocardial infarction*. This study was recently described in comprehensive reports (645, 721, 777). Every effort was made to take necessary precautions particularly proper matching of placebo and treated groups, as outlined above. All patients are men under age 50 with proved coronary disease. The groups are in fact matched closely —

PRE TREATMENT SERUM LIPIDS





FIG 62 Stamler J, Pick R, Katz L, N. Kaplan B and Pick A (721). Note that the 4 groups in this long term therapeutic evaluation had similar control serum lipid levels.

by age, ethnic origin, occupation, disease history, height, weight, pre-treatment plasma cholesterol, lipid, lipoprotein levels (Figs 62, 63 and 64). Since June 1954 the standard maintenance dosage has been 10 mg of conjugated equine estrogens daily by mouth. It is our current practice to arrive at this dosage level gradually in a stepwise fashion over months beginning with 1.25 mg, then increasing to 2.5, 4.0 and then 10 mg.

The anticipated effects on serum lipids, lipoproteins have been consistently observed: i.e. an increase in α lipoprotein and phospholipids with a

* It was recently reported that estrogens are ineffective in the treatment of intermittent claudication and angina pectoris (778, 779). On the other hand, a preliminary report suggests favorable results with prolonged low-dosage estrogen therapy in idiopathic hyperlipemia and hypercholesterolemia (780).

Survival during First 3½ Years of Study (data as of March 1958)



	Survived %	Risk %			No. with Diabetes	Age	Serum Cholesterol / mm	Height	Weight	No. started on low dosage on 10 mg	Total No. test subjects	Total No. of Deaths	No. of non-CV deaths	No. CV deaths	No. of total deaths
		G	P	U											
PLACEBO % 58	58	30		7		4	96	90		0		4		0	
ESTROGEN % 61	44	24	10	10		29	24							2	5

FIG 66 Stamler J Pick R Katz L N Kaplan B and Pick A (721) The total number of patients under observation for as long as 3½ years were 40 and 61 in the placebo and estrogen treated groups respectively. Data on the comparability of the two groups are summarized in the table (cf text and ref 721). The patients were divided into good (G) poor (P) and unknown (U) risk based on the presence or absence of complications during the acute episode of myocardial infarction: all patients with more than one infarction prior to the onset of the study were classified as poor risk. There were no non-cardiovascular renal deaths in the placebo group; hence the upper graph presents only 3 curves of survival. There were 3 non-cardiovascular renal deaths in the estrogen treated group; the survival curve for the estrogen treated patients excluding these 3 patients (a total of 58 patients) is the upper curve in the upper graph. The number immediately to the right of the survival curve is the number of patients represented on the given curve. The lower graph plots the survival curve for valid deaths, i.e. cardiovascular renal deaths occurring after 2 months of medication had transpired. There were no non-cardiovascular renal deaths and no deaths during the first 2 months in the placebo group; hence the curve plots the data for the entire group of 40 patients. There were 3 non-cardiovascular renal deaths after and 1 cardiovascular renal death during the first 2 months of drug administration in the estrogen treated group; hence the survival curve for the estrogen treated patients in the lower graph excludes these 4 cases (leaving a total of 57 patients). Although these data indicate that estrogen therapy may be effecting a better prognosis for survival, the size of the groups followed for as long as 3½ years is too small to yield a statistically significant clearcut result. A longer period of follow-up is essential to evaluate whether estrogens are or are not efficacious in the therapy of coronary heart disease in middle aged males.

tendency for cholesterol to fall (Figs 64 to 66) (1, 4 5, 12-14, 645, 646, 704-719, 721) Hence, the ratios β/α lipoprotein and cholesterol/phospholipids are lowered These effects have been maintained for at least three, and as many as four years of treatment The α lipoprotein level—together with signs of gynecomastia—is therefore utilized as a reliable objective criterion of patient adherence to the estrogen regimen

SERIAL EFFECTS OF ESTROGENS (100 MG/DAY) ON SERUM LIPOPROTEINS—SERIES PATIENTS




FIG 65 Stamler J Pick R Katz L N Kaplan B and Pick A (721) Note that the estrogen induced increase in $-S : 1-10$ (α) lipoprotein levels tended to persist over as long as 2 years Similar data were obtained for an additional third year as of April 1957 (777)

As of March 1, 1958 276 patients had been brought under study, with the cooperation of several Veterans Administration hospitals throughout the Middle West An approximate 27 per cent attrition (drop out of patients) has occurred in the estrogen treated group compared to 19 per cent in the placebo group a difference attributable to the side effects of the hormone gynecomastia depression of libido and potency However these have certainly not been insurmountable problems On the dosage of 10 mg of the conjugated equine estrogens toxic effects (as distinguished from side effects)—e.g nausea lassitude malaise diarrhea—have been minimal With respect to effects on recurrence rates of myocardial infarction and on survival, the data to date indicate the following trends While the overall rate of recurrence of myocardial infarction is similar in the two groups, the survival rate is higher in the treated group under observation for over 3 years (Fig 66) These differences are not at the level of statistical significance A longer period of follow up is essential We are therefore, not yet prepared to advocate use of estrogens routinely

of these actuarial predictions of risk will be false positives and false negatives respectively. However, these "errors" are of minor importance, reckoned against the ability to identify high risk individuals—with the consequent possibility of successful prophylactic intervention.

Recent investigations are making it possible to estimate the scope of this problem of high risk individuals in specific quantitative terms. Based on the data of the Framingham Heart Study (Fig. 18), for example, it can be roughly estimated that a low risk middle aged male—normal in weight, blood pressure and serum cholesterol—has 1 chance in 25 of

PERCENT OF MALES AGE 45-62 WITH HYPERCHOLESTEROLEMIA OBESITY HYPERTENSION -- FRAMINGHAM STUDY




FIG. 6/ Dawber T R, Moore F E and Mann G V (209). Note the extent to which these defects individually and in combination are present in the middle aged male population of Framingham, Massachusetts.

developing clinical coronary heart disease during the 45 to 64 age period. In contrast a middle aged male with two or three abnormalities (obesity, hypercholesterolemia, hypertension) stands almost 1 chance in 2. These are very different risks.

About 1 of every 8 middle aged American males falls into this high risk category (Fig. 67) (209). When it is further recognized that 20 to 30 per cent of first attacks of myocardial infarction are fatal within the first few weeks, the serious nature of the high risk situation is even more starkly apparent. Is it not extremely valuable—for both patients and physicians—to be able to identify these high risk individuals in order to attempt to do something for them prophylactically?

In high risk subjects can risk be prophylactically reduced several fold by correcting defects? It is known that these defects are amenable to

E Overall Evaluation

It is evident that the research activity of the last decade has led to several specific approaches, particularly to definitive therapy of clinical coronary heart disease. However, for each of these approaches, unequivocal proof is not yet available concerning their efficacy. Only the work in the next decade can complete this complex task of evaluation. However, some suggestions—as distinct from scientifically clear proof—have been *derived and serve as the basis for certain tentative recommendations* we have made above for the physician who is in practice. Time alone will tell how sound they are.

F An Interim Approach to Prophylaxis for High Risk Persons, and to Therapy

Throughout the foregoing discussion the key role of diet has been emphasized. Obviously, routine therapy other than this is not yielding satisfactory results (781-784). There is room for improvement. This conclusion in no way gainsays the importance of such established constituents of the therapeutic armamentarium as indicated medications and specific hygienic admonitions—e.g. reasonable schedules of work, relaxation, recreation and rest, avoidance of tension, frustration, stress, over exertion, fatigue, undue competitiveness, prescription of appropriate physical exercise to maintain physical fitness and the like (see below). These recommendations continue to be of fundamental importance even as the medical profession goes beyond them to employ the more definitive therapeutic approaches. The cornerstone of modern therapy for patients with coronary disease today seems to be an increasingly scientific dietary prescription aimed at correcting the nutritional metabolic derangements intensifying atherogenesis. This is obviously of value for the person with clinical manifestations of coronary and cerebrovascular atherosclerosis. It is valuable also for persons with a genetic handicap as regards atherosclerosis.

For persons not yet victimized by clinical coronary disease it is also widely agreed among most investigators (6, 195, 372, 373) that one sizable sector of the American population particularly needs prophylactic advice from the medical profession, i.e. the middle aged male group with an inordinately high risk of developing clinical coronary heart disease. This is the group with two or more derangements—obesity, hypercholesterolemia, hypertension, renal damage, diabetes, hypothyroidism, heavy smoking and/or a poor family history.

While it is perfectly correct that no one can definitively predict whether a given person will or will not develop clinical coronary heart disease in the next year or two, long term prognostic predictions of the actuarial type—predictions of risk—can be made. Inevitably a small percentage

through utilization of vegetable oils, *e g*, corn oil. As one important measure towards this end, the ordinary bacon eggs-buttered toast creamed coffee breakfasts—with their huge intake of saturated fats, calories and cholesterol—should become the exception and not the rule. They should be largely superseded by breakfasts made up of fruits, whole grain cereals, skim milk (enriched with nonfat dry milk solids), waffles or wheat cakes (made with enriched white flour and prepared with vegetable oil and skim milk), and spreads of honey, jam or marmalade used in moderation on whole grain or enriched breads.

5 At other meals, the use of meats in moderation—preferably lean—prepared by broiling, roasting, boiling, steaming, roasting with trimming of visible fat and avoidance of fat laden gravies and sauces. Meat tenderizers may be used for flavoring—as can relishes and spices. The utilization of poultry and sea foods in quantity prepared according to the foregoing recommendations. The ample intake of vegetables and fruits. The minimal use of highly processed refined foodstuffs having a low ratio of nutrients to calories. These measures, in addition to the other objectives they achieve, will insure the ingestion of a high protein diet.

There is no doubt that this diet should be prescribed for every person who has had one or more clinical episodes of atherosclerotic disease such as angina pectoris, myocardial infarction, certain forms of cerebral vascular diseases (those due to atherosclerosis) and atherosclerotic peripheral vascular disease. It would seem logical to us that the above diet should also be followed by the following groups of persons—all of whom are subjects with a high risk of developing clinical atherosclerosis:

- 1 Those who are obese on a typical American diet
- 2 Those who have a hypercholesterolemia over 260 or 275 mg per cent
- 3 Those who are hypertensive, have diabetes mellitus, or have renal damage (provided there is no specific contraindication in any particular patient)
- 4 Those who have a poor genetic history, *i e*, a familial predilection for clinical atherosclerosis in middle age

Such a program, we feel convinced, will reduce clinical atherosclerosis in this country to a significant extent within 5 years. However, this remains to be proven.

This dietary regimen at first requires constant attention and adjustment on the part of the physician. During this time the physician can combine it with simple and sound advice on a way of life for his patient who is approaching the second half of his life. Emphasis should be on moderation—in his vocation and avocation. There should be a reduction of excessive competition, acceptance of one's limitation and pride in what one has accomplished—without too much smugness. The physician should preach a habit of not so much ambition as to make the person constantly unhappy, nor too little to make him bored—and boring. The

correction in whole or in part by relatively simple diet therapy. It is not yet known definitively whether coronary heart disease can be significantly lowered thereby, although the findings of the life insurance companies on the positive results of correcting obesity are highly suggestive in this regard (Fig 58).

There is widespread agreement among investigators on the individualized prescription of reasonable, moderate, safe, palatable dietary correction in high risk patients, particularly those with obesity, hypercholesterolemia, hypertension, renal damage, diabetes, poor family history, and other factors (6, 195, 372, 373, 720).

G Specific Recommendations

Recommendations for prophylaxis should be virtually without danger of any type—biological, sociological or psychological. For this reason—and because the accumulated evidence strongly suggests that the best results are likely to be achieved with *dietary* prophylaxis (coupled with a good overall hygienic regimen including regular physical exercise)—our general recommendations focus on this approach.

Every effort should be made to advise prophylactic dietary corrections that are simple, moderate, familiar and feasible. The average high risk person should not be enjoined to abandon the pleasures of eating. Rather, he should be encouraged to graduate to a modified appetizing cuisine including a wide enough variety of palatable foods to allow even the gourmet ample play for his proclivities. Stringent dietary restrictions (and related adjuvant measures) should be undertaken only when absolutely necessary, i.e. when moderate measures fail to elicit a reduction in weight, cholesterolemia, blood pressure.

The general objective then is to correct the imbalances in the American diet, cutting down on excesses and overcoming inadequacies. This approach calls for the following simple, safe, reasonable, nutritional recommendations:

1. The curtailment of the culinary custom of adding large amounts of fats, particularly saturated fats, to foods; the avoidance of frying and deep fat frying; of preparing salads with cheese or sour cream dressings, and of putting butter, margarine or cream on vegetables.

2. For dessert, adherence to the French tradition of serving fruit, instead of the American routine of pies, cakes, ice cream and other similar desserts.

3. The correction of obesity, if present, by reducing total caloric intake, total fat intake and ingestion of 'empty calories' from refined carbohydrates, processed fats, spirits and so forth.

4. The reduction of fat intake from the present 40 to 60 per cent of total calories down to 20 to 25 per cent. Along with this, the accomplishment of an increase in the ratio of unsaturated fats to saturated fats—

Bibliography

General Reviews on Atherosclerosis

- 1 KATZ L N and STAMLER J *Experimental Atherosclerosis* Springfield Ill Charles C Thomas 1953
- 2 Measuring the Risk of Coronary Heart Disease in Adult Population Groups—A Symposium *Am J Pub Health* 47 No 4 Part 2 April 1957
- 3 PAGE I H *Atherosclerosis* *Circulation* 10 1 1954
- 4 Symposium on Atherosclerosis National Academy of Sciences-National Research Council Washington D C 1954 NAS NRC Publication No 338 1954
- 5 KEYS A Editor *A Symposium on Arteriosclerosis* Minneapolis Minnesota Heart Association and University of Minnesota 1956
- 6 Symposium Fats in Human Nutrition *JAMA* 164 1890 1957
- 7 PAGE I H Editor *Chemistry of Lipides as Related to Atherosclerosis* Springfield Ill Charles C Thomas 1958
- 8 PINCUS G Editor *Hormones and Atherosclerosis* Proceedings of a Conference New York Academic Press in press
- 9 Symposium on Atherosclerosis *Am J Med* 23 120 269 463 769 898 928 1957
- 10 COWDRY E V *Arteriosclerosis* New York The Macmillan Co 1943
- 11 HUEPER W C *Arteriosclerosis* *AMA Arch Path* 38 167 245 350 1944 and 39 51 117 187 1945

Nutrition and Atherosclerosis

A *The Nutritional Metabolic Cholesterol Lipid Lipoprotein Theory of Atherosclerosis*

- 12 STAMLER J KATZ L N and PICK R Nutritional Endocrinological Interrelationships in the Pathogenesis of Hypertension and Atherosclerosis over the Life Span Fourth Annual Report on Stress Selye and Heuser Editors Montreal Acta p 245 1954
- 13 STAMLER J KATZ L N PICK R and ROBBARD S Dietary and Hormonal Factors in Experimental Atherogenesis and Blood Pressure Regulation *Recent Progress in Hormone Research* 11 401 1955 New York Academic Press Inc 1955
- 14 KATZ L N STAMLER J and PICK R The Role of the Hormones in Atherosclerosis *Symposium on Atherosclerosis* National Academy of Sciences—National Research Council Washington D C 1954 NAS NRC Publication 338 236 1954
- 15 KATZ L N PICK R and STAMLER J *Atherosclerosis* Modern Concepts of Card ovascular Disease 23 239 1954
- 16 STAMLER J Basic Research on Atherosclerosis *J Am Pharm Mfgs Assoc* p 34 1955
- 17 KATZ L N STAMLER J and PICK R Research Approach to Atherosclerosis (An Editorial) *JAMA* 161 536 1956
- 18 ——— Atherosclerosis and Coronary Heart Disease *J Lancet* 76 355 1956
- 19 STAMLER J Current Epidemiological Clinical and Laboratory Research Findings on the Etiology of Atherosclerosis *Nebraska State Med J* 41 75 1956

patient should run his job, not have the job run him. He should have peace of mind and plenty of physical exercise to overcome our civilization's hectic pace and sedentary trend. Frustrations should be kept at a minimum, competition—while useful as a driving force—should not become too much of a compulsion. In short, the high risk individual must be reminded that it is not how high he (or she) climbs that is harmful, but rather how fast and over how many obstacles. Physicians can help high risk patients with general advice of this sort, if they are willing to spend enough time with their patients over their troublesome day by day problems. The ability of a patient to unburden himself to a sympathetic and knowledgeable physician is obviously beneficial psychologically—and doubtlessly medically. This role, we believe, should be assumed by more physicians.

As more and more knowledge comes to the fore, generalities in medical advice will become more specific. There is great need, in our opinion, to study more intensively the role of the psychological and emotional make up of the individual, and of physical and psychical stresses in the atherosclerotic process.

Finally, our attention has been on the etiology, prophylaxis and treatment of the basic process of atherosclerosis. There is adequate coverage of the handling of sequelæ and clinical manifestations in other books, monographs and reports. However, we must stress that if atherosclerosis is absent, its complications and sequelæ cannot occur!

- 51 THOMASON M J TOBAR R OLMEDO R SANCHEZ M and VARGAS C Dietary Studies in Ecuador *Am J Clin Nutr* 5 295 1957
- 52 WHITE H S COLLAZOS C WHITE P L HUENEMANN R L BENITES R CASTELLANOS A BRAVO Y MOSCOSO I and DIESELDORFF A Dietary Surveys in Peru II Yurimaguas A Jungle Town on the Huallaga River *J Am Dietet Assoc* 30 856 1954
- 53 NICOL B M Tribal Nutrition and Health in Nigeria A Comparative Clinical Study of Primitive and Urban Nutrition *J Clin Nutr* 1 364 1953
- 54 WALKER A R P and ARVIDSSON U B Studies on Human Bone from South African Bantu Subjects I Chemical Composition of Pibs from Subjects Habituated to a Diet Low in Calcium *Metabolism* 3 385 1954
- 55 BROCK J F and AUTRET M *Aashstorkor in Africa* Geneva World Health Organization 1952
- 56 GILLMAN J and GILLMAN T *Perspectives in Human Malnutrition A Contribution to the Biology of Disease from a Clinical and Pathological Study of Chronic Malnutrition and Pellagra in the African* New York Grune and Stratton 1951
- 57 GRANT F W Nutrition and Health of Gold Coast Children I Food in Four Communities *J Am Dietet Assoc* 31 685 1955
- 58 FORBES W H Food and Population Illustrated by Observations in India *Fed Proc* 11 667 1952
- 59 KRIDAKARA O BOOZAYAANGOL R YUKTANANDA I and VOLKER J F Dental Survey of Selected Thai Children *Am J Clin Nutr* 4 280 1956
- 60 MAYER J ROY P and MITRA K P Relation Between Caloric Intake Body Weight and Physical Work Studies in an Industrial Male Population in West Bengal *Am J Clin Nutr* 4 169 1956
- 61 ARIMOTO K Nutrition in Japan *Nutr Revs* 10 321 1957
- 62 POLLACK H Special Editor Symposium on Nutrition—Studies on Nutrition in the Far East *Metabolism* 5 203 1956
- 63 YUDKIN J Food Population and the Future *Triangle* 11 1 1955
- 64 JOLLIFFE N Recent Advances in Nutrition of Public Health Significance *Metabolism* 4 191 1955
- 65 LEGROS CLARA F *Feeding the Human Family—Science Plans for the World Larder* London Sigma 1947
- 66 SAND R *The Advance to Social Medicine* London Staples Printers Ltd 1952
- 67 BROZEK J BUZINA R and MIXIC F Population Studies on Serum Cholesterol and Dietary Fat in Yugoslavia *Am J Clin Nutr* 5 279 1957

D The Epidemiologic Findings

- 68 Annual Epidemiological and Vital Statistics 1950 Part I Vital Statistics and Causes of Death Geneva World Health Organization 1953
- 69 ROSENTHAL S R Atherosclerosis Chemical Experimental and Morphologic *A M A Arch Path* 18 473 660 827 1934
- 70 RAAB W Diet Hormones and Arteriosclerosis Data and References from the Older European and Oriental Literature Assembled and published by the Author
- 71 KEYS A The Diet and the Development of Coronary Heart Disease *J Chronic Dis* 4 364 1956
- 72 KEYS A and WHITE P D Editors *World Trends In Cardiology I Cardiovascular Epidemiology* Selected Papers from Second World Congress and Twenty Seventh Annual Scientific Sessions of the American Heart Assoc New York Hoeber Harper 1956
- 73 BECKER B J P Cardio Vascular Disease in the Bantu and Coloured Races of South Africa V Hypertensive Heart Disease *S Afr J Med Sci* 11 107 1946
- 74 ——— Cardio Vascular Disease in the Bantu and Coloured Races of South Africa IV Atheromatosis *S Afr J Med Sci* 11 97 1946
- 75 ——— Cardio Vascular Disease in the Bantu and Coloured Races of South Africa I Incidence Pathology and General Features *S Afr J Med Sci* 11 1 1946

- 20 KATZ L N STAMLER J and PICK R Nutrition and Atherosclerosis Fed Proc 15 885 1956
- 21 STAMLER J Evaluation of Current Approaches to the Therapy of Atherosclerotic Disease In Cerebral Vascular Diseases—Transactions of the 2nd Conference p 98 New York Grune and Stratton 1958
- 22 ——— Diet and Atherosclerotic Disease Recent Pathologic Clinical Animal Experimental and Epidemiologic Findings J Am Dietet Assoc 34 701 814 929 1053 1060 1958

B The Epidemiologic Method

- 23 STAMLER J Epidemiology as an Investigative Method for the Study of Human Atherosclerosis J Nat Med Assoc 50 161 1958
- 24 MORRIS J N Uses of Epidemiology Baltimore The Williams and Wilkins Co 1957
- 25 ——— Uses of Epidemiology Brit Med J 2 395 1955
- 26 GARLAND J The Epidemiologic Method New Eng J Med 254 1044 1956
- 27 GREGG A The Future Health Officers Responsibility Past Present and Future Am J Pub Health 46 1385 1956
- 28 CLARK E G Modern Concepts of Epidemiology J Chronic Dis 2 593 1955
- 29 GILLIAM A G Epidemiology in Noncommunicable Disease Pub Health Rep 69 907 1954
- 30 CRICKSHANK R Epidemiological Methods in Preventive Medicine Proc Roy Soc Med 45 621 1952
- 31 LILLENFELD A M Epidemiological Methods and Inferences in Studies of Non Infectious Disease Pub Health Rep 72 51 1957
- 32 SINGER H E A History of Medicine Vol I Primitive and Archaic Medicine New York Oxford University Press 1951

C Human Nutrition—Historical and Ecological Background

- 33 JENSEN L B Man's Foods Champaign Ill The Garrard Press 1953
- 34 PRICE W A Nutrition and Physical Degeneration—a Comparison of Primitive and Modern Diets and Their Effects Am Acad of Applied Nutr Los Angeles 1948
- 35 CHILDE V G What Happened in History? Harmondsworth Eng Penguin Books Ltd 1942
- 36 ——— Man Makes Himself New York New American Library 1951
- 37 HOWELLS W Back of History Garden City N Y Doubleday and Company Inc 1954
- 38 The Bible Old Testament Genesis 45 17–18
- 39 STEFANSSON V The Fat of the Land New York The Macmillan Co 1956
- 40 Demographic Yearbook Statistical Office of the United Nations Department of Economic Affairs New York 1953
- 41 Statistical Yearbook Statistical Office of the United Nations Department of Economic Affairs New York 1953
- 42 The State of Food and Agriculture 1954 Review and Outlook Food and Agriculture Organization of the United Nations Rome Italy 1954
- 43 HANLON J J The Design of Public Health Programs for Underdeveloped Countries Pub Health Rep 69 1078 1954
- 44 BRODY S Facts Fables and Fallacies on Feeding the World Population Fed Proc 11 631 1952
- 45 SLURE B The Critical World Food Situation Am J Clin Nutr 4 211 1956
- 46 Anonymous Disease and Nutritional Barriers to Health Pub Health Rep 68 1127 1954
- 47 WILLIAMS R R The World Beriberi Problem Today J Clin Nutr 1 513 1953
- 48 JENNEY E R Public Health in Indonesia Pub Health Rep 68 409 1953
- 49 Anonymous Demographic Characteristics of Latin America Pub Health Rep 68 1123 1953
- 50 Anonymous Nutrition in the Americas Pub Health Rep 69 277 1954

- 104 HAMMOND E G and LUNDBERG W O The Effect of Low Fat Diet and Atherosclerosis on the Polyunsaturated Fatty Acids of Human Plasma Arch Biochem and Biophys 57 517 1955
- 105 KEYS A The Age Trend of Serum Concentrations of Cholesterol and of S_f 10-20 (G) Substances in Adults J Geront 7 201 1952
- 106 KEYS A FIDANZA F SCARDI V and BERGAMI G The Trend of Serum Cholesterol Levels with Age Lancet 2 209 1952
- 107 KEYS A Diet and the Epidemiology of Coronary Heart Disease J A M A 164 1912 1957
- 108 SCRIMSHAW N S TRULSON M TEJADA C HEGSTED D M and STARE F J Serum Lipoprotein and Cholesterol Concentrations Comparison of Rural Costa Rican Guatemalan and U S Populations Circulation 15 805 1957
- 109 MANN G V MILNOZ J A and SCRIMSHAW N S The Serum Lipoprotein and Cholesterol Concentrations of Central and North Americans with Different Dietary Habits Am J Med 19 25 1955
- 110 GUBNER R and UNGERLEIDER H Arteriosclerosis Am J Med 6 60 1949
- 111 LEWIS L A OLIMSTED F PAGE I H LAWRY E Y MANN G V STARE F J HANIG M LAUFFER M A GORDON T and MOORE F E Serum Lipid Levels in Normal Persons—Findings of a Cooperative Study of Lipoproteins and Atherosclerosis Circulation 16 227 1957
- 112 LAWRY E Y MANN G V PETERSON A WYSOCKI A P O'CONNELL R and STARE F J Cholesterol and Beta Lipoproteins in the Serum of Americans Well Persons and Those with Coronary Heart Disease Am J Med 22 605 1957
- 113 KEYS A Mode of Life and the Prevalence of Coronary Heart Disease A Symposium on Arteriosclerosis p 28 Minneapolis Minnesota Heart Association and University of Minnesota 1956
- 114 NATH H P GUPTA K K and IYER P V K Serum Cholesterol in Indian Army Personnel Ind J Med Res 45 217 1957 Cited by Nutr Revs 15 375 1957
- 115 GORE I and TEJADA C The Quantitative Appraisal of Atherosclerosis Am J Path 33 875 1957
- 116 KIMURA N Analysis of 10 000 Postmortem Examinations in Japan World Trends In Cardiology I Cardiovascular Epidemiology p 22 New York Hoeber Harper 1956
- 117 KUSAKAWA A Statistical Findings on the Incidence of Coronary Heart Disease in Japan World Trends In Cardiology I Cardiovascular Epidemiology p 159 New York Hoeber Harper 1956
- 118 KEYS A KIMURA N KUSAKAWA A and YOSHITOMI M Serum Cholesterol in Japanese Coal Miners Am J Clin Nutr 5 245 1957
- 119 KEYS A FIDANZA F SCARDI V BERGAMI G KEYS M H and DI LORENZO F Studies on Serum Cholesterol and Other Characteristics of Clinically Healthy Men in Naples A M A Arch Int Med 93 328 1954
- 120 KEYS A FIDANZA F and KEYS M H Further Studies on Serum Cholesterol of Clinically Healthy Men in Italy Voeding 16 497 1955
- 121 KEYS A Field Studies in Italy 1954 World Trends in Cardiology I Cardiovascular Epidemiology p 50 New York Hoeber Harper 1956
- 122 ——— Diet and Coronary Heart Disease World Trends in Cardiology I Cardiovascular Epidemiology p 135 New York Hoeber Harper 1956
- 123 WHITE P D Clinical Studies in Italy 1954 World Trends in Cardiology I Cardiovascular Epidemiology p 67 New York Hoeber Harper 1956
- 124 KEYS A VIVANCO F RODRIGUEZ MINON J L KEYS M H and CASTRO-MENDOZA H Studies on the Diet Body Fatness and Serum Cholesterol in Madrid Spain Metabolism 3 195 1954
- 125 STAMLER J KJELSBERG M and HALL Y Epidemiologic Studies on Coronary Disease in the Chicago and Illinois Populations—Relationship of Sex Race Nationality Occupation Income Urban vs Rural Residence to Mortality Rates J Lab Clin Med 50 956 1957
- 126 STAMLER J KJELSBERG M HALL Y and SCOTCH N Epidemiological Studies on Cardiovascular Renal Diseases III Analysis of Mortality by Age Sex Nationality In preparation

- 76 DAVIES I G *Atherosclerosis at Autopsy in East Africa* Josiah Macy Liver Injury Symposium p 151 New York Josiah Macy Found 1950
- 77 MANN G V Coronary Artery Disease in Uganda Brit Med J 1 690 1956
- 78 TROWELL H C Coronary Artery Disease in Uganda Brit Med J 1 46 1956
- 79 VOGELPOLL I and SCHIRRL V Myocardial Infarction Its Racial Incidence in Cape Town Lancet 2 1108 1955
- 80 PALLISTER R A Cardiac Infarction in Malaya Brit Med J 1 27 1957
- 81 LARSEN N P The Animal Fat Diet and Atherosclerosis Hawaii Med J 14 485 1955
- 82 JALILI M A Heart Disease in Iraq In *World Trends In Cardiology I Cardiovascular Epidemiology* New York Hoeber Harper p 73 1956
- 83 ALIMURUNG M M HERRERA F Jr GUYTINGCO A and CRUZ P M Heart Disease in the Philippines Am Heart J 50 293 1955
- 84 ZUIDEMA P J Angina Pectoris and Coronary Thrombosis in Djakarta 1947-1951 Docum Med Geog et Trop 4 3/8 1952
- 85 VAKIL R J Heart Disease in the Tropics A Statistical Study of 1 860 Cases in India from a total Medical Population of 30 104 Cases The Indian Practitioner 7 1 1954
- 86 ——— Heart Disease in India Am Heart J 48 439 1954
- 87 VAN UNNIK J and STRAUB M The Frequency of Thrombosis and Pulmonary Embolism in East and West Docum Med Geog et Trop 5 261 1953
- 88 FERNANDEZ MENDIA J A Frecuencia de Cardiopatias en Guatemala Rev Colegio Guatemala 4 344 1953
- 89 TORRE J M CARELLO J O and VILDOSOLA D M Incidencia de las Cardiopatias en San Luis Potosi Anales de la Sociedad Potosina de Estudios Medicos 2 9 1953
- 90 WHITE P D Prevalence of Heart Disease and Its Etiologic Types Chapter 12 p 266 in *Heart Disease* 4th edition New York The Macmillan Co 1951
- 91 KEYS A Coronary Heart Disease and the Mode of Life Modern Concepts of Cardiovascular Dis 25 317 1956
- 92 WALKER A R P and ARVIDSSON U B Fat Intake Serum Cholesterol Concentration and Atherosclerosis in the South African Bantu Part I Low Fat Intake and the Age Trend of Serum Cholesterol Concentration in the South African Bantu J Clin Invest 33 1358 1954
- 93 HIGGINSON J and PEPLER W J Fat Intake Serum Cholesterol Concentration and Atherosclerosis in the South African Bantu Part II Atherosclerosis and Coronary Artery Disease J Clin Invest 33 1366 1954
- 94 HIGGINSON J Atherosclerosis Diet and Serum Cholesterol in the South African Bantu *World Trends In Cardiology I Cardiovascular Epidemiology* p 34 New York Hoeber Harper 1956
- 95 BRONTE STEWART B KEYS A and BROCK J F Serum Cholesterol the Diet and the Relationship to the Incidence of Coronary Heart Disease Lancet 2 1103 1955
- 96 BROCK J F and BRONTE STEWART B Arteriosclerosis in African Populations *A Symposium on Arteriosclerosis* p 102 Minneapolis Minnesota Heart Association and University of Minnesota 1956
- 97 SCRIMSHAW N S Measuring the Risk of Coronary Heart Disease—Implications to the Nutritionist Am J Pub Health 47 Part 2 60 1957
- 98 GOPALEN C and RAMANATHAN K S (Letter to the editor) Lancet 2 1212 1956
- 99 LARSEN N P Diet and Atherosclerosis—A Field Study AMA Arch Int Med 100 436 1957
- 100 TEJADA G and GORE I Comparison of Atherosclerosis in Guatemala City and New Orleans Am J Path 33 887 1957
- 101 KEYS A Epidemiologic Aspects of Coronary Artery Disease J Chronic Dis 6 552 1957
- 102 STARE F J Dietary Aspects Fed Proc 15 900 1956
- 103 MANN G V NICOL B M and STARE F J The Beta Lipoprotein and Cholesterol Concentrations in Sera of Nigerians Brit Med J 2 1008 1955

- 104 HAMMOND E G and LUNDBERG W O The Effect of Low Fat Diet and Atherosclerosis on the Polyunsaturated Fatty Acids of Human Plasma Arch Biochem and Biophys 57 517 1955
- 105 KEYS A The Age Trend of Serum Concentrations of Cholesterol and of S₁₀₋₂₀ (G) Substances in Adults J Geront 7 201 1952
- 106 KEYS A FIDANZA F SCARDI V and BERGAMI G The Trend of Serum Cholesterol Levels with Age Lancet 2 209 1952
- 107 KEYS A Diet and the Epidemiology of Coronary Heart Disease J A M A 164 1912 1957
- 108 SCRIMSHAW N S TRULSON M TEJADA C HEGSTED D M and STARE F J Serum Lipoprotein and Cholesterol Concentrations Comparison of Rural Costa Rican Guatemalan and U S Populations Circulation 15 805 1957
- 109 MANN G V MUÑOZ J A and SCRIMSHAW N S The Serum Lipoprotein and Cholesterol Concentrations of Central and North Americans with Different Dietary Habits Am J Med 19 25 1955
- 110 GUBNER R and UNGERLEIDER H Arteriosclerosis Am J Med 6 60 1949
- 111 LEWIS L A OLINSTEAD F PAGE I H LAWRY E Y MANN G V STARE F J HANIG M LAUFFER M A GORDON T and MOORE F E Serum Lipid Levels in Normal Persons—Findings of a Cooperative Study of Lipoproteins and Atherosclerosis Circulation 16 227 1957
- 112 LAWRY F Y MANN G V PETERSON A WYSOCKI A P O'CONNELL R and STARE F J Cholesterol and Beta Lipoproteins in the Serum of Americans Well Persons and Those with Coronary Heart Disease Am J Med 22 605 1957
- 113 KEYS A Mode of Life and the Prevalence of Coronary Heart Disease A Symposium on Arteriosclerosis p 28 Minneapolis Minnesota Heart Association and University of Minnesota 1956
- 114 NATH H P GUPTA K K and IYER P V K Serum Cholesterol in Indian Army Personnel Ind J Med Res 45 217 1957 Cited by Nutr Revs 15 325 1957
- 115 GORE I and TEJADA C The Quantitative Appraisal of Atherosclerosis Am J Path 33 875 1957
- 116 KIMURA N Analysis of 10 000 Postmortem Examinations in Japan World Trends In Cardiology I Cardiovascular Epidemiology p 22 New York Hoeber Harper 1956
- 117 KUSABAWA A Statistical Findings on the Incidence of Coronary Heart Disease in Japan World Trends In Cardiology I Cardiovascular Epidemiology p 150 New York Hoeber Harper 1956
- 118 KEYS A KIMURA N KUSABAWA A and YOSHITOMI M Serum Cholesterol in Japanese Coal Miners Am J Clin Nutr 5 245 1957
- 119 KEYS A FIDANZA F SCARDI V BERGAMI G KEYS M H and DI LORENZO F Studies on Serum Cholesterol and Other Characteristics of Clinically Healthy Men in Naples A M A Arch Int Med 93 328 1954
- 120 KEYS A FIDANZA F and KEYS M H Further Studies on Serum Cholesterol of Clinically Healthy Men in Italy Voeding 16 497 1955
- 121 KEYS A Field Studies in Italy 1954 World Trends in Cardiology I Cardiovascular Epidemiology p 50 New York Hoeber Harper 1956
- 122 ——— Diet and Coronary Heart Disease World Trends in Cardiology I Cardiovascular Epidemiology p 135 New York Hoeber Harper 1956
- 123 WHITE P D Clinical Studies in Italy 1954 World Trends in Cardiology I Cardiovascular Epidemiology p 62 New York Hoeber Harper 1956
- 124 KEYS A VIVANCO F RODRIGUEZ MINON J L KEYS M H and CASTROMENDOZA H Studies on the Diet Body Fatness and Serum Cholesterol in Madrid Spain Metabolism 3 195 1954
- 125 STAMLER J KJELSBERG M and HALL Y Epidemiologic Studies on Coronary Disease in the Chicago and Illinois Populations—Relationship of Sex Race Nationality Occupation Income Urban vs Rural Residence to Mortality Rates J Lab Clin Med 50 956 1957
- 126 STAMLER J KJELSBERG M HALL Y and SCOTCH N Epidemiological Studies on Cardiovascular Renal Diseases III Analysis of Mortality by Age Sex Nationality In preparation

- 127 TOOR M AGMON J and ALLALOUF D Changes of Serum Total Lipids Total Cholesterol and Lipid Phosphorus in Jewish Yemenite Immigrants after 20 Years in Israel *Bull Res Council of Israel* 4 202 1954
- 128 DREYFUSS F TOOR M AGMON J and ZLOTNICK A Observations on Myocardial Infarction in Israel *Cardiologia* 30 387 1957
- 129 Anonymous Serum Lipids and Atherosclerosis (foreign letter from Israel reporting on a symposium paper by Toor M) *JAMA* 166 817 1953
- 130 BRUNNER D IOEHL K FISCHER M and SCHICK G Cholesterol Phospholipids Total Lipids and the Cholesterol Phospholipid Ratio in Healthy Males of Various Origin in Israel Harefuah (Tel Aviv) 48 1 1955
- 131 BRUNNER D and LOEHL K Serum Cholesterol Electrophoretic Lipid Pattern Diet and Coronary Artery Disease *Ann Int Med* 49 732 1958
- 132 DREYFUSS F The Incidence of Myocardial Infarction in Various Communities in Israel *Am Heart J* 45 749 1953
- 133 KEYS A KIMURA N KUSUKAWA A BRONTE STEWART B LARSEN N P and KEYS M H Diet and Serum Cholesterol of Japanese Men in Japan Hawaii and California *Fed Proc* 16 204 1957
- 134 EPSTEIN F H and BOAS E P Prevalence of Manifest Atherosclerosis among Randomly Chosen Italian and Jewish Garment Workers *J Geront* 10 331 1955
- 135 EPSTEIN F H Relations between Diet and Atherosclerosis among a Working Population of Different Ethnic Origins *Am J Clin Nutr* 4 10 1956
- 136 EPSTEIN F H BOAS E P and SIMPSON R The Epidemiology of Atherosclerosis among a Random Sample of Clothing Workers of Different Ethnic Origins in New York City I Prevalence of Atherosclerosis and Some Associated Characteristics *J Chronic Dis* 5 300 1957
- 137 EPSTEIN F H SIMPSON R and BOAS E P The Epidemiology of Atherosclerosis among a Random Sample of Clothing Workers of Different Ethnic Origins in New York City II Associations between Manifest Atherosclerosis Serum Lipid Levels Blood Pressure Overweight and some other Variables *J Chronic Dis* 5 329 1957
- 138 TOOR M KATCHALSKY A AGMON J and ALLALOUF D Serum Lipids and Atherosclerosis among Yemenite Immigrants in Israel *Lancet* 1 1270 1957
- 139 YERUSHALMY J and HILLEBOE H E Fat in the Diet and Mortality from Heart Disease—a Methodological Note *New York State J Med* 57 2343 1957
- 140 HILLEBOE H E Some Epidemiologic Aspects of Coronary Artery Disease *J Chronic Dis* 6 210 1957
- 141 Anonymous National Food Supplies and Vital Statistics Dury Council Digests 28 1 1956
- 142 YUDKIN J Diet and Coronary Thrombosis *Lancet* 2 155 1957
- 143 MANN G V The Epidemiology of Coronary Heart Disease *Am J Med* 23 463 1957
- 144 RYLE J A and RUSSELL W T The Natural History of Coronary Disease A Clinical and Epidemiological Study *Brit Heart J* 11 370 1949
- 145 MORRIS J N Recent History of Coronary Disease *Lancet* 1 1 60 1951
- 146 MORRIS J N and DALE R A Epidemiology of Coronary Atherosclerosis *Proc Roy Soc Med* 48 667 1955
- 147 MORRIS J N HEADY J A and BARLEY R G Coronary Heart Disease in Medical Practitioners *Brit Med J* 1 503 1952
- 148 MORRIS J N HEADY J H RAFFLE P A B ROBERTS C G and PARKS J W Coronary Heart Disease and Physical Activity of Work *Lancet* 2 1053 1111 1953
- 149 MORRIS J N HEADY J A and RAFFLE P A B Physique of London Busmen Epidemiology of Uniforms *Lancet* 2 569 1956
- 150 LOGAN W P D Mortality from Coronary and Myocardial Disease in Different Social Classes *Lancet* 1 758 1952
- 151 Social Class Variations in Mortality *Brit J Prev Soc Med* 8 128 1954
- 15 Social Class Variations in Mortality *Pub Health Rep* 69 1217 1954
- 153 MARTIN W J The Distribution in England and Wales of Mortality from Coronary Disease *Brit Med J* 1 1523 1950

- 154 THOMAS A J COTES J E and HIGGINS I T T Prevalence of Coronary Heart Disease in Elderly Coal Workers *Lancet* 270 414 1956
- 155 OLIVER M F and BOYD G S Coronary Atherogenesis—An Endocrine Problem? *A Symposium on Arteriosclerosis* p 64 Minneapolis Minnesota Heart Association and University of Minnesota 1956
- 156 KEYS A and KEYS M H Serum Cholesterol and the Diet in Clinically Healthy Men at Slough Near London *Brit J Nutr* 8 138 1954
- 157 BJORCK G Further Studies in Sweden *World Trends In Cardiology* 1 *Cardiovascular Epidemiology* p 69 New York Hoeber Harper 1956
- 158 MALMROS H BJÖRCK G and SWAEN B Hypertension Atherosclerosis and the Diet *Acta Med Scand* 154 Supp 312 71 1956
- 159 KEYS A ANDERSON J T ARESU M BJÖRCK G BROCK J F BRONTE STEWART B FIDANZA F KEYS M H MALMROS H POPPI A POSTELI T SWAEN B and DEL VECCHIO A Physical Activity and the Diet in Populations Differing in Serum Cholesterol *J Clin Invest* 35 1173 1956
- 160 MULLER C USTVEDT H J *et al* Myocardial Infarction—An Epidemiologic and Prognostic Study of Patients from Five Departments of Internal Medicine in Oslo 1935–1949 *Acta Med Scand* 154 Supp 315 1956
- 161 BJÖRCK G Wartime Lessons on Arteriosclerotic Heart Disease from Northern Europe *World Trends In Cardiology* I *Cardiovascular Epidemiology* p 8 New York Hoeber Harper 1956
- 162 SCHORNAGEL H E The Connection between Nutrition and Mortality from Coronary Sclerosis during and after World War II *Docum Med Geog et Trop* 5 173 1953
- 163 VARTRANEN I and KANERVA K Arteriosclerosis and Wartime *Ann Med Int Fenniae* 36 748 1947
- 164 MALMROS H The Relation of Nutrition to Health—A Statistical Study of Effect of the War Time on Arteriosclerosis Cardiosclerosis Tuberculosis and Diabetes *Acta Med Scand* 138 Supp 246 137 1950
- 165 STROM A and JENSEN R A Mortality from Circulatory Diseases in Norway 1940–1945 *Lancet* 1 126 1951
- 166 GROEN J TJONG B K and HAMMINGA C E The Influence of Nutrition Individuality and Some Other Factors Including Various Forms of Stress on the Serum Cholesterol an Experiment of Nine Months Duration in 60 Normal Human Volunteers *Voeding* 13 556 1957
- 167 JOLLIFFE N Fats Cholesterol and Coronary Heart Disease New York State J Med 57 2684 1957
- 168 KEYS A BROZEK J HENSCHEL A MICHELSON O and TAYLOR H L *The Biology of Human Starvation* Minneapolis University of Minnesota Press 1950
- 169 Malnutrition and Starvation in Western Netherlands—September 1944 July 1945 Parts I and II General State Printing Office The Hague 1948
- 170 A CHOFF L *Lectures in Pathology* New York Hoeber Harper 1974
- 171 DOCK W *Heart Sounds Cardiac Pulsations and Coronary Disease* Lawrence Kan University of Kansas Press 1956
- 172 MORIYAMA I M and GOVER M Statistical Studies of Heart Diseases I Heart Diseases and Allied Causes of Death in Relation to Age Changes in the Population *Pub Health Rep* 63 537 1948
- 173 WOOLSEY T D and MORIYAMA I M Statistical Studies of Heart Diseases II Important Factors in Heart Disease Mortality Trends *Pub Health Rep* 63 1247 1948
- 174 GOVER M Statistical Studies of Heart Disease III Heart Disease Associated with Other Major Causes of Death as Primary or Contributory Cause *Pub Health Rep* 64 104 1949
- 175 ——— Statistical Studies of Heart Disease IV Mortality from Heart Disease (All Forms) Related to Geographic Section and Size of City *Pub Health Rep* 64 439 1949
- 176 COLLINS S D Statistical Studies of Heart Disease V Illness from Heart and Other Cardiovascular Renal Diseases Recorded in General Morbidity Surveys of Families *Pub Health Rep* 64 1439 1949

- 177 WOOLSEY T D Statistical Studies of Heart Disease VI Age at Onset of Heart and Other Cardiovascular Renal Diseases Pub Health Rep 65 555 1950
- 178 GOVER M and PENNELL M Y Statistical Studies of Heart Disease VII Mortality from Eight Specific Forms of Heart Disease among White Persons Pub Health Rep 65 819 1950
- 179 PENNELL M Y and LEHMAN J L Statistical Studies of Heart Disease VIII Mortality from Heart Disease among Negroes as Compared with White Persons Pub Health Rep 66 57 1951
- 180 MORIYAMA I M and WOOLSEY T D Statistical Studies of Heart Disease IX Race and Sex Differences in the Trend of Mortality from the Major Cardiovascular Renal Diseases Pub Health Rep 66 355 1951
- 181 KAUFMAN G and WOOLSEY T D Sex Differences in the Trend of Mortality from Certain Chronic Diseases Pub Health Rep 68 761 1953
- 182 SOWDER W T Why is Sex Difference in Mortality Increasing? Pub Health Rep 69 860 1954
- 183 LEW E A Some Implications of Mortality Statistics Relating to Coronary Artery Disease J Chronic Dis 6 192 1957
- 184 MORIYAMA I WOOLSEY T and STAMLER J Observations on Possible Causative Factors Responsible for the Sex and Race Trends in Cardiovascular Renal Disease Mortality in the United States J Chronic Dis 7 401 1958
- 185 LEE K T and THOMAS W A Myocardial Infarction Changing Sex Ratio and other Factors A M A Arch Int Med 97 421 1956
- 186 ——— Factors Associated with Changing Sex Ratio of Myocardial Infarction A M A Arch Int Med 98 80 1956
- 187 THOMAS W A Fatal Acute Myocardial Infarction Sex Race Diabetes and Other Factors Nutr Revs 15 97 1957
- 188 HUTCHESON J M JR HEJTMANCIK M R and HERRMANN G R Changes in the Incidence and Types of Heart Disease Am Heart J 46 565 1953
- 189 SAPHIR O OHRINGER L and SILVERSTONE H Coronary Arteriosclerotic Heart Disease in the Younger Age Group Its Greater Frequency in this Group among an Increasingly Older Necropsy Population Am J Med Sci 231 494 1956
- 190 WHITE F D Heart Disease a Generation Ago and Now Cardiologia 21 218 1952
- 191 JAMES G PATTON R E and HESLIN A S Accuracy of Cause of Death State Reports on Death Certificates Pub Health Rep 70 39 1955
- 192 KRAUS A S The Use of Hospital Data in Studying the Association between a Characteristic and a Disease Pub Health Rep 69 1211 1954
- 193 MAINLAND D The Risk of Fallacious Conclusions from Autopsy Data on the Incidence of Diseases with Applications to Heart Disease Am Heart J 45 644 1953
- 194 MILLER D C STARE F J WHITE P D and GORDON J E The Community Problem in Coronary Heart Disease A Challenge for Epidemiologic Research Am J Med Sci 232 329 1956
- 195 PAGE I H STARE F J CORCORAN A C POLLACK H and WILKINSON C F JR Atherosclerosis and the Fat Content of the Diet Circulation 16 163 1957
- 196 STAMLER J KJELSBERG M and HALL Y Epidemiological Studies on Cardiovascular Renal Diseases in Chicago and Illinois I Analysis of Mortality Trends by Age Race Sex Occupation In preparation
- 197 KJELSBERG M and STAMLER J Epidemiological Studies on Cardiovascular Renal Diseases in Chicago and Illinois II Analysis of Mortality by Age Race Sex Place of Residence Including Urban Rural Comparisons In preparation
- 198 GLENDY R E LEVINE S A and WHITE P D Coronary Disease in Youth J A M A 109 1175 1937
- 199 ROBERTS D W MEEK P G and KRUEGER D E Prevalence of Heart Disease in the Population of Baltimore Circulation 14 988 1956
- 200 JAMES T N POST H W and SMITH F J Myocardial Infarction in Women Ann Int Med 43 153 1955
- 201 WEINREB H L GERMAN E and ROSENBERG B A Study of Myocardial Infarction in Women Ann Int Med 46 285 1957

- 202 GERTLER M M and WHITE P D *Coronary Heart Disease in Young Adults—A Multidisciplinary Study* Cambridge Commonwealth Fund Harvard University Press 1954
- 203 MASTER A M SIMON D and JAFFE H L Age Sex and Hypertension in Myocardial Infarction due to Coronary Occlusion *Arch Int Med* 64 767 1939
- 204 FORRSSELL J and BRUNILA T Cardiac Infarction and Pregnancy *Acta Med Scand* 157 387 1957
- 205 PEEL A A F Age and Sex Factors in Coronary Heart Disease *Brit Heart J* 17 319 1955
- 206 Anonymous United States Vital Statistics 1951-1952 Prepared by National Office of Vital Statistics USPHS Pub Health Rep 68 68 1953
- 207 Vital Statistics of the United States 1950—Vol III National Office of Vital Statistics Public Health Service U S Department of Health Education and Welfare
- 208 LILLENFELD A M Variation in Mortality from Heart Disease—Race Sex and Socioeconomic Status *Pub Health Rep* 71 545 1956
- 209 DAWBER T R MOORE F E and MANN G V Coronary Heart Disease in the Framingham Study *Am J Pub Health* 47 No 4 Part 2 1957
- 210 CUMMINGS R O *The American and His Food* Chicago University of Chicago Press 1940
- 211 MARTIN E W *The Standard of Living in 1860* Chicago University of Chicago Press 1942
- 212 Consumption of Food in the United States—1909-57 Agriculture Handbook No 62 U S Department of Agriculture Bureau of Agricultural Economics Washington D C 1953
- 213 CLARK F MURRAY J WEISS G S and GROSSMAN E Food Consumption of Urban Families in the United States—with an Appraisal of Methods of Analysis Agriculture Information Bulletin No 132 United States Department of Agriculture Washington D C 1954
- 214 CLARK F and MURRAY J *et al* Household Food Consumption Survey 1955—Reports 1-10 U S Department of Agriculture Washington D C 1957
- 215 COONS G M Family Food Consumption Studies *Pub Health Rep* 67 788 1952
- 216 BABCOCK M J CHURCH H N and GATES L O Nutritional Status of Industrial Workers Effects of Education Age Income and Ethnic Groups *Milbank Mem Fund Quart* 33 137 1955
- 217 MABROUK A F and BROWN J B The Trans Fatty Acids of Margarines and Shortenings *J Am Oil Chemists Soc* 33 98 1956
- 218 SREENIVASAN B and BROWN J B Octadecadienoic Acids of Shortenings and Margarines *J Am Oil Chemists Soc* 33 341 1956
- 219 MURLIN J R Diet of the United States Army Soldier in the Training Camp *JAMA* 71 950 1918
- 220 KINISHI F *et al* The Dietary Composition and Adequacy of the Food Consumed by Soldiers under an Ad Libitum Regimen *Medical Nutrition Laboratory Report* 184 July 1956
- 221 CONSOLAZIO C F *et al* Nutrition Survey of Two Consecutive Training Cycles of the Airborne Training Battalion Co G Ft Benning Ga *Medical Nutrition Laboratory Report* 166 May 1955
- 222 RYER R CONSOLAZIO C F and BERGER F M Report of Nutrition Survey of Two Company Messes 10 Infantry Divisions Fort Riley Kansas *Medical Nutrition Laboratory U S Army Fitzsimmons Army Hospital Report* No 128 April 1954
- 223 SCHOR H C and SWAIN H L Simultaneous Surveys of Food Consumption in Various Camps of the United States Army *J Nutr* 38 51 1949
- 224 Anonymous Bread—White and Brown *Nutr Revs* 14 16 1956
- 225 Editorial Nutrients in Bread *Brit Med J* 1 1223 1956
- 226 McCANCE R A and WIDDOWSON E M *Breads White and Brown Their Place in Thought and Social History* London Pitman Medical Publishing Co Ltd 1956
- 227 ROBINSON C H Fiber in the Diet *Am J Clin Nutr* 4 288 1956

- 228 STAMLER J Epidemiological Investigation of Group Mortality Trends in Chicago due to Arteriosclerotic and Hypertensive Cardiovascular Disease *Circulation* 12 497 1955
- 229 STAMLER J and HALL Y Epidemiological Analysis of Vital Statistics on the Cardiovascular Renal Diseases in Chicago *Circulation* 14 1006 1956
- 230 STAMLER J Problems of the Research Utilization of Vital Statistics on the Cardiovascular Renal Diseases—Draft Report Submitted to the Chicago Heart Association Committee on Morbidity and Mortality Classification of Cardiovascular Disorders 1957
- 231 FRIEDMAN M and ROSENMAN R H Comparison of Fat Intake of American Men and Women—Possible Relationship to Incidence of Clinical Coronary Artery Disease *Circulation* 16 339 1957
- 232 PARKHURST E Differential Mortality in New York State Exclusive of New York City by Sex Age and Cause of Death According to Degree of Urbanization *Am J Pub Health* 46 959 1956
- 233 ENTERLINE P E and STEWART W H Geographic Patterns in Deaths from Coronary Heart Disease *Pub Health Rep* 71 849 1956
- 234 MORRI W H M Heart Disease in Indiana Farmers *Proc First Wisc Conf on Work and the Heart* In Press
- 235 GOVER M Physical Impairments of Members of Low Income Farm Families *Pub Health Rep* 63 1083 1948
- 236 PEMBERTON J and MACLEOD K I Rural Health Survey of Men over Forty *Pub Health Rep* 71 1213 1956
- 237 DOYLE H N Occupational Health on Farms *Pub Health Rep* 72 145 1957
- 238 ADELSON S F and BLAKE E C Diets of Families in the Open Country a Georgia and Ohio County Summer 1945 U S Dept of Agriculture Misc Publication No 704 April 1950
- 239 Anonymous Family Food Consumption in Three Types of Farming Areas of the South II An Analysis of Weekly Food Records Late Winter and Early Spring 1948 Clemson So Car Southern Cooperative Series Bulletin 20 1951
- 240 BABCOCK M J CHURCH H N and GATES I O Nutritional Status of Industrial Workers I Dietary Blood and Physical Findings *Milbank Mem Fund Quart* 32 323 1954
- 241 BILLINGTON L K Dietary Survey Wisconsin Children *Wisc Med J* 54 312 1955 Abstracted by *Am J Clin Nutr* 3 430 1955
- 242 EPPRIGHT E S SIDWELL V D and SWANSON P P Nutritive Value of the Diets of Iowa School Children *J Nutr* 54 371 1954
- 243 WARNICK K P BRING S V and WOODS E Nutritional Status of Adolescent Idaho Children *J Am Dietet Assoc* 31 1143 1955
- 244 Anonymous Diets of Low Income Families *Nutr Revs* 13 109 1955
- 245 BLOCK R J *Amino Acid Handbook* Springfield Ill Charles C Thomas 1956
- 246 HARD M M and ESSELBAUGH N C Nutritional Status of Selected Adolescent Children *Am J Clin Nutr* 4 261 1956
- 247 GILLUM H MORGAN A F and JEROME D W Nutritional Status of the Aging IV Serum Cholesterol and Diet *J Nutr* 55 449 1955
- 248 Anonymous Nutritional Status of an Aged Population *Nutr Revs* 14 10 1956
- 249 DARBY W J MCGANITY W J MARTIN M P BRIDGFORTH E B DENSEN P M KASER M M OGLE P J NEWBILL J A STOCKELL A FERGUSON M E TOUSTER O MCLELLAN G S WILLIAMS C and CANNON R O The Vanderbilt Cooperative Study of Maternal and Infant Nutrition IV Dietary Laboratory and Physical Findings in 2122 Delivered Pregnancies *J Nutr* 51 565 1953
- 250 MCGANITY W J BRIDGFORTH E B MARTIN M P NEWBILL J A and DARBY W J The Vanderbilt Cooperative Study of Maternal and Infant Nutrition VIII Some Nutritional Implications *J Am Dietet Assoc* 31 582 1955
- 251 HARDINGE M G and STAKE F J Nutritional Studies of Vegetarians II Dietary and Serum Levels of Cholesterol *J Clin Nutr* 2 83 1954
- 252 KAYS A and ANDERSON J T Dietary Protein and the Serum Cholesterol Level in Man *Am J Clin Nutr* 5 29 1957

- 253 MIRON L Hemoglobin Level and Dietary Intake of Adults J Clin Nutr 3 38 1954
- 254 Anonymous Estimated Prevalence of Overweight in the United States Pub Health Rep 69 1084 1954
- 255 GORDON T Mortality Experience among the Japanese in the United States Hawaii and Japan Pub Health Rep 72 543 1957
- 256 GILBERT J Absence of Coronary Thrombosis in Navajo Indians California Med 82 114 1955
- 257 PAGE I H LEWIS L A and GILBERT J Plasma Lipids and Proteins and Their Relationship to Coronary Disease among Navajo Indians Circulation 13 675 1956
- 258 HADLEY J N Health Conditions among Navajo Indians Pub Health Rep 70 831 1955
- 259 SMITH R L Cardiovascular Penal and Diabetes Deaths among the Navajos Pub Health Rep 72 33 1957
- 260 STOCKS P Race and Climate as Possible Factors In Cowdry E V Editor Arteriosclerosis New York The Macmillan Co 1933
- 261 LEWIS J H The Biology of the Negro Chicago University of Chicago Press 1942
- 262 SMITH T M Coronary Atherosclerosis in the Negro J Nat Med Assoc 38 193 1946
- 263 BLACHE J O and HANDLER F P Coronary Artery Disease A Comparison of the Rates and Patterns of Development of Coronary Arteriosclerosis in the Negro and White Races and its Relation to Clinical Coronary Artery Disease A M A Arch Path 50 189 1950
- 264 HANDLER F P BLACHE J O and BLUMENTHAL H T Comparison of Aging Processes in the Renal and Splenic Arteries in the Negro and White Races A M A Arch Path 53 29 1952
- 265 GRAY S H HANDLER F P BLACHE J O ZUCKNER J and BLUMENTHAL H T Aging Processes of Aorta and Pulmonary Artery in Negro and White Races A M A Arch Path 56 238 1953
- 266 WEISS M M and GRAY W R Hypertension and Myocardial Infarction in the Negro Am J Med Sci 227 186 1954
- 267 KEIL P G and McVAY L V JR A Comparative Study of Myocardial Infarction in the White and Negro Races Circulation 13 712 1956
- 268 McVAY L V and KEIL P G Myocardial Infarction with Special Reference to the Negro A M A Arch Int Med 96 767 1955
- 269 THOMAS W A BLACHE J O and LEE K I Race and the Incidence of Acute Myocardial Infarction A M A Arch Int Med 100 473 1957
- 270 WEISSLER A M and SHAPIRO O W Alimentary Lipemia and Coronary Artery Disease in Two Racial Groups Am J Med 27 976 1957
- 271 HOLMAN R L MCGILL H C JR STRONG J P GRIFFIN O R and GEER J C Natural History of Atherosclerosis Fed Proc 16 360 1957
- 272 CHAPMAN J M GOERKE L S DIXON W LOVELAND D B and PHILLIPS E The Clinical Status of a Population Group in Los Angeles under Observation for Two to Three Years Am J Pub Health 47 No 4 part 2 33 1957
- 273 BUECKLEY R DUNN J E JR LINDEN G and BRESLOW L Death Certificate Statement of Occupation Its Usefulness in Comparing Mortalities Pub Health Rep 71 1105 1956
- 274 KENT A P MCCARROLL J R SCHWEITZER M D and WILLARD H N Coronary Artery Disease (Arteriosclerotic Heart Disease) Deaths in Manhattan New York City Prepublication Report
- 275 SPAIN D M and BRADSHAW V A Sudden Death from Coronary Atherosclerosis Age Race Sex Physical Activity and Alcohol A M A Arch Int Med 100 228 1957
- 276 STRONG J P TEJADA G MCGILL H C JR and HOLMAN R L Comparison of Early Lesions of Atherosclerosis in New Orleans Guatemala and Costa Rica Circulation 16 516 1957
- 277 GROOM D McNEE E E PEAN V HOUDICOURT E WEBB C and GRANT F W Coronary and Aortic Atherosclerosis Comparative Incidence in the Negro Race in Haiti and the United States Circulation 16 888 1957

- 278 DOYLE J T HESLIN A S HILLEBOE H F FORMEL P F and KORN S R F A Prospective Study of Degenerative Cardiovascular Disease in Albany Report of Three Years Experience—I Ischemic Heart Disease *Am J Pub Health* 47 No 4 Part 2 25 1957
- 279 EDER H A The Lipoproteins of Human Serum *Am J Med* 23 269 1957
- 280 MANN G V The Variability of Serum Lipoproteins and Cholesterol in Normal and Diseased People *Proc 7th Ann Conf on the Nephrotic Syndrome* 1955 New York National Nephrosis Foundation Inc 1956
- 281 GOFMAN J W *et al* and ANDRUS F C *et al* Evaluation of Serum Lipoprotein and Cholesterol Measurements as Predictors of Clinical Complications of Atherosclerosis Report of a Cooperative Study of Lipoproteins and Atherosclerosis *Circulation* 14 691 1956
- 282 ALBRITTON E C Editor *Standard Values in Nutrition and Metabolism* Philadelphia W B Saunders Co 1954
- 283 GOFMAN J W and JONES H B Obesity Fat Metabolism and Cardiovascular Disease *Circulation* 5 514 1952
- 284 ARMSTRONG D B DUBLIN L I WHEATLEY G M and MARKS H H Obesity and Its Relation to Health and Disease *JAMA* 147 100 1951
- 285 SHEPARD W P and MARKS H H Life Insurance Looks at the Arteriosclerosis Problem *A Symposium on Arteriosclerosis* p 6 Minneapolis Minnesota Heart Association and University of Minnesota 1956

E The Clinical Pathological Findings

- 286 HARTROFT W S and THOMAS W A Pathological Lesions Related to Disturbances of Fat and Cholesterol Metabolism in Man *JAMA* 164 1899 1951
- 287 McCANDLESS E L and ZILVERSMIT D B Aortic Synthesis of Plaque Phospholipids in the Eviscerated Nephrectomized Cholesterol Fed Rabbit *Circulation* 16 483 1957
- 288 ZILVERSMIT D B McCANDLESS E I and SHORE M L Phospholipide Metabolism in Various Tissues of Cholesterol Fed Rabbits *Proc Soc Exp Biol and Med* 89 48 1955
- 289 SHORE M L ZILVERSMIT D B and ACKERMAN R F Plasma Phospholipide Deposition and Aortic Phospholipide Synthesis in Experimental Atherosclerosis *Am J Physiol* 181 527 1955
- 290 BLUMGART H L Coronary Disease Clinical Pathological Correlations and Physiology *Bull New York Acad Med* 27 693 1951
- 291 ZOLL P M WESSLER S and BLUMGART H L Angina Pectoris—A Clinical and Pathological Correlation *Am J Med* 11 331 1951
- 292 SPAIN D M BRADESS V A and HUSS G Observations on Atherosclerosis of Coronary Arteries in Males under Age of 46 Necropsy Study with Special Reference to Somatotypes *Ann Int Med* 38 251 1953
- 293 WHITE N K EDWARDS J E and DRY T J The Relationship of the Degree of Coronary Atherosclerosis with Age in Men *Circulation* 1 645 1950
- 294 EDWARDS J E Correlations in Coronary Artery Disease *Bull New York Acad Med* 33 199 1957
- 295 YOUNG W GOFMAN J W MALAMUD N and SIMON A The Interrelationship between Cerebral Atherosclerosis and Coronary Atherosclerosis *Geriatrics* 11 413 1956
- 296 ROBERTS J C JR and MOSES C The Distribution of Atherosclerosis in 200 Autopsies with Special Reference to the Location and Severity of the Vascular Lesions *Circulation* 14 485 1956
- 297 FOLEY W T McDEVITT E and WRIGHT I S Incidence of Combined Arteriosclerosis in the Arteries of the Brain and the Heart *Circulation* 16 882 1957
- 298 ROBERTS J C JR and MOSES C Development of Atherosclerosis in the Major Cerebral Arteries *Circulation* 16 483 1957
- 299 YATER W M TRALM A H BROWN W G FITZGERALD R P GEISLER M A and WILCOX B B Coronary Artery Disease in Men Eighteen to Thirty Nine Years of Age *Am Heart J* 36 334 481 683 1948

- 300 ENOS W F JR, HOLMES R H and BEYER J Coronary Disease among United States Soldiers Killed in Action in Korea *JAMA* 152 1090 1953
- 301 JONES R J, COHEN L and CORBUS H The Serum Lipid Pattern in Hyperthyroidism Hypothyroidism and Coronary Atherosclerosis *Am J Med* 19 71 1955
- 302 MARFORI SAVINI L, VUTERINI S, MAGALINI S and COLTORTI M The Serum Lipoproteins (by Salting out with Sodium Sulphate) in Subjects Suffering from Coronary Atherosclerosis *Acta Med Scand* 146 148 1953
- 303 SPAIN D M, BRADESS V A and GREENBLATT J J Postmortem Studies on Coronary Atherosclerosis Serum Beta Lipoproteins and Somatotypes *Am J Med Sci* 229 294 1955
- 304 GOTTFRIED S P, POPE R H, FRIEDMAN N H, AKERSON I B and DiMAURO S Lipoprotein Studies in Atherosclerotic and Lipemic Individuals by Means of Paper Electrophoresis *Am J Med Sci* 229 34 1955
- 305 EDER H A Plasma Lipoproteins in Atherosclerosis and Related Diseases *A Symposium on Atherosclerosis* p 228 National Academy of Sciences-National Research Council Washington D C 1954
- 306 LITTLE J A, SHANOFF H M, VAN DER FLIER R W and RYKERT H E Serum Lipid Fractionations in Selected Atherosclerotic and Normal Males *Circulation* 14 500 1956
- 307 CHAPMAN I, GOLDBLOOM A A, MIRRELL G and EIBER H B Autopsy Correlation with Clinically Determined Atherogenic Index *AMA Arch Path* 61 357 1956
- 308 PATERSON J C, CORNISH B R and ARMSTRONG E C The Serum Lipids in Human Atherosclerosis—An Interim Report *Circulation* 13 224 1956
- 309 ——— The Gofman Indices in Coronary Atherosclerosis *Canadian Med Assoc J* 74 538 1956
- 310 BJÖRCK G, BLUMQUIST G and SIEVERS J Cholesterol Values in Patients with Myocardial Infarction and in a Normal Control Group *Acta Med Scand* 156 493 1957
- 311 ROSENFELD L Macromolecular Lipoproteins in Atherosclerosis A Review *Am J Clin Nutr* 5 286 1957
- 312 TAMPLIN A R, STRISOWER B, DELALLA O F, GOFMAN J W and GLAZIER F W Lipoproteins Aging and Coronary Artery Disease *J Geront* 9 403 1954
- 313 GOFMAN J W, GLAZIER F W, TAMPLIN A R, STRISOWER B and DELALLA O F Lipoproteins Coronary Heart Disease and Atherosclerosis *Physiol Revs* 34 589 1954
- 314 GOFMAN J W Serum Lipoproteins and the Evaluation of Atherosclerosis *Ann New York Acad Sci* 64 590 1956
- 315 KATZ L B, RHODES G J, GEORGE R S and MOSES C Total Serum Cholesterol Cholesterol Lipid Phosphorus Ratio and Sr^{125}I Concentration in Hypertension Diabetes and Coronary Artery Disease *Am J Med Sci* 25 120 1953
- 316 OLIVER M F and BOYD G S The Plasma Lipids in Coronary Artery Disease *Brit Heart J* 15 387 1953
- 317 ——— Serum Lipoprotein Patterns in Coronary Sclerosis and Associated Conditions *Brit Heart J* 17 299 1955
- 318 GOTTFRIED S P, FRIEDMAN N H, AKERSON I B, POPE R H and DiMAURO S The Relation between Blood Organic Acid Soluble Phosphate Fractions Citric Acid Protein Fractions and Lipid Fractions in Atherosclerotic Heart Diseases and in Lipemias *Circulation* 10 271 1954
- 319 WALKER W J, LAWRY E Y, LOVE D E, MANN G V, LEVINE S A and STARE F J Effect of Weight Reduction and Caloric Balance on Serum Lipoprotein and Cholesterol Levels *Am J Med* 14 654 1953
- 320 WALKER W J Relationship of Adiposity to Serum Cholesterol and Lipoprotein Levels and their Modification by Dietary Means *Ann Int Med* 39 405 1953
- 321 MANN G V and STARE F J Nutrition and Atherosclerosis *A Symposium on Atherosclerosis* p 169 National Academy of Sciences-National Research Council Washington D C 1954 NAS-NRC Publication No 338 1954

- 322 MANN G V, TEEL K, HAYES O, McNALLY A and BRUNO D Exercise in the Disposition of Dietary Calories Regulation of Serum Lipoprotein and Cholesterol Levels in Human Subjects *New Eng J Med* 253 349 1955
- 323 ANDERSON J T, LAWLER A and KEYS A Weight Gain from Simple Overeating II Serum Lipids and Blood Volume *J Clin Invest* 36 81 1957
- 324 MANN G V Lack of Effect of High Fat Intake on Serum Lipid Levels *Am J Clin Nutr* 3 230 1955
- 325 KEYS A The Claim of Lack of Effect of High Fat Intake on Serum Lipid Levels *Am J Clin Nutr* 4 74 1956
- 326 WALDRON J M, BEIDELMAN B and DUNCAN C G The Local and Systemic Effects of Cream on Blood Coagulation A Physiological Basis for Early Feeding in Gastrointestinal Bleeding *Gastroenterology* 17 360 1951
- 327 WALDRON J M and NICHOLS W Plasma Lipids and Whole Blood Clotting Time *Am J Physiol* 171 776 1952
- 328 WALDRON J M and DUNCAN C G Variability of the Rate of Coagulation of Whole Blood *Am J Med* 17 365 1954
- 329 TULLOCH J A, OVERMAN R S and WRIGHT I S Failure of Ingestion of Cream to Affect Blood Coagulation *Am J Med* 14 674 1953
- 330 FULLERTON H W, DAVIE W J A and ANASTASOPOULOS G Relationship of Alimentary Lipemia to Blood Coagulability *Brit Med J* 2 250 1953
- 331 MANNING P R and WALFORD R L Lack of Effect of Fat Ingestion on Blood Coagulation *Am J Med Sci* 228 652 1954
- 332 CULLIN C F and SWANK R L Intravascular Aggregation and Adhesiveness of the Blood Elements Associated with Alimentary Lipemia and Injections of Large Molecular Substances *Circulation* 9 335 1954
- 333 POOLE J C F The Effect of Certain Fatty Acids on the Coagulation of Plasma *In Vitro* *Brit J Exper Path* 36 248 1955
- 334 O'BRIEN J R Relation of Blood Coagulation to Lipaemia *Lancet* 2 690 1955
- 335 MUSTARD J F Increased Activity of the Coagulation Mechanism during Alimentary Lipaemia Its Significance with Regard to Thrombosis and Atherosclerosis *Canad Med Assoc J* 77 308 1957
- 336 O'BRIEN J R Effect of a Meal of Eggs and Different Fats on Blood Coagulability *Lancet* 2 232 1956
- 337 HALL G H Reassessment of Effect of Fatty Meals on Blood Coagulability *Brit Med J* 2 207 1956
- 338 MACLAGAN N F and BILLIMORIA J D Food Lipids and Blood Coagulation *Lancet* 2 235 1956
- 339 BARKHAN P, NEWLANDS M J and WILD F Phospholipids from Brain Tissue as Accelerators and Inhibitors of Blood Coagulation *Lancet* 2 234 1956
- 340 GREIG H B W Etiology of Atherosclerosis *Nature* 178 422 1956
- 341 BUZINA R. and KEYS A Blood Coagulation after a Fat Meal *Circulation* 14 854 1956
- 342 LASCH H G and SCHIMPF K *Blutgerinnung und Alimentäre Fettbelastung* *Deutsch Arch f klin Med* 203 146 1956
- 343 WATSON W C Lipaemia Heparin and Blood Viscosity *Lancet* 2 366 1957
- 344 HIGGINBOTHAM A C and WILLIAMS A V Intravascular Agglutination of the Blood Following Fat Ingestion *J South Carolina Med Assoc* 53 1 1957
- 345 MAYER G A, CONNELL W F and BEVERIDGE J M R Effects of Homogeneous Formula Diets on the Clotting Time of Whole Blood *Circulation* 16 511 1957
- 346 WALKER W J and WIER J A Plasma Cholesterol Levels during Rapid Weight Reduction *Circulation* 3 864 1951
- 347 WALKER W J, WEINER N and MILCH L J Differential Effect of Dietary Fat and Weight Reduction on Serum Levels of Beta Lipoprotein *Circulation* 15 31 1955
- 348 POMERANZE J, BEINFELD W H and CHESSIN M Serum Lipid and Fat Tolerance Studies in Normal Obese and Atherosclerotic Subjects *Circulation* 10 42 1954
- 349 THANNHAUSER S J and MAGENDANTZ H The Different Clinical Groups of Xanthomatous Diseases a Clinical Physiological Study of 22 Cases *Ann Int Med* 2 1662 1938

- 350 STARKE H Effect of the Rice Diet on the Serum Cholesterol Fractions of 154 Patients with Hypertensive Vascular Disease *Am J Med* 9 494 1950
- 351 GOFMAN J W TAMPLIN A and STRISOWER B Relation of Fat and Caloric Intake to Atherosclerosis *J Am Dietet Assoc* 30 317 1954
- 352 PETERS J P and VAN SLYKE D D *Quantitative Clinical Chemistry Vol I Interpretations* 2nd Ed Baltimore The Williams and Wilkins Co 1946
- 353 KEMPNER W Treatment of Hypertensive Vascular Disease with Rice Diet *Am J Med* 4 545 1948
- 354 KEYS A MICKELSEN O MILLES E V O and CHAPMAN C B The Relation in Man between Cholesterol Levels in the Diet and in the Blood *Science* 117 19 1950
- 355 WATKIN D M FROEB H F HATCH F T and GUTMAN A B Effect of Diet in Essential Hypertension II Results with Unmodified Kempner Rice Diet in Fifty Hospitalized Patients *Am J Med* 9 441 1950
- 356 MELLINKOFF S M MACHELLA T E and REINHOLD J G The Effect of a Fat Free Diet in Causing Low Serum Cholesterol *Am J Med Sci* 220 203 1950
- 357 HILDRETH E A MELLINKOFF S M BLAIR G W and HILDRETH D M The Effect of Vegetable Fat Ingestion on Human Serum Cholesterol Concentration *Circulation* 3 641 1951
- 358 HILDRETH E A HILDRETH D M and MELLINKOFF S M Principles of a Low Fat Diet *Circulation* 4 899 1951
- 359 URBACH F HILDRETH E A and WACKERMAN M T The Therapeutic Uses of Low Fat Low Cholesterol Diets I Treatment of Essential Familial Xanthomatosis *J Clin Nutr* 1 52 1952
- 360 WILMOT V A and SWANK R L The Influence of Low Fat Diet on Blood Lipid Levels in Health and in Multiple Sclerosis *Am J Med Sci* 223 25 1952
- 361 HATCH F ABELL L and KENDALL F Effects of Restriction of Dietary Fat and Cholesterol upon Serum Lipids and Lipoproteins in Patients with Hypertension *Am J Med* 19 48 1955
- 362 WILKINSON C F JR BOYLE E JACKSON R S and BENJAMIN M R The Effect of Varying the Intake of Dietary Fat and the Ingestion of Sitosterol on the Lipid Fractions of Human Serum *Circulation* 8 444 1953
- 363 ——— Effect of Varying the Intake of Dietary Fat and the Ingestion of Sitosterol on Lipid and Lipoprotein Fractions of Human Serum *Metabolism* 4 302 1955
- 364 KEYS A ANDERSON J T FIDANZA F KEYS M H and SWANK B Effects of Diet on Blood Lipids in Man—Particularly Cholesterol and Lipoproteins *Clin Chem* 1 34 1955
- 365 MALMROS H and WIGAND C Treatment of Hypercholesteremia *A Symposium on Arteriosclerosis* p 114 Minneapolis Minnesota Heart Association and University of Minnesota 1955
- 366 MORRISON L M A Nutritional Program for Prolongation of Life in Coronary Atherosclerosis *JAMA* 159 1475 1955
- 367 LYON T P YANLEY A GOFMAN J W and STRISOWER B Lipoproteins and Diet in Coronary Heart Disease *California Med* 84 325 1956
- 368 NELSON A M Treatment of Atherosclerosis by Diet Part I Results in Patients Followed from 36 to 72 Months *Northwest Med* 55 643 1956 Abstracted by *JAMA* 161 1416 1956
- 369 VAN HANDEL E and NEUMANN H A Diet Restricted in Refined Cereals and Saturated Fats Its Effect on the Serum Lipid Level of Atherosclerotic Patients *Lancet* 1 245 1957
- 370 NICHOLS A V DOBBIN V and GOFMAN J W The Influence of Dietary Factors upon Human Serum Lipoprotein Concentrations *Geriatrics* 12 7 1957
- 371 ENGELBERG H Studies of Serum Cholesterol and Low Density Lipoprotein Levels Previously Lowered by a Reduced Fat Intake after the Addition of Corn Oil to the Diet *J Chronic Dis* 6 229 1957
- 372 AHRENS E H JR HIRSCH J INSULL W JR TSALTA T T BLOMSTRAND R and PETERSON M L Dietary Control of Serum Lipids in Relation to Atherosclerosis *JAMA* 164 1905 1957

- 373 STARE F J VAN ITALLIE T B MCCANN M B and PORTMAN O W Therapeutic Implications of Nutritional Studies Relating to the Serum Lipids and Atherosclerosis *JAMA* 164 1920 1957
- 374 ROBINSON C H Dietotherapy The Low Cholesterol Low Fat Diet *Am J Clin Nutr* 2 353 1954
- 375 KEYS A ANDERSON J T MICHELSEN O ADERSON S F and FIDANZA F Diet and Serum Cholesterol in Man Lack of Effect of Dietary Cholesterol *J Nutr* 59 39 1956
- 376 MAYER G A CONNELL W F DeWOLFE M S and BEVERIDGE J M R Diet and Plasma Cholesterol Levels *Am J Clin Nutr* 2 316 1954
- 377 BEVERIDGE J M R CONNELL W F MAYER G A FIRSTBROOK J B and DeWOLFE M S The Effects of Certain Vegetable and Animal Fats on the Plasma Lipids of Humans *J Nutr* 56 311 1955
- 378 BEVERIDGE J M R CONNELL W F and MAYER G A Dietary Factors Affecting the Level of Plasma Cholesterol in Humans The Role of Fat *Canadian J Biochem & Physiol* 34 441 1956
- 379 BRONF STEWART B ANTONIS A EALES L and BROCK J F Effects of Feeding Different Fats on Serum Cholesterol Level *Lancet* 1 521 1956
- 380 HORLICK L Effect of Long Chain Polyunsaturated and Saturated Fatty Acids on the Blood Lipids in Man *Circulation* 16 491 1957
- 381 KINSELL L W PARTRIDGE J BOLING L MARGEN S and MICHAELS G Dietary Modification of Serum Cholesterol and Phospholipid Levels *J Clin Endocrinol* 12 909 1952
- 382 KINSELL L W Effects of High Fat Diets on Serum Lipids *J Am Dietet Assoc* 30 685 1954
- 383 KINSELL L W MICHAELS G D COCHRANE G C PARTRIDGE J W JAHN J J and BALCH H E Effect of Vegetable Fat on Hypercholesterolemia and Hyperphospholipemia *Diabetes* 3 113 1954
- 384 KINSELL L W MICHAELS G D PARTRIDGE J W BOLING L A BALCH H E and COCHRANE G C Effect upon Serum Cholesterol and Phospholipids of Diets Containing Large Amounts of Vegetable Fat *J Clin Nutr* 1 224 1953
- 385 COCHRANE G C MICHAELS G D and KINSELL L W Dietary Modifications of Plasma Cholesterol and Phospholipid Levels in Diabetic Patients The Effects of Mixed Diets High in Vegetable Fat *J Clin Nutr* 1 295 1953
- 386 KINSELL L W Some Thoughts Regarding Obesity *Am J Clin Nutr* 2 350 1954
- 387 KINSELL L W and MICHAELS G D Letter to the Editor—In Reply to Comments in Nutrition Reviews *Am J Clin Nutr* 3 247 1955
- 388 ——— Hormonal Nutritional Lipid Relationships *Fed Proc* 14 661 1955
- 389 BEVERIDGE J M R CONNELL W F MAYER G A FIRSTBROOK J B and DeWOLFE M S The Effects of Certain Vegetable and Animal Fats on Plasma Lipids of Humans *Circulation* 10 593 1954
- 390 BEVERIDGE J M R CONNELL W F and MAYER G A The Nature of the Plasma Cholesterol Elevating and Depressant Factors in Butter and Corn Oil *Circulation* 14 484 1956
- 391 SHAPIRO S L and FREEDMAN L Effect of Essential Unsaturated Fatty Acids and Methionine on Hypercholesterolemia *Am J Physiol* 181 441 1955
- 392 AHRENS E H JR BLANKENHORN D H and TSALTAS T T Effect on Human Serum Lipids of Substituting Plant for Animal Foods in Diet *Proc Soc Exper Biol and Med* 86 872 1954
- 393 AHRENS E H JR INSULL W BLONSTRAND R HIRSCH J TSALTAS T T and PETERSON M L The Influence of Dietary Fats on Serum Lipid Levels in Man *Lancet* 1 913 1957
- 394 KEYS A ANDERSON J T and GRANDE F Essential Fatty Acids Degree of Unsaturation and the Effect of Corn (Maize) Oil on the Serum Cholesterol in Man *Lancet* 1 66 1957
- 395 ANDERSON J T KEYS A and GRANDE F The Effects of Different Food Fats on Serum Cholesterol Concentration in Man *J Nutr* 62 421 1957
- 396 KEYS A ANDERSON J T and GRANDE F (Letter to the Editor) *Lancet* 1 992 1957

- 397 ANDERSON J T GRANDE F and KEYS A Dietary Fat and Serum Cholesterol in Man II The Effect of the Nature of the Dietary Fat Personal communication
- 398 SHAPIRO W ESTES E H JR and HILDERMAN H Corn Oil Lowers Cholesterol Levels Dietary Study in Volunteers Shows Report of paper delivered before the Southern Section Meeting of the American Federation for Clinical Research Scope p 5 March 1957
- 399 TOBIAN L and TUNA N The Efficacy of Corn Oil in Lowering the Serum Cholesterol of Patients with Coronary Atherosclerosis Clin Res Proc 5 182 1957
- 400 MALMROS H and WIGAND G The Effect on Serum Cholesterol of Diets Containing Different Fats Lancet 2 1 1957
- 401 MANN G V Essential Fatty Acids and Atherosclerosis a Critique of the Present Knowledge AMA Arch Int Med 100 77 1957
- 402 STEINER A VARSON A and RUDMAN D Effect of a Formula Diet Containing Various Vegetable Oils upon the Serum Lipids of Human Subjects Circulation 16 495 1957
- 403 FARQUHAR J W and SOKOLOW M A Comparison of the Effects of Beta Sitosterol and Safflower Oil Alone and in Combination on Serum Lipids of Humans A Long Term Study Circulation 16 494 1957
- 404 POMERANZE J GOMLIN A and SLOBODY L B Effect of Dietary Fat on Serum Cholesterol Levels in Infancy and Childhood Circulation 16 481 1957
- 405 KINSELL L W MICHAELS G D and DAILEY J P Effects of Ethyl Linoleate Ethyl Oleate Triolein and of a Phosphatide Mixture Containing Tetraoleic Acid upon Fatty Acid Composition of Plasma Lipids in Normal and Abnormal Subjects Circulation 16 479 1957
- 406 KEYS A ANDERSON J T and GRANDE F Serum Cholesterol Response to Natural and Hydrogenated Fats Circulation 16 480 1957
- 407 SHAPIRO W ESTES E H JR and HILDERMAN H L Serum Total Cholesterol Response to Corn Oil Feeding in Patients with Acute Coronary Occlusion Circulation 16 937 1957
- 408 BOZIAN R C LAUFER I J STUTMAN L J HIRSCHHORN K and WILKINSON C F Jr Clinical Studies with Unsaturated Fats in Patients with Hypercholesterolemia and Hyperlipemia Circulation 16 494 1957
- 409 MOSES C JABLONSKI J R DANOWSKI T S and KUNKELE M H Changes in Lipid and Lipoprotein Partition Induced by Feeding a Skim Milk Corn Oil Mixture Circulation 16 511 1957
- 410 NOTHMAN M M BELLIN L and PROGER S Effect of Certain Unsaturated Fatty Acids on Serum Lipids Circulation 16 920 1957
- 411 GORDON E S Blood Cholesterol Lowering Effect of Essential Fatty Acid Supplements in Diabetic Patients J Lab Clin Med 50 818 1957
- 412 KRAMAR J and LEVINE V E Influence of Fats and Fatty Acids on the Capillaries J Nutr 50 149 1953
- 413 BLAKE T M Intramural Hemorrhage in Coronary Arteries Circulation 16 496 1957
- 414 BEVERIDGE J M R CONNELL W F MAYER G A and HAUST H Further Assessment of the Role of Sitosterol in Accounting for the Plasma Cholesterol Depressant Action of Corn Oil Circulation 16 491 1957
- 415 KINSELL L W FRISKEY R W MICHAELS G D and BROWN F R Effect of a Synthetic Triglyceride on Lipid Metabolism Am J Clin Nutr 4 285 1956
- 416 HARRIS R S HOLMAN R T HANSEN A E WIESE H F and EMERSON G A Essential Fatty Acids In *The Vitamins* Vol 2 p 267 Sebrell W H Jr and Harris R S Editors New York Academic Press Inc 1954
- 417 DEUEL H J JR Fat as a Required Nutrient of the Diet Fed Proc 14 639 1955
- 418 DEUEL H J JR and REISER R The Physiology and Biochemistry of the Essential Fatty Acids *Vitamins and Hormones* 13 29 1955
- 419 RATHMANN D M *Vegetable Oils in Nutrition* New York Corn Products Refining Company Publisher 1957
- 420 DEUEL H J JR *The Lipids—Their Chemistry and Biochemistry* Vols I and II New York Interscience Publishers Inc 1951 and 1955

- 421 KEYS A Deficiency of Essential Fatty Acids *Lancet* 1 516 1956
- 422 YUDKIN J Deficiency of Essential Fatty Acid *Lancet* 1 506 1956
- 423 HIRSCH J INSULL W JR and AHRENS E H JR A Chromatographic Study of the Chemical Structure of Human Serum Lipids *J Lab Clin Med* 50 826 1957
- 424 HELLMAN L ROSENFELD R S INSULL W JR and AHRENS E H JR Regulation of Plasma Cholesterol Levels by Fecal Sterol Excretion *Circulation* 16 497 1957
- 425 FRANTZ I D JR and CAREY J B JR Observations on the Pathways of Cholesterol Excretion in Man *J Lab Clin Med* 50 814 1957
- 426 SINCLAIR H M Deficiency of Essential Fatty Acids and Atherosclerosis Etcetera *Lancet* 1 381 1956
- 427 CORNWELL D G BACKDERF R WILSON C L and BROWN J B The Trans Octadecanoic Acid Content of Butter Fat *Arch Biochem and Biophys* 46 364 1953
- 428 AAS JØRGENSEN E ENGEL P F FUNCH J P and DAM H The Role of Fat in the Diet of Rats 7 The Influence on Growth of Diets Supplemented with Raw Skim Milk Linoleic Acid or Both and of Raw Casein Compared with Alcohol Extracted Casein *Brit J Nutr* 9 42 1955
- 429 PRIVETT O S PUSCH B J and LUNDBERG W O Fatty Acid Composition and Oxidative Deterioration during Storage of Fats in Cuts of Beef Lamb Pork and Turkey Food Technology 9 347 1955
- 430 JAMES A T LOVELOCK J E WEBB J and TROTTER W R The Fatty Acids of the Blood in Coronary Disease *Lancet* 1 105 1957
- 431 HUNDLEY J M Diabetes Overweight U S Problems *J Am Dietet Assoc* 32 417 1956
- 432 JOHNSON M L BURKE B S and MAYER J The Prevalence and Incidence of Obesity in a Cross Section of Elementary and Secondary School Children *Am J Clin Nutr* 4 231 1956
- 433 Editorial The Prevalence of Obesity *Diabetes* 4 155 1955
- 434 BARR D P Health and Obesity *New Eng J Med* 248 967 1953
- 435 DUBLIN L I Relation of Obesity to Longevity *New Eng J Med* 248 971 1953
- 436 MASTER A M JAFFE H L and CHESEKY K Relationship of Obesity to Coronary Disease and Hypertension *JAMA* 157 1499 1953
- 437 Editorial Obesity and Heart Disease *J Chronic Dis* 1 456 1955
- 438 *Weight Control—Colloquium Iowa State College Ames Ia Iowa State College Press* 1955
- 439 LEE K T and THOMAS W A Relationship of Body Weight to Acute Myocardial Infarction *Am Heart J* 52 581 1956
- 440 Anonymous Obesity and Coronary Artery Disease *Nutr Revs* 14 3 1956
- 441 MASTER A M and JAFFE H L Physiologic Effects of Obesity upon Heart Incidence of Obesity in Coronary Disease *J Am Geriatrics Soc* 3 709 1955
- 442 KEYS A Obesity and Heart Disease *J Chronic Dis* 1 456 1955
- 443 POLLACK H Obesity and Heart Disease—A Perennial Problem Modern Concepts of Cardiovascular Dis 26 371 1957
- 444 MASTER A M Hypertension and Coronary Occlusion *Circulation* 8 170 1953
- 445 SIGLER L H The Role of Hypertension in the Etiology and Prognosis of Coronary Occlusion *Ann Int Med* 46 369 1955
- 446 THOMAS C B and COHEN B H The Familial Occurrence of Hypertension and Coronary Artery Disease with Observations Concerning Obesity and Diabetes *Ann Int Med* 47 90 1955
- 447 MASTER A M GARFIELD C I and WALTERS M B *Normal Blood Pressure and Hypertension Philadelphia Lea & Febiger* 1952
- 448 TOBIAN L Hypertension and Atherosclerosis A Symposium on Arteriosclerosis p 54 Minneapolis Minnesota Heart Association and University of Minnesota 1956
- 449 DUCUID J B Pathogenesis of Atherosclerosis *Lancet* 2 925 1949
- 450 MORGAN A D *The Pathogenesis of Coronary Occlusion Oxford Eng Blackwell* 1956

F *The Animal Experimental Findings*

- 451 ANITSCHKOW N Experimental Arteriosclerosis in Animals In Cowdry E V Editor *Arteriosclerosis* New York The Macmillan Co 1933
- 452 STAMLER J and KATZ L N Production of Experimental Cholesterol Induced Atherosclerosis in Chicks with Minimal Hypercholesterolemia and Organ Lipidosis *Circulation* 2 705 1950
- 453 STEINER A and KENDALL F E Atherosclerosis and Arteriosclerosis in Dogs Following Ingestion of Cholesterol and Thiouracil *A M A Arch Path* 42 433 1946
- 454 MANN G V ANDRUS S B McNALLY A and STARE F J Experimental Atherosclerosis in Cebus Monkeys *J Exper Med* 98 195 1953
- 455 FILLIOS L C and MANN G V Influence of Sulfur Amino Acid Deficiency on Cholesterol Metabolism *Metabolism* 3 16 1954
- 456 MANN G V and ANDRUS S B Xanthomatosis and Atherosclerosis Produced by Diet in an Adult Rhesus Monkey *J Lab Clin Med* 48 533 1956
- 457 COX G E NELSON L G and TAYLOR C B Hypercholesterolemia in Monkeys Ingesting an Adequate Diet *Circulation* 10 603 1954
- 458 WISSLER R W EILERT M L SCHROEDER M A and COHEN L Production of Lipomatous and Atheromatous Arterial Lesions in the Albino Rat *A M A Arch Path* 57 333 1954
- 459 WISSLER R W ALLEN R MOY R and BRADFORD W The Relationships between Arterial and Renal Injury and Diet in the Production of Atheromatous Lesions in the Coronary Arteries of the Rat *Circulation* 12 502 1955
- 460 MALINOW M R HOJMAN D and PELLEGRINO A A Production of Atherosclerosis in the Rat an Omnivorous Animal by Forced Feeding of Vegetable Oil Preliminary Report *Circulation* 10 302 1954
- 461 STAMLER J Experimental Atherosclerosis Proc 5th Interamerican Cardiological Congress 1956
- 462 HARTROFT W S and THOMAS W A Production of Coronary Thromboses and Myocardial Infarcts in Rats by Dietary Means *Circulation* 16 481 1957
- 463 WILGRAM G F Calcified Atheromatosis in the Rat *Fed Proc* 16 270 1957
- 464 ——— Fibrinous Type of Atheroma in Rats *Circulation* 16 500 1957
- 465 PICK R KATZ L N ROBBARD S and STAMLER J Influences of Neutral Fat and Plant Sterols on Exogenous and Endogenous Hypercholesterolemia and Hyperphosphipemia in Chicks *Circulation* 10 610 1954
- 466 BRAGDON J H Hyperlipemia and Atheromatosis in a Hibernator *Citellus Columbianus* *Circulation Res* 2 520 1954
- 467 WOLFFE J B HYMAN A S PLUNGIAN M B DALE A D MCGINNIS G E and WALKOW M B Studies in Experimental Atheromatosis Atheromatosis and Hepato Atherosclerosis in Geese Possible Reversibility and Clinical Implications *J Geront* 7 13 1952
- 468 STEINER A and DAYTON S Production of Hyperlipemia and Early Atherosclerosis in Rabbits by a High Vegetable Fat Diet *Circulation Res* 4 62 1956
- 469 LAMBERT G F and SCHMIDT J L Personal Communication
- 470 HORLICK L and KATZ L N Retrogression of Atherosclerotic Lesions on Cessation of Cholesterol Feeding in the Chick *J Lab Clin Med* 34 1427 1949
- 471 STAMLER J PICK R and KATZ L N Effects of Various Dietary Alterations on Cholesterol and Atherogenesis in Cockerels on a High Fat High Cholesterol Ration In preparation
- 472 ROBBARD S PICK R and KATZ L N The Rate of Regression of Hypercholesterolemia and Atherosclerosis in Chicks Effects of Diet Pancreatectomy Estrogens and Thyroid *Circulation* 10 597 1954
- 473 STAMLER J PICK R and KATZ L N Insulin Inhibition of Diet Induced Regression of Coronary Atherosclerosis in Chicks *Circulation* 12 487 1955
- 474 BEVANS M DAVIDSON J D and KENDALL F E Regression of Lesions in Canine Arteriosclerosis *A M A Arch Path* 51 288 1951
- 475 DAUBER D V and KATZ L N Experimental Atherosclerosis in the Chick *A M A Arch Path* 36 473 1943

- 476 STAMLER J PICK R and KATZ L N Effects of Various Fats on Cholesterolemia and Atherogenesis in Cholesterol Fed Cockerels Fed Proc 16 123 1957
- 477 ——— Effects of Oleic Acid on Plasma Lipids and Atherogenesis of Cholesterol Fed Cockerels Circulation 16 483 1957
- 478 VAN HANDEL E and ZILVERSMIT D B Effect of Various Dietary Fats on Blood Lipids of Rabbits Circulation 16 516 1957
- 479 DILLER E R WOODS B L and HARVEY O A Effect of Degree of Unsaturation of Vegetable Lipid upon the Plasma and Hepatic Cholesterol Content in the Rat Circulation 16 505 1957
- 480 AVICAN J and STEINBERG D Effect of Corn Oil Feeding on Cholesterol Metabolism in the Rat Circulation 16 492 1957
- 481 HEGSTED D M GOTSIS A and STARE F J Relation of Oil Composition to Serum Cholesterol Levels in Hypercholesterolemic Rats Circulation 16 479 1957
- 482 KLUMMEROW F A The Relation of Adequate Nutrition to Atherosclerosis Food and Nutrition News 27 1 1955
- 483 KRITCHEVSKY D MOYER A W LOGAN J B and McCANDLESS R F J Effect of State of Administered Cholesterol in Experimental Atherosclerosis Arch Biochem and Biophys 59 526 1955
- 484 KRITCHEVSKY D MOYER A W TESAR W C McCANDLESS R F J LOGAN J B BROWN R A and ENGLERT M E Cholesterol Vehicle in Experimental Atherosclerosis II Influence of Unsaturation Am J Physiol 185 279 1956
- 485 SENKIND C R SCHROEDER M T RASMUSSEN R and WISSLER R W Effect on Serum Lipids in Rats Fed Plant Oils with and without Cholesterol Fed Proc 16 371 1957
- 486 JONES R J REISS O K and HUFFMAN S Corn Oil and Hypercholesteremic Response in the Cholesterol Fed Chick Proc Soc Exper Biol and Med 93 85 1956
- 487 PORTMAN O W STARE F J and BRUNO D Level and Type of Dietary Fat and Experimental Hypercholesteremia in the Cebus Monkey Fed Proc 15 510 1956
- 488 LIN T M KARVINEN E and IVY A C Effect of Certain Fatty Acids on Excretion of Endogenous Cholesterol Am J Physiol 187 173 1956
- 489 OKEY R and STONE M M Lard vs Vegetable Fat in Relation to Liver and Serum Cholesterol J Am Dietet Assoc 32 807 1956
- 490 AFTERGOOD L ALFIV SLATER R B and DEUEL H J JR Comparative Effect of Cottonseed Oil and Lard on Cholesterol Metabolism in Rat Fed Proc 15 541 1956
- 491 KING J S JR CLARKSON T B and WARNOCK N H Effect on Experimental AVIAN Atherosclerosis of Dietary Oils and Gallogens Proc Soc Exper Biol and Med 93 443 1956
- 492 PHIL A The Effect of Dietary Fat on the Intestinal Cholesterol Absorption and on the Cholesterol Metabolism in the Liver of Rats Acta Physiol Scand 34 183 1955
- 493 DAM H PRANGE I and SONDERGAARD E Influence of Various Levels of Dietary Cholesterol on the Cholesterol Content of Certain Organs and of Bile of Chicks Fed Fat Free and Peanut Oil Containing Diets Acta Physiol Scand 34 141 1955
- 494 DAM H KRISTENSEN G KOFOED N PRANGE I and SONDERGAARD E Influence of Varied Levels of Peanut Oil and Cholesterol on Cholesterol and Polyenoic Acids in Tissues of the Chicks Acta Physiol Scand 36 319 1956
- 495 HEGSTED D M ANDRUS S B GOTSIS A and PORTMAN O W The Quantitative Effects of Cholesterol Cholic Acid and Types of Fat on Serum Cholesterol and Vascular Sudanophilia in the Rat J Nutr 63 273 1957
- 496 HEGSTED D M GOTSIS A and STARE F J The Effects of Various Fats upon Experimental Hypercholesteremia in the Rat J Nutr 63 377 1957
- 497 BURR G O and BURR M M A New Deficiency Disease Produced by the Rigid Exclusion of Fat from the Diet J Biol Chem 82 345 1929
- 498 RODRIGUEZ R R CATTANEO P HOLSSAY B A and UNO B Unsaturated Fatty Acids and Alloxan Diabetes J Nutr 51 441 1953
- 499 PEIFER J J and HOLMAN R T Essential Fatty Acids Diabetes and Cholesterol Arch Biochem and Biophys 57 520 1955

- 500 ——— Relation of Dietary Cholesterol to Essential Fatty Acid Deficiency Fed Proc 15 326 1956
- 501 AAES JØRGENSEN E and HOLMAN R T Effects of Cholesterol Cholic Acid and EFA Deficiency upon Adult Male Rats Fed Proc 16 142 1957
- 502 AAES-JØRGENSEN E FRENCH J P ENGEL P F and DAM H Role of Fat in the Diet of Rats Influence of Growth and Histologic Findings of Diets with Hydrogenated Arachis Oil or No Fat Supplemented with Linoleic Acid A Raw Skim Milk and of Crude Casein Compared with Vitamin Test Casein Part 9 Brit J Nutr 10 297 1956
- 503 HOLMAN R T and AAES JØRGENSEN E Effects of *Trans* Fatty Acid Isomers upon Essential Fatty Acid Deficiency in Rats Proc Soc Exper Biol and Med 93 175 1956
- 504 STAMLER J PICK R and KATZ L N Experimental Studies on the Effects of Protein and Carbohydrate on Hypercholesterolemia and Atherogenesis in Cockerels on a High Fat High Cholesterol Ration Clin Res Proc 5 296 1957
- 505 ——— Effects of Dietary Proteins Methionine and Vitamins on Plasma Lipids and Atherogenesis in Cholesterol Fed Cockerels Circulation Res 6 442 1958
- 506 ——— Effects of Dietary Protein and Carbohydrate Level on Cholesterolemia and Atherogenesis in Cockerels on a High Fat High Cholesterol Mash Circulation Res 6 447 1958
- 507 PICK R STAMLER J and KATZ L N Effects of High Protein High Vitamin Supplements on Cholesterolemia and Atherogenesis in Cholesterol Fed Cockerels Fed Proc 16 101 1957
- 508 ——— Effects of Dietary and Hormonal Imbalance on Coronary Atherogenesis in Cholesterol Fed Cockerels Circulation 14 983 1956
- 509 ——— Effects of Dietary Imbalance on Atherogenesis in Cholesterol Fed Cockerels Circulation 14 507 1956
- 510 LOEWE L GOLDNER M G RAPOPORT S M and STERN I Failure of Protein to Protect Against Cholesterol Atherogenesis in Underfed Rabbits Proc Soc Exper Biol and Med 87 360 1954
- 511 KOLATNUR M RAND N T KUMMEROW F A and SCOTT H M Effect of Dietary Protein and Fat on Changes of Serum Cholesterol in Mature Roosters Fed Proc 16 389 1957
- 512 RAND N T SHAH S N ARGOUEDELIS C J and KUMMEROW F A Effect of Dietary Protein on the Serum and Total Cholesterol and the Carcass Fat Composition of Growing Chicks Circulation 16 500 1957
- 513 BROWN H B and LEWIS L A Effect of High Fat and High Protein Diets on Tissue Lipids Circulation 14 488 1956
- 514 JONES R J and HUFFMAN S The Effect of Various Chronic Levels of Dietary Casein on Hypercholesterolemia in the Rat Circulation 14 488 1956
- 515 OLSON R E JABLONSKI J R and TAYLOR E Effect of Dietary Fat Protein and Choline upon Serum Lipids and Lipoproteins of the Rat Fed Proc 16 395 1957
- 516 GRANT W C and FAHRENBAUGH M J Influence of Sucrose and Casein on Plasma Cholesterol in Animals on Purified Diets Fed Proc 16 50 1957
- 517 HSU J M and CHOW B F Vitamin B₂ Deficiency and Hypercholesterolemia Fed Proc 16 63 1957
- 518 EADES C H JR and GALLO D G Effect of Vanadyl Sulfate on Plasma and Tissue Cholesterol Levels and Atherosclerosis in the Chicken Fed Proc 16 176 1957
- 519 TENNENT D M MUSHETT C W KURON G W OTT W H and SIEGEL H Influence of Cobaltous Chloride on Aortic Atheromatosis and Plasma Lipid Pattern in Cholesterol Fed Chickens Fed Proc 16 399 1957
- 520 HELLERSTEIN E E VITALE J J and ZANCHECK N Influence of Dietary Magnesium on Cardiac and Renal Lesions of Weanling Rats Fed an Atherogenic Diet Fed Proc 16 359 1957
- 521 RINEHART J F and GREENBERG L D Vitamin B₆ Deficiency in the Rhesus Monkey with Particular Reference to the Occurrence of Atherosclerosis Dental Caries and Hepatic Cirrhosis Am J Clin Nutr 4 318 1956

- 522 MUSHETT C W and EMERSON G A Arteriosclerosis in Pyridoxine Deficient Monkeys and Dogs Fed Proc 16 367 1951
- 523 SCHROEDER H A Is Atherosclerosis a Conditioned Pyridoxal Deficiency? J Chronic Dis 28 1955
- 524 HARTROFT W S RIDOUT J H SELLERS E A and BEST C H Atheromatous Changes in Aorta Carotid and Coronary Arteries of Choline Deficient Rats Proc Soc Exper Biol and Med 81 384 1952
- 525 HARTROFT W S Effects of Various Types of Lipids in Experimental Hypolipotropic Diets Fed Proc 14 655 1955
- 526 WILGRAM G P BEST C H and BLUMENSTEIN J Aggravating Effect of Cholesterol on Cardiovascular Changes in Choline Deficient Rats Proc Soc Exper Biol and Med 89 476 1955
- 527 VAN HAARLEM E Atherosclerosis Overnutrition or Malnutrition? Am J Digest Dis 22 206 1955
- 528 BELAVADY B and BANERJEE S Metabolism of Cholesterol in Scorbatic Guinea Pigs J Biol Chem 209 641 1954
- 529 SWELL L BOSTER T A FIELD H JR and TREADWELL C R Pantothenate and Dietary Cholesterol in the Maintenance of Blood and Tissue Cholesterol Esters J Nutr 57 121 1955
- 530 MARCH B E and BIELY J Fat Studies in Poultry 3 Folic Acid and Fat Tolerance in the Chick Poultry Sci 34 39 1955
- 531 FOX M R S BRIGGS G M and ORTIS L O High Dietary Fat and the Chick's Requirement for Vitamin B₁₂ Fed Proc 14 433 1955
- 532 SUNDE M L A Relationship Between Protein Level and Energy Level in Chick Rations Poultry Sci 35 350 1956
- 533 MARCH B and BIELY J Fat Studies in Poultry 5 The Effect of Dietary Fat Level on the Choline Requirement of the Chick Poultry Sci 35 545 1956
- 534 ——— Folic Acid Supplementation of High Protein High Fat Diets Poultry Sci 35 550 1956
- 535 FOX H M SWANSON P P and BIRDS E Role of Fat in Metabolism under Stress Conditions Fed Proc 15 331 1956
- 536 GRIMINGER P SCOTT H M and FORBES R M The Effect of Protein Level on the Tryptophan Requirement of the Growing Chick J Nutr 59 67 1956
- 537 KLEINERMAN J KLAYNS J V KOBAYASHI N and MARSTERS R Effects of Ethionine and High Fat Diets with and without Choline Supplementation on Production of Aortic Sclerosis in Rats Fed Proc 16 302 1957
- 538 PARSONS W B JR and FLINN J H Success of Niacin and Failure of Niacinamide in Reducing Plasma Cholesterol Levels in Patients with Hypercholesterolemia Circulation 16 923 1957
- 539 WILGRAM G P LEWIS L A and BLUMENSTEIN J Lipoproteins in Choline Deficiency Circulation Res 3 549 1955
- 540 STAMLER J PICKER R NASH H A BUNDE C A and KATZ L N Corn Germ Inhibition of Hypercholesterolemia and Atherogenesis in Cholesterol Fed Cockerels —Experiments on Mechanism In preparation
- 541 PETERSON D W Effect of Soybean Sterols in the Diet on Plasma and Liver Cholesterol in Chicks Proc Soc Exper Biol and Med 78 143 1951
- 542 PETERSON J E and HIRST A E Studies on the Relation of Diet Cholesterol and Atheroma in Chickens Circulation 3 116 1951
- 543 PETERSON D W NICHOLS C W JR and SHNEOUR E A Some Relationships among Dietary Sterols Plasma and Liver Cholesterol Levels and Atherosclerosis in Chicks J Nutr 47 57 1952
- 544 PETERSON D W SHNEOUR E A PEEK N F and GAFFER H W Dietary Constituents Affecting Plasma and Liver Cholesterol in Cholesterol Fed Chicks J Nutr 50 191 1953
- 545 PETERSON D W SHNEOUR E A and PEEK N F Effects of Dietary Sterols and Sterol Esters on Plasma and Liver Cholesterol in the Chick J Nutr 53 451 1954
- 546 POLLAL O J Successful Prevention of Experimental Hypercholesterolemia and Cholesterol Atherosclerosis in the Rabbit Circulation 7 696 1953

- 547 SWELL L BOITER T A FIELD H JR and TREADWELL C R Esterification of Soybean Sterols *in vitro* and their Influence on Blood Cholesterol Levels Proc Soc Exper Biol and Med 86 295 1954
- 548 ROSENMAN R H BYERS S O and FRIEDMAN M The Effect of Soybean Sterols on the Absorption of Cholesterol by the Rat Circulation Res 2 160 1954
- 549 ALVIN SLATER R B WELLS A F AFTERGOOD L MELNICK D and DEUEL H J JR The Effect of Plant Sterols on Cholesterol Levels in the Rat Circulation Res 2 471 1954
- 550 BURKE K A McCANDLESS R F J and KRITCHEVSKY D Effect of Soybean Sterols on Liver Deposition of Cholesterol C¹⁴ Proc Soc Exper Biol and Med 87 87 1954
- 551 BEHER W T and ANTHONY W L Effects of Beta Sitosterol and Ferric Chloride on Accumulation of Cholesterol in Mouse Liver Proc Soc Exper Biol and Med 90 223 1955
- 552 STAMLER J PICK R. and KATZ L N Effect of Sitosterols on Diet Induced Exogenous and Hormone Induced Endogenous Hypercholesterolemia in Cockerels In preparation
- 553 HERNANDEZ H H PETERSON D W CHAIKOFF I L and DAUBEN W G Absorption of Cholesterol-4 C¹⁴ in Rats Fed Mixed Soybean Sterols and Beta Sitosterol Proc Soc Exper Biol and Med 83 498 1953
- 554 HERNANDEZ H H and CHAIKOFF I L Do Soy Sterols Interfere with Absorption of Cholesterol? Proc Soc Exper Biol and Med 87 541 1954
- 555 GOULD R G Absorbability of Dihydrocholesterol and Sitosterol Circulation 10 589 1954
- 556 DAVIS W W The Physical Chemistry of Cholesterol and Beta Sitosterol Related to the Intestinal Absorption of Cholesterol Trans N Y Acad Sci 18 123 1955
- 557 BISSET S K and COOK R P Faecal Excretion in Man after the Administration of Mixed Plant Sterols (Sitosterols) Biochem J 63 13 1956
- 558 Anonymous Plant Sterols and Cholesterol Absorption Nutr Revs 15 45 1957
- 559 GOULD R G Absorbability of β Sitosterol Trans N Y Acad Sci 18 129 1955
- 560 BEHER W T and ANTHONY W L Effects of Dihydrocholesterol and Soybean Sterols on Elevated Tissue Cholesterol Proc Soc Exper Biol and Med 86 589 1954
- 561 BEHER W T ANTHONY W L and BAKER G D Effects of Beta Sitosterol on Regression of Cholesterol Atherosclerosis in Rabbits Circulation Res 4 485 1956
- 562 SIPERSTEIN M D NICHOLS C W and CHAIKOFF I L Prevention of Plasma Cholesterol Elevation and Atheromatosis in the Cholesterol Fed Bird by the Administration of Dihydrocholesterol Circulation 7 37 1953
- 563 ROSENMAN R H BYERS S O and FRIEDMAN M The Effect of Dihydrocholesterol on the Absorption of Cholesterol by the Rat Circulation Res 2 45 1954
- 564 BEHER W T and ANTHONY W L Prevention of the Cholesterol Type of Fatty Livers in Mice by Dietary Dihydrocholesterol J Nutr 52 519 1954
- 565 IVY A C LIN T and KARVINER E Absorption of Dihydrocholesterol and Soya Sterols by the Rat's Intestine Am J Physiol 183 79 1955
- 566 SWELL L BOITER T A FIELD H JR and TREADWELL C R The Absorption of Plant Sterols and their Effect on Serum and Liver Sterol Levels J Nutr 58 385 1956
- 567 NICHOLS C W JR LINDSAY S and CHAIKOFF I L Production of Arteriosclerosis in Birds by the Prolonged Feeding of Dihydrocholesterol Proc Soc Exper Biol and Med 89 609 1955
- 568 CURRAN G L and COSTELLO R L Effect of Dihydrocholesterol and Soybean Sterols on Cholesterol Metabolism in Rabbit and Rat Proc Soc Exper Biol and Med 91 52 1956
- 569 COOK R P KLIMAN A and FIESER L F The Absorption and Metabolism of Cholesterol and its Main Companions in the Rabbit—with Observations on the Atherogenic Nature of the Sterols Arch Biochem and Biophys 52 439 1954

- 570 TOMKINS G M NICHOLS C W JR CHAPMAN D D HOFFA S and CHAIKOFF I L Use of Δ^4 -Cholestenone to Reduce the Level of Serum Cholesterol in Man *Science* 125 936 1957
- 571 STEINBERG D and FREDRICKSON D S Inhibitors of Cholesterol Biosynthesis and the Problem of Hypercholesterolemia *Ann New York Acad Sci* 64 579 1956
- 572 GORDON R B KRAFT S C and JONES R J The Effect of a Brain Fraction on Serum Cholesterol and the Serum Lipid Pattern *J Lab Clin Med* 41 583 1953
- 573 JONES R J Serum Cholesterol Reduction in Patients by the Oral Administration of a Brain Extract *J Lab Clin Med* 47 261 1957
- 574 STAMLER J PICK R and KATZ L N Effects of Various Brain Fractions on Cholesterolemia and Atherogenesis in Cholesterol Fed Cockerels *In preparation*
- 575 JONES R J KEOUGH T F CUMMINGS D and KRAFT S Factors Determining the Hypercholesteremic Response in Patients Fed a Brain Extract *Circulation* 16 497 1957
- 576 CARROLL H K Effect of Cerebroside Preparations on Excretion of Fecal Cholesterol *Circulation* 16 497 1957
- 577 ROBBARD S BOLENE C and KATZ L N Hypercholesteremia and Atheromatosis in Chicks on a Restricted Diet Containing Cholesterol *Circulation* 4 43 1951
- 578 ROBBARD S BOLENE WILLIAMS C PICK R and KATZ L N The Beneficial Effects of Intermittent Dietary Regimes on the Tendency to Atherosclerosis *J Lab Clin Med* 41 587 1953
- 579 GOLDNER M G LOEWE L LASSER R and STERN I Effect of Calorie Restriction on Cholesterol Atherogenesis in the Rabbit *Proc Soc Exper Biol and Med* 87 105 1954
- 580 LI T and FREEMAN S Experimental Lipemia and Hypercholesterolemia by Protein Depletion and by Cholesterol Feeding in Dogs *Am J Physiol* 145 660 1946
- 581 JONES R J and HUFFMAN S Chronic Effect of Dietary Protein on Hypercholesteremia in the Rat *Proc Soc Exper Biol and Med* 93 519 1956
- 582 MOYER A W KRITCHEVSKY D LOGAN J B and COV H R Dietary Protein and Serum Cholesterol in Rats *Proc Soc Exper Biol and Med* 92 736 1956
- 583 WILLIS G C and FISHMAN S Ascorbic Acid Content of Human Arterial Tissue *Canad Med Assoc J* 72 500 1955
- 584 MOUNTAIN J T STOCKELL F R JR and STOKINGER H E Effect of Ingested Vanadium on Cholesterol and Phospholipid Metabolism in the Rabbit *Proc Soc Exper Biol and Med* 92 582 1956
- 585 CURRAN G L and COSTELLO R L Reduction of Excess Cholesterol in the Rabbit Aorta by Inhibition of Endogenous Cholesterol Synthesis *J Exper Med* 103 49 1956
- 586 PARSONS W B JR ACHOR R W P BERGE H G MCKENZIE B F and BARKER N W Changes in Concentration of Blood Lipids Following Prolonged Administration of Large Doses of Nicotinic Acid to Persons with Hypercholesterolemia Preliminary Observations *Proc Staff Mayo Clin* 31 377 1956
- 587 ——— The Effect of Large Doses of Nicotinic Acid on the Plasma and Serum Lipids of Human Beings with Hypercholesterolemia *Circulation* 14 495 1956
- 588 ALTSCHUL R Influence of Nicotinic Acid on Cholesterol Metabolism *Circulation* 14 494 1956
- 589 ——— Influence of Nicotinic Acid on Serum Cholesterol in Man *Arch Biochem and Biophys* 54 558 1955
- 590 Anonymous Niacin and Blood Cholesterol *Nutr Revs* 15 137 1957
- 591 ACHOR R W P BERGE H G BARKER N W and MCKENZIE B F Treatment of Hypercholesteremia with Nicotinic Acid *Circulation* 16 499 1957
- 592 PARSONS W B JR and FLINN J H Success of Niacin and Failure of Niacinamide in Reducing Plasma Cholesterol Levels in Patients with Hypercholesterolemia *Circulation* 16 499 1957
- 593 ALTSCHUL R and HOFFER A Effect of Large Doses of Nicotinic Acid on Serum Cholesterol and on BMR *Circulation* 16 499 1957

- 594 MERRILL J M and LEMLEY STONE J Prevention of Cholesterol Deposition in the Rabbit Aorta by Oral Nicotinic Acid Circulation 16 915 1957
- 595 FAILEY R B JR. Effect of Large Doses of Pyridoxine on Serum Cholesterol in the Human Circulation 16 506 1957
- 596 GREENBLATT I J Use of Massive Doses of Vitamin E in Humans and Rabbits to Reduce Blood Lipids Circulation 16 508 1957

Hormonal and Other Endogenous Factors and Atherosclerosis

- A The Key but not Exclusive Role of Diet in Atherogenesis Interrelationships between Diet and Other Factors (Heredity Physical Activity Stress Endocrine Functions Smoking and the Like)*
- 597 WILLIS G C Localizing Factors in Atherosclerosis Canad Med Assoc J 70 1 1954
 - 598 MORRIS J N and HARDY J A Mortality in Relation to Physical Activity of Work Brit J Indust Med 10 245 1953
 - 599 KEYS A ANDERSON J T and MICKELSEN O Serum Cholesterol in Men in Basal and Nonbasal States Science 123 29 1956
 - 600 KEYS A and BUZINA R Blood Coagulability Effects of Meals and Differences Between Populations Circulation 14 479 1956
 - 601 WARNOCK N H CLARKSON T B and STEVENSON R Effect of Exercise on Blood Coagulation Time and Atherosclerosis of Cholesterol Fed Cockerels Circulation Res 5 478 1957
 - 602 WONG H Y C ANDERSON M B KIM J K LIU D J and HAWTHORNE E W Hypocholesterolizing Effect of Exercise on Cholesterol Fed Cockerels Fed Proc 16 138 1957
 - 603 PELTONEN L and KARVONEN M Effects of Dietary Cholesterol Dietary Fat and Exercise on Mouse Plasma Cholesterol Ann Med Exper Biol Fenniae 34 246 1956
 - 604 BROWN C E HUANG T C BORTZ E L and MCCAY C M Observations on Blood Vessels and Exercise J Geront 11 797 1956
 - 605 FILATOV V P and BARG TS M (Appearance of Biogenic Stimulators during Muscular Work) Abstracted in Abstracts of Sov Med Part A—Basic Med Sci 1 26 1957
 - 606 KRAUS H HIRSCHLAND B P and HIRSCHHORN K Role of Inactivity in Production of Disease Hypokinetic Disease J Am Geriatr Soc 4 463 1956
 - 607 MANN G V and WHITE H S The Influence of Stress on Plasma Cholesterol Levels Metabolism 2 47 1953
 - 608 SELLERS E A and YOW R W Deposition of Fat in Coronary Arteries after Exposure to Cold Brit Med J 1 815 1956
 - 609 HOLLISTER L E The Effect of Adrenergic Blocking Agents (Including Chlorpromazine) on Serum Lipid Levels of Patients with Disorders of Fat Metabolism J Chron c Dis 6 234 1957
 - 610 WILENS S L MCCLUSKEY A T and SOMOZA C Inhibitory Effect of Chlorpromazine on Arterial Lipid Deposition in Cholesterol Fed Rabbits Proc Soc Exper Biol and Med 93 121 1956
 - 611 KAPLAN A and GANT M Effect of Epinephrine Administration upon Blood Lipids Fed Proc 14 87 1955
 - 612 DURY A and MOSS L D Experimental Arteriosclerosis Effect of Repeated Intravenous Injections of Epinephrine on Lipid Partition of Plasma and Liver and Aortic Degenerative Changes in Intact Rabbits J Geront 9 287 1954
 - 613 ROSENMAN R H and FRIEDMAN M Change in the Serum Cholesterol and Blood Clotting Time in Men Subjected to Cyclic Variation of Emotional Stress Circulation 16 931 1957
 - 614 Capitalist Cholesterol MD Medical Newsmagazine 1 14 1957
 - 615 HAMMARSTEN J F CATHEY C W REDMOND R F and WOLF S G JR Serum Cholesterol Diet and Stress in Patients with Coronary Artery Disease Circulation 16 890 1957

- 616 ALTSCHUL R Inhibition of Experimental Cholesterol Arteriosclerosis by Ultra Violet Irradiation *New Eng J Med* 249 96 1953
- 617 YOUNG D R and COOK S F Body Lipids in Small Mammals following Prolonged Exposures to High and Low Temperatures *Am J Physiol* 181 72 1955
- 618 SODEMAN W A and LOGUF J T Effect of Environmental Temperature upon Cholesterol Induced Atherosclerosis in the Rabbit *J Lab Clin Med* 46 954 1955
- 619 REDMOND R P MILCH L J and CALHOUN W W Plasma Lipoprotein Changes Induced by Mild Local Cold Injury *J Lab Clin Med* 42 112 1953
- 620 HEWITT J E HAGES T L GOFMAN J W JONES H B and PIERCE F T Effect of Total Body Irradiation upon Lipoprotein Metabolism *Cardiologia* 21 353 1952
- 621 MILCH L J YURNELL R A and STENSON J V Relationship between Blood Lipids and Radiation Injury in Rabbits *Science* 120 713 1954
- 622 MILCH L J REDMOND R F and CALHOUN W W Blood Lipoproteins in Traumatic Injury *J Lab Clin Med* 43 603 1954
- 623 HAMMOND E C and HORN D The Relationship Between Human Smoking Habits and Death Rates A Follow Up Study of 187 766 Men *J A M A* 155 1316 1954
- 624 ——— Smoking and Death Rates—Report on Forty four Months of Follow up of 187 783 Men *J A M A* 166 1159 1958
- 625 RUSSEK H I ZOHMAN B L and DORSET V J Effects of Tobacco and Whiskey on the Cardiovascular System *J A M A* 157 563 1955
- 626 GOFMAN J W LINDGREN F T STRISOWER B DE LALLA O GLAZIER F and TAMPLIN A Cigarette Smoking Serum Lipoproteins and Coronary Heart Disease *Geriatrics* 10 349 1955
- 627 DOLGOFF S SCHREK R BALLARD G P and BAKER L A Tobacco Smoking as an Etiologic Factor in Disease II Coronary Disease and Hypertension *Angiology* 3 323 1952
- 628 SIGLER L H Tobacco as a Contributing Cause of Degenerative Coronary Disease *New York State Med J* 55 310 1955
- 629 SCHAEFER L E DRACHMAN S R STEINBERG A G and ADLERSBERG D Genetic Studies on Hypercholesteremia Frequency in a Hospital Population and in Families of Hypercholesteremic Index Patients *Am Heart J* 46 99 1953
- 630 SCHAEFER L E and ADLERSBERG D Genetic and Epidemiologic Studies in Cholesterol Metabolism Contributions of a Prepaid Medical Care Program *Am J Pub Health* 46 966 1956
- 631 ZEMAN F D Genetic Factors in the Diseases of Later Life *J Chronic Dis* 2 11 1955
- 632 THOMAS C B and COHEN B H The Familial Occurrence of Hypertension and Coronary Artery Disease with Observations Concerning Obesity and Diabetes *Ann Int Med* 4 90 1955
- 633 Editorial The Genetic Background of Chronic Diseases *J Chronic Dis* 2 96 1955
- 634 TANNER J M The Relation Between Serum Cholesterol and Physique in Healthy Young Men *J Physiol* 115 371 1951
- 635 WONG H Y C JOHNSON F B WONG A ANDERSON J and LIU D Effect of Exercise and Androgen on Cholesterol Fed Capons *Circulation* 16 501 1957
- 636 FISHER L M MOGENSON G J and JAKUES L B Effect of Stress and Dicumaryl on Prothrombin Time of Rats *Fed Proc* 16 296 1957
- 637 SHERBER D A and MARCUS M Studies on the Physical State of the Serum Lipids under Stress *Circulation* 16 491 1957
- 638 WEXLER B C and MILLER B F Severe Arteriosclerosis Produced in the ACTH Treated Rat *Circulation* 16 950 1957
- 639 TAYLOR H L ANDERSON J T and KEYS A Diet Physical Activity and Serum Cholesterol in Man *Circulation* 16 516 1957
- 640 WOLFFE J B Continued Vigorous Physical Activity as a Possible Factor in the Prevention of Atherosclerosis *Circulation* 16 517 1957
- 641 ADLERSBERG D SCHAEFER L E and STEINBERG A G Studies on Genetic and Environmental Control of Serum Cholesterol Level *Circulation* 16 487 1957

- 642 WHEELER E O The Genetic Aspects of Atherosclerosis *Am J Med* 23 653 1957
- 643 HIRSCHHORN K and WILKINSON C F JR Mode of Inheritance in Essential Familial Hypercholesterolemia *Circulation* 16 509 1957
- 644 BOYLE E JR and McLAUGHLIN H M Differentiation of Types of Idiopathic Hypercholesterolemia by Serum Lipoprotein Distribution and Response to Therapy *Circulation* 16 867 1957
- 645 STAMLER J PICK R and KATZ L N Experiences in Assessing Estrogen Antiatherogenesis in the Chick the Rabbit and Man *Ann New York Acad Sci* 64 596 1956
- 646 BARR D P Influence of Sex and Sex Hormones upon the Development of Atherosclerosis and upon the Lipoproteins of Plasma *J Chronic Dis* 1 63 1955
- 647 STAMLER J PICK R and KATZ L N Effects of Desoxycorticosterone Acetate on Cholesterolemia Blood Pressure and Atherogenesis in Chicks *Circulation* 4 262 1951
- 648 STAMLER J and KATZ L N The Effect of Salt Hypertension on Atherosclerosis in Chicks Fed Mash without a Cholesterol Supplement *Circulation* 3 859 1951
- 649 WAKERLIN G E MOSS W G and KIELY J P Effect of Experimental Renal Hypertension on Experimental Thiouracil Cholesterol Atherosclerosis in Dogs *Circulation Res* 5 426 1957
- 650 STAMLER J and HALL Y Epidemiological Studies on the Cardiovascular Renal Diseases IV Findings of a Special Query Concerning Chicago Deaths In preparation

B Blood Clotting Mechanisms

- 651 LEVER W F SMITH P A J and JURLEY N A Effects of Intravenous Heparin on the Plasma Lipoproteins in Primary Hypercholesteremic Xanthomatosis and Idiopathic Hyperlipemia *Science* 118 653 1953
- 652 LINDGREN F T NICHOLS A V and FREEMAN N K Physical and Chemical Composition Studies on the Lipoproteins of Fasting and Heparinized Human Sera *J Physical Chem* 59 930 1955
- 653 ANFINSEN C B JR Biochemical Aspects of Atherosclerosis *Fed Proc* 15 894 1956
- 654 BROWN W D Effects of Heparin and Protamine on Alimentary Lipaemia in Rats *Acta Physiol Scand* 30 374 1954
- 655 SHORE B Heparin Activated Enzyme System of Human Plasma *Proc Soc Exper Biol and Med* 88 73 1955
- 656 GROSSMAN M I The Effect of Heparin on the Fate of Intravenously Administered Fat Emulsions *J Lab Clin Med* 40 805 1955
- 657 GROSSMAN M I STADLER J CUSHING A and PALM L Relation of Lipolysis to Deturbidification and Returbidification in Heparin Induced Lipemia Clearing Phenomenon *Proc Soc Exper Biol and Med* 88 137 1955
- 658 GROSSMAN M I MOELLER H C and PALM L Effects of Lipemia and Heparin on Free Fatty Acid Concentration of Serum in Humans *Proc Soc Exper Biol and Med* 90 106 1955
- 659 McDANIEL R A and GROSSMAN M I Paper Electrophoretic Study of C¹⁴ Fat Emulsion Cleared from Post Heparin Rat Plasma *Proc Soc Exper Biol and Med* 89 442 1955
- 660 YOUNG W and FREEMAN N K Effects of Heparin on Triglyceride and Free Fatty Acid Concentrations in Lymph *Proc Soc Exper Biol and Med* 90 463 1955
- 661 SPITZER J J and MILLER H J Unesterified Fatty Acids and Lipid Transport in Dogs *Proc Soc Exper Biol and Med* 97 124 1956
- 662 FREEMAN L ENGELBERG H and DUDLEY A Plasma Heparin Levels *Am J Clin Path* 24 599 1954

- 663 ENGELBERG H Studies of Normal Endogenous Lipemia Clearing Factor Am J Physiol 181 309 1955
- 664 ——— Correlation of Plasma Heparin Levels with Serum Lipoproteins Acta Med Scand 151 161 1955
- 665 PIERCE F T JR The Interconversion of Serum Lipoproteins *In Vivo* Metabolism 3 142 1954
- 666 BRADGON J H and HAVEL R J *In Vivo* Effects of Anti Heparin Agents on Serum Lipids and Lipoproteins Am J Physiol 177 128 1954
- 667 FULTON G P MAYNARD F L RILEY J F and WEST G B Humoral Aspects of Tissue Mast Cells Physiol Revs 37 221 1957
- 668 NIKKILA E A and MAJANEN S The Blood Heparinoid Substances in Human Atherosclerosis Scand J Clin and Lab Invest 4 204 1952
- 669 THOMAS W A and OLWIN J H Variations in Protamine Titration Results in the Blood of Atherosclerotic Patients Personal Communication
- 670 ENGELBERG H Human Plasma Heparin Levels Following Intravenous Fat Emulsions Circulation 16 481 1957
- 671 ANGERVALL G and HOOD B Studies on Heparin and Lipemia Clearing Factor Inhibition towards Clearing in Plasma from Normal and Atherosclerotic Individuals Acta Med Scand 157 407 1957
- 672 THOMAS W A Quantitative Protamine Titration Test (Heparin) of Blood in Atherosclerosis J Lab Clin Med 50 958 1957
- 673 DAY A J WILKINSON G K GILMORE H R and SCHWARTZ C J Effect of Protamine on Alimentary Lipemia Circulation 16 72 1957
- 674 KORN E D and QUIGLEY T W JR Studies on Lipoprotein Lipase of Rat Heart and Adipose Tissue Biochem Biophys Acta Amst 18 143 1955
- 675 ANFINSEN C B JR BOYLE E and BROWN R K The Role of Heparin in Lipoprotein Metabolism Science 115 583 1952
- 676 GRAHAM D M LYON T P GOFMAN J W JONES H B YANKLEY A SIMON J and WHITE S Blood Lipids and Human Atherosclerosis II The Influence of Heparin upon Lipoprotein Metabolism Circulation 4 666 1951
- 677 HORITA A and LOOMIS T A The Influence of Anticoagulant on the Formation and Regression of Experimental Atherosclerosis J Exper Med 100 381 1951
- 678 CONSTANTINIDES P SZASZ G and HARDER F Retardation of Atheromatosis and Adrenal Enlargement by Heparin in the Rabbit A M A Arch Path 56 36 1953
- 679 CONSTANTINIDES P Mast Cells and Susceptibility to Experimental Atherosclerosis Science 117 505 1953
- 680 CAIRNS A and CONSTANTINIDES P Mast Cells in Human Atherosclerosis Science 120 31 1954
- 681 MENG H C and DAVIS W S Effects of Heparin on Development of Atherosclerosis and Fatty Liver a Report of Studies on Cholesterol Fed Rabbits A M A Arch Path 60 276 1955
- 682 DAVIS O F OESTER Y T and FRIEDMAN B Influence of Adenosine Triphosphate Adenosine Monophosphate and Heparin in Experimental Arteropathy Circulation Res 3 374 1955
- 683 OPDYKE D F ROSENBERG A SILBER R OTT W H and SIEGEL H Effect of Chronic Injection of a Heparin Complex on Aortic Atherosclerosis in Cholesterol Fed Chickens J Lab Clin Med 45 271 1955
- 684 HORLICK L and DUFF G L Heparin in Experimental Cholesterol Atherosclerosis in the Rabbit A M A Arch Path 57 417 1954
- 685 CLOSS K and DEDICHEN J Vascular Disease Lancet 257 1242 1949
- 686 STANLER J PICK R and KATZ L N Effects of Heparin and Dicumarol on Atherogenesis in Cholesterol Fed Cockerels Clin Res Proc 5 175 1957

C Thyroid Hormone

- 687 PICK R STANLER J and KATZ L N Effects of Hypothyroidism on Estrogen Induced Inhibition of Coronary Atherogenesis in Cholesterol Fed Cockerels Circulation Res 5 510 1957

- 688 STAMLER J PICK R and KATZ L N Influence of Thyroid Pancreatic and Adrenal Hormones on Lipid Metabolism and Atherosclerosis in Experimental Animals *Hormones and Atherosclerosis* New York Academic Press in press
- 689 PICK R STAMLER J and KATZ L N Influence of Estrogens on Lipids and Atherosclerosis in Experimental Animals *Hormones and Atherosclerosis* New York Academic Press in press

D Estrogens

- 690 STAMLER J and KATZ L N The Plasma Cholesterol/Phospholipid Ratio and Atherosclerosis in the Chick *Am J Physiol* 163 752 1950
- 691 PICK R STAMLER J ROBBARD S and KATZ L N The Inhibition of Coronary Atherosclerosis by Estrogens in Cholesterol Fed Chicks *Circulation* 6 276 1952
- 692 ——— Estrogen Induced Regression of Coronary Atherosclerosis in Cholesterol Fed Chicks *Circulation* 6 585 1952
- 693 MALINOW R HOJMAN D PELLEGRINO A PICK R STAMLER J and KATZ L N Histological and Histochemical Studies on the Effects of Estrogens on Coronary Atherogenesis in Cholesterol Fed Cockerels In preparation
- 694 ROBBARD S PICK R STAMLER J and KATZ L N Hypercholesterolemia and Atherogenesis in Sexually Immature Male and Female Chicks In preparation
- 695 STAMLER J PICK R and KATZ L N Inhibition of Cholesterol Induced Coronary Atherogenesis in the Egg Producing Hen *Circulation* 10 251 1954
- 696 PICK R STAMLER J and KATZ L N Susceptibility of the Ovariectomized Hen to Cholesterol Induced Coronary Atherogenesis *Circulation Res* 5 515 1957
- 697 STAMLER J PICK R and KATZ L N Prevention of Coronary Atherosclerosis by Estrogen Androgen Administration in the Cholesterol Fed Chick *Circulation Res* 1 94 1953
- 698 ——— Estrogen Prophylaxis of Cholesterol Induced Coronary Atherogenesis in Chicks Given Adrenal Corticoids or ACTH *Circulation* 10 241 1954
- 699 STAMLER J Estrogen Inhibition of Corticoid Hypertension in Chickens *Circulation* 10 896 1954
- 700 MOSKOWITZ M S MOSKOWITZ A E BRADFORD W L JR and WISSLER R W Changes in Serum Lipids and Coronary Arteries of the Rat in Response to Estrogens *A M A Arch Path* 61 245 1956
- 701 MALINOW M R and PELLEGRINO A A The Effect of Estrogens on Spontaneous and Experimental Atherosclerosis in the Rat *Circulation* 14 491 1956
- 702 Anonymous Influence of Sex Hormones on Coronary Artery Disease in Rats *Nutr Revs* 15 140 1957
- 703 CHAKRAVARTI R N DE U N and MUKERJI B Studies in Experimental Atherosclerosis III Effect of Desiccated Thyroid and Oestrogen on the Regression of Experimental Cholesterol Atherosclerosis *Ind J Med Res* 41 683 1956
- 704 EILERT M L Effect of Estrogens on the Partition of Serum Lipids in Female Patients *Metabolism* 2 137 1953
- 705 BARR D P Some Chemical Factors in the Pathogenesis of Atherosclerosis *Circulation* 8 641 1953
- 706 ——— Hormonal Factors in the Pathogenesis of Atherosclerosis *A Symposium on Arteriosclerosis* p 58 Minneapolis Minnesota Heart Association and University of Minnesota 1955
- 707 OLIVER M F and BOYD G S Changes in the Plasma Lipids during the Menstrual Cycle *Clin Sci* 12 217 1953
- 708 ——— Plasma Lipid and Serum Lipoprotein Patterns during Pregnancy and Puerperium *Clin Sci* 14 15 1955
- 709 ——— Effect of Estrogens on the Plasma Lipids in Coronary Artery Disease *Am Heart J* 47 348 1954
- 710 ——— The Influence of the Sex Hormones on the Circulating Lipids and Lipoproteins in Coronary Sclerosis *Circulation* 13 82 1956
- 711 ——— Endocrine Aspects of Coronary Sclerosis *Lancet* 2 1273 1956
- 712 STEINER A PAYSON H and KENDALL F E Effect of Estrogenic Hormone on Serum Lipids in Patients with Coronary Arteriosclerosis *Circulation* 11 784 1955
- 713 WLEST J DRY T J and EDWARDS J E The Degree of Coronary Sclerosis in Bilaterally Oophorectomized Women *Circulation* 7 801 1953

- 714 RIVIN A U and DIMITROFF S P The Incidence and Severity of Atherosclerosis in Estrogen Treated Males and in Females with a Hypoestrogenic or a Hyperestrogenic State *Circulation* 9 533 1954
- 715 MARMORSTON J LEWIS J J BERNSTEIN J L SOBEL H KUZMA O ALEXANDER R MAGIDSON O and MOORE F J Excretion of Urinary Steroid by Men and Women with Myocardial Infarction *Geriatrics* 12 297 1957
- 716 AIDIN R Coronary Artery Disease *Lancet* 270 447 1956
- 717 FURMAN R H HOWARD R P and IMAGAWA R Serum Lipid and Lipoprotein Concentrations in Castrate and Non Castrate Institutionalized Male Subjects *Circulation* 14 490 1956
- 718 FURMAN R H HOWARD R P SHETLAR M R and IMAGAWA R Serum Lipoproteins Glycoproteins and Lipids in Institutionalized Eunuchs and Non Castrate Male Subjects *Am J Med* 22 965 1956
- 719 COOPER E E Sex Hormones and Atherogenesis *Am Pract* 7 436 1956

Definitive Approaches to Prophylaxis and Therapy of Atherosclerotic Disease in Man

- 720 STAMLER J Diet in the Long Term Clinical Management of Myocardial Infarction (Part of a Panel Discussion) *J Am Geriatrics Soc* 5 879 1957
- 721 STAMLER J PICK R KATZ L N PICK A and KAPLAN B Interim Report on Clinical Experiences with Long Term Estrogen Administration to Middle Aged Men with Coronary Heart Disease *Hormones and Atherosclerosis* New York Academic Press in press
- 722 STEAD E S and WARREN C K *Low Fat Cookery* New York McGraw Hill Book Co Inc 1956
- 723a DOBBIN E V GOFMAN H F JONES H C LYON L and YOUNG C B *The Low Fat Low Cholesterol Diet* New York Doubleday & Co 1951
- 723b GOFMAN J W NICHOLS A V and DOBBIN E V *Dietary Prevention and Treatment of Heart Disease* New York G P Putman's Sons 1958
- 724 KUO P T and JOYNER C R Angina Pectoris Induced by Fat Ingestion in Patients with Coronary Artery Disease *JAMA* 158 1008 1955
- 725 POLLAK O J Reduction of Blood Cholesterol in Man *Circulation* 7 702 1953
- 726 BEST M M DUNCAN C H VAN LOON E J and WATHEN J Lowering of Serum Cholesterol by the Administration of a Plant Sterol *Circulation* 10 210 1954
- 727 BEST M M DUNCAN C H WATHEN J D VAN LOON E J and SHIPLEY R E Effects of Prolonged Administration of Sitosterol on Serum Lipids *Circulation* 10 590 1954
- 728 BEST M M DUNCAN C H VAN LOON E J and WATHEN J D The Effects of Sitosterol on Serum Lipids *Am J Med* 19 61 1955
- 729 BEST M M and DUNCAN C H Modification of Abnormal Serum Lipid Patterns in Atherosclerosis by Administration of Sitosterol *Ann Int Med* 45 614 1956
- 730 JOYNER C R JR. and KUO P T The Serum Cholesterol Level and Lipoprotein Pattern Following the Administration of Plant Sterols *Circulation* 10 589 1954
- 731 ——— The Effect of Sitosterol Administration upon the Serum Cholesterol Level and Lipoprotein Pattern *Am J Med Sci* 230 636 1955
- 732 KUO P T and JOYNER C R JR The Effect of Low Fat Diet and Sitosterol on the Serum Lipids of Patients with Coronary and Peripheral Arterial Diseases *Circulation* 14 499 1956
- 733 LESESNE J M CASTOR C W and HOOBLER S W Prolonged Reduction in Human Blood Cholesterol Levels Induced by Plant Sterols *U Michigan Med Bull* 21 13 1955
- 734 BARBER J M and GRANT A P The Serum Cholesterol and Other Lipids after Administration of Sitosterol *Brit Heart J* 17 296 1955
- 735 SACHS B A and WESTON R E Sitosterol Administration in Normal and Hypercholesteremic Subjects The Effect in Man of Sitosterol Therapy on Serum Lipids and Lipoproteins *AMA Arch Int Med* 97 738 1956

- 736 FARQUHAR J W SMITH R E and DEMPSEY M E The Effect of Beta Sitosterol on the Serum Lipids of Young Men with Arteriosclerotic Heart Disease *Circulation* 14 77 1956
- 737 SHIPLEY R E The Effects of Sitosterol on Serum Cholesterol Concentration *Trans New York Acad Sci* 18 111 1955
- 738 WILKINSON C F BOYLE E JACKSON R S and BENJAMIN M R Effect of Varying Intake of Dietary Fat and the Ingestion of Sitosterol on Lipid and Lipoprotein Fractions of Human Serum Metabolism 4 307 1955
- 739 WILKINSON C F Effects of Lipotropes and Sitosterol on the Level of Blood Lipids and the Clinical Course of Angina Pectoris *J Am Geriatrics Soc* 3 381 1955
- 740 Anonymous The Effect of Sitosterols on the Serum Cholesterol of Man *Nutr Revs* 14 39 1956
- 741 SCHROEDER H A A Practical Method for the Reduction of Plasma Cholesterol in Man *J Chronic Dis* 4 461 1956
- 742 BEST M M and DUNCAN C H Observations on the Mechanism of the Hypocholesterolemic Effect of Sitosterol *Circulation* 14 911 1956
- 743 LEHMANN J H Clinical Experiences with Beta Sitosterol a New Anticholesterolic Agent *Northwest Med* 56 46 1957
- 744 PERRY H M JR and SCHROEDER H A Depression of Cholesterol Levels in Human Plasma Following Ethylenediamine Tetracetate and Hydralazine *J Chronic Dis* 2 520 1955
- 745 KING J S JR CLARKSON T B and WARNOCK N H Effect on Experimental Avian Atherosclerosis of Dietary Oils and Gallogen *Proc Soc Exper Biol and Med* 93 443 1956
- 746 ——— The Hypocholesterolizing Effects of Drugs—Fed Singly and in Combination—on Cholesterol Fed Cockerels *Circulation Res* 4 162 1956
- 747 COTTET J and MATHIVAT A L'Acide Phenyl Ethyl Acetique Est Il Anti Atherogene? *Presse Med* 63 1005 1955
- 748 KRITCHEVSKY D MAYER A W TESAR W C MCCANDLESS R F J LOGAN J B BROWN R A and ENGLERT M The Effect of Sodium 2 Phenyl Butyrate in Experimental Atherosclerosis *Angiology* 7 156 1956
- 749 ENGELBERG H and MASSELL T B Heparin in the Treatment of Advanced Peripheral Atherosclerosis *Am J Med Sci* 225 14 1953
- 750 ENGELBERG H KUHN R and STEINMAN M A Controlled Study of the Effect of Intermittent Heparin Therapy on the Course of Human Coronary Atherosclerosis *Circulation* 13 489 1956
- 751 SUZMAN M M RUSKIN H D and GOLDBERG B An Evaluation of the Effect of Continuous Long Term Anticoagulant Therapy on the Prognosis of Myocardial Infarction A Report of 82 Cases *Circulation* 12 338 1955
- 752 KEYES J W DRAKE E H and SMITH F J Survival Rates after Acute Myocardial Infarction with Long Term Anticoagulant Therapy *Circulation* 14 254 1956
- 753 MANCHESTER B The Prevention of Myocardial Infarction *A M A Arch Int Med* 100 959 1957
- 754 DEDICHER J Long Term Anticoagulant Treatment of Cardiovascular Disorders with Phenylindandrone (Pid Trombantin) *Scand J Clin and Lab Invest* 5 324 1953
- 755 LUND E Long Term Dicumarol Therapy in Thromboembolic Conditions Especially Coronary Thrombosis *Acta Med Scand* 146 252 1953
- 756 BAY E B WRIGHT P A JONES R J PAGE R G and SERDE M Use of Anticoagulant Drugs in Ambulatory Patients *Circulation* 9 741 1954
- 757 NICHOL E S PHILLIPS W C and JENKINS V E Symposium on Efficacy of New Drugs Anticoagulants in Coronary Heart Disease *Med Clin N Amer* 38 399 1954
- 758 OWREN P A The Protracted and Permanent Anticoagulant Therapy *Nord Med cin* 54 1733 1955
- 759 BENGTSEN K K and ASPENSTRÖM G Preliminary Report on Long Term Treatment of Cardiosclerosis with Dicumarol *Acta Med Scand* 157 217 1957

- 714 RIVIN A U and DIMITROFF S P The Incidence and Severity of Atherosclerosis in Estrogen Treated Males and in Females with a Hypoestrogenic or a Hyperestrogenic State *Circulation* 9 533 1954
- 715 MARMORSTON J LEWIS J J BERNSTEIN J L SOBEL H KUZMA O ALEXANDER R MAGIDSON O and MOORE F J Excretion of Urinary Steroids by Men and Women with Myocardial Infarction *Geriatrics* 12 297 1957
- 716 AIDIN R Coronary Artery Disease *Lancet* 270 447 1956
- 717 FURMAN R H HOWARD R P and IMAGAWA R Serum Lipid and Lipoprotein Concentrations in Castrate and Non Castrate Institutionalized Male Subjects *Circulation* 14 490 1956
- 718 FURMAN R H HOWARD R P SHETLAR M R and IMAGAWA R Serum Lipoproteins Glycoproteins and Lipids in Institutionalized Eunuchs and Non Castrate Male Subjects *Am J Med* 22 965 1956
- 719 COOPER E E Sex Hormones and Atherogenesis *Am Pract* 7 436 1956

Definitive Approaches to Prophylaxis and Therapy of Atherosclerotic Disease in Man

- 20 STAMLER J Diet in the Long Term Clinical Management of Myocardial Infarction (Part of a Panel Discussion) *J Am Geriatrics Soc* 5 879 1957
- 721 STAMLER J PICK R KATZ L N PICK A and KAPLAN B Interim Report on Clinical Experiences with Long Term Estrogen Administration to Middle Aged Men with Coronary Heart Disease *Hormones and Atherosclerosis* New York Academic Press in press
- 722 STEAD E S and WARREN G K *Low Fat Cookery* New York McGraw Hill Book Co Inc 1956
- 723a DOBBIN E V GOFMAN H F JONES H C LYON L and YOUNG C B *The Low Fat Low Cholesterol Diet* New York Doubleday & Co 1951
- 723b GOFMAN J W NICHOLS A V and DOBBIN E V *Dietary Prevention and Treatment of Heart Disease* New York G P Putman & Sons 1958
- 724 KLO P T and JOYNER C R Angina Pectoris Induced by Fat Ingestion in Patients with Coronary Artery Disease *JAMA* 158 1008 1955
- 725 POLLAK O J Reduction of Blood Cholesterol in Man *Circulation* 7 702 1953
- 726 BEST M M DUNCAN C H VAN LOON E J and WATHEN J Lowering of Serum Cholesterol by the Administration of a Plant Sterol *Circulation* 10 210 1954
- 727 BEST M M DUNCAN C H WATHEN J D VAN LOON E J and SHIPLEY R E Effects of Prolonged Administration of Sitosterol on Serum Lipids *Circulation* 10 590 1954
- 728 BEST M M DUNCAN C H VAN LOON E J and WATHEN J D The Effects of Sitosterol on Serum Lipids *Am J Med* 19 61 1955
- 729 BEST M M and DUNCAN C H Modification of Abnormal Serum Lipid Patterns in Atherosclerosis by Administration of Sitosterol *Ann Int Med* 45 614 1956
- 730 JOYNER C R JR and KLO P T The Serum Cholesterol Level and Lipoprotein Pattern Following the Administration of Plant Sterols *Circulation* 10 589 1954
- 731 ——— The Effect of Sitosterol Administration upon the Serum Cholesterol Level and Lipoprotein Pattern *Am J Med Sci* 230 636 1955
- 32 KLO P T and JOYNER C R JR The Effect of Low Fat Diet and Sitosterol on the Serum Lipids of Patients with Coronary and Peripheral Arterial Diseases *Circulation* 14 499 1956
- 733 LESENE J M CASTOR C W and HOOBLER S W Prolonged Reduction in Human Blood Cholesterol Levels Induced by Plant Sterols *U Michigan Med Bull* 21 13 1955
- 734 BARBER J M and GRANT A P The Serum Cholesterol and Other Lipids after Administration of Sitosterol *Brit Heart J* 17 296 1955
- 735 SACHS B A and WESTON R E Sitosterol Administration in Normal and Hypercholesteremic Subjects The Effect in Man of Sitosterol Therapy on Serum Lipids and Lipoproteins *AMA Arch Int Med* 97 738 1956

- 784 MORRIS J N and HEADY J A Prognosis of Coronary Heart Disease in Medical Practitioners *Brit Heart J* 19 227 1957
- 785 KATZ L N STAMLER J and PICK R Approaches to the Problem of the Relation of Emotions to Hormonal Function and Atherosclerosis *Hormones and Atherosclerosis* New York Academic Press in press

Addenda

- 786 PICK R STAMLER J and KATZ L N Production of Ulcerated Hemorrhagic Atherosclerotic Plaques and Dissecting Aneurysms in the Abdominal Aorta of Cockerels *Fed Proc* 17 124 1958
- 787 PICK R and KATZ L N Further Observations on Ulcerated Arteriosclerotic Lesions of the Abdominal Aorta in Cockerels *Circulation* 18 481 1958

- 160 WAALER B A The Effect of Permanent Anticoagulant Therapy on Symptoms and Mortality in Angina Pectoris *Acta Med Scand* 157 289 1957
- 761 FOLEY W T McDEVITT E SYMONS C and WRIGHT I S Further Experience with Long Term Anticoagulant Therapy *A M A Arch Int Med* 95 497 1955
- 762 TULLOCH J and WRIGHT I S Long Term Anticoagulant Therapy—Further Experiences *Circulation* 9 823 1954
- 763 MORROW E J PAINTER J T and LEDBETTER P V Prolonged Anticoagulant Therapy in the Management of Coronary Artery Disease *Circulation* 17 750 1955
- 764 GRIFFITH G C DOOLEY J V ENGELBERG H ANDERSON R and ZINN W J Treatment of Myocardial Infarction with Heparin *Circulation* 16 508 1957
- 765 GILCHRIST R Personal Communication
- 766 CHANDLER H L and MANN G V Heparin Treatment of Patients with Angina Pectoris—Failure to Influence either the Clinical Course or the Serum Lipids *New Eng J Med* 249 1045 1953
- 767 GRUNER A HILDEN T RASSCHOU F and VOGELIUS H Heparin Treatment of Angina Pectoris *Am J Med* 14 433 1953
- 768 RINZLER S H TRAVELL J BAKST H BENJAMIN Z H ROSENTHAL R L ROSENFELD S and HIRSCH B B Effect of Heparin in Effort Angina *Am J Med* 14 438 1953
- 769 POST M P KATZ A HELLMAN E and ENSELBERG C P The Heparin Treatment of Angina Pectoris *Am Heart J* 45 769 1953
- 770 KORY R C TOWNES A S MABE R E DORRIS E R and MENEELY G R Drug Evaluation in Angina Pectoris *Akella Heparin Penitrate Am Heart J* 50 308 1955
- 771 BROWN K W G and RYKERT H E Heparin in the Treatment of Angina Pectoris *Canad Med Assoc J* 70 617 1954
- 772 CORCORAN A C LEWIS L A DUSTAN H P and PAGE I H Atherosclerotic Complications in Hypertensive Disease *Ann N Y Acad Sci* 64 463 1956
- 773 SKOUBY A P Vascular Lesions in Diabetics with a Special Reference to the Influence of Treatment *Acta Med Scand* 155 Supp 317 1956
- 774 STRISOWER B GOFMAN J W GALTON E RUBINGER J H O'BRIEN G W and SIMON A Effect of Long Term Administration of Desiccated Thyroid on Serum Lipoprotein and Cholesterol Levels *J Clin Endocrinol* 15 73 1955
- 775 STRISOWER B GOFMAN J W CALSON E F PLININGER J H POLTEAL J and GUZVICH P Long Term Effect of Desiccated Thyroid Substance upon Serum Lipoproteins and Serum Cholesterol Printed for the U S Atomic Energy Commission UCRL—3534 1956
- 776 BLOMGART H L FREEBERG A S and KURLAND G S Hypothyroidism Produced by Radioactive Iodine (I^{131}) in the Treatment of Euthyroid Patients with Angina Pectoris and Congestive Failure *Circulation* 1 1105 1950
- 777 STAMLER J PICK R KATZ L N KAPLAN B and PICK A Evaluation of Estrogen Therapy in Males with Previous Myocardial Infarction—Interim Report Four Year Follow Up *Circulation* 16 940 1957
- 778 GODDEN J O and HINES E A JR Studies of Intermittent Claudication II Effect of Androgen and an Androgen Estrogen Combination in the Treatment of Intermittent Claudication *Proc Staff Meet Mayo Clin* 30 491 1955
- 779 OLIVER B B and FRIEDMAN B The Effect of Estrogen Therapy on Patients with Angina Pectoris *Am J Med Sci* 231 205 1956
- 780 BOSSAE E T WANG C and ADLERSBERG D Prolonged Low Dose Estrogen Therapy in Idiopathic Hyperlipemia and Hypercholesterolemia *Circulation* 16 503 1957
- 781 ROBB G P and MARKS H H What Happens to Men Disabled by Heart Disease? *Tr Assoc Life Insur Med Dir America* (1953) 37 171 1954
- 782 MILLS G Y SIMON A J CISNEROS F and KATZ L N Myocardial Infarction—Observations on One Hundred Patients Who Survived up to Six Years *A M A Arch Int Med* 84 632 1949
- 783 COLE D R SINGAN E B and KATZ L N The Long Term Prognosis Following Myocardial Infarction and Some Factors which Affect It *Circulation* 9 321 1954

Index

A

- ANIMAL fats cholesterolemia and 53 54
- Anticoagulants 98
- Antihypertensives 99
- Atherogenesis diet and 15 61
 - experimental 60 63 70
 - prerequisites for 15
- Atherosclerosis animal experimental
 - findings in 61
 - blood clotting and 82
 - clinical pathological findings and 44
 - diet and 80
 - economics and 19
 - epidemiologic findings in 16 19
 - estrogens and 83
 - experimental 64
 - fibrinolytic mechanisms and 82
 - heredity and 77
 - high cholesterol diet and 62
 - high fat diet and 62
 - hormonal factors in 76
 - in Bantu 22
 - income and 38
 - occupation and 37
 - physical activity and 79
 - prophylaxis of 93
 - smoking and 78
 - therapy of 93
 - thyroid hormone and 83

B

- BRAIN extracts 69

C

- CARIOVASCULAR RENAL mortality foreign
 - born and 36
 - Negro and 35
 - sex and 32
 - trend of 31
- Cholesterol dietary 63
 - serum in several countries 25
- Cholesterolemia diet and 25
 - dietary effects on 50 52

- Corn germ 69
- Coronary artery disease unsaturated
 - fatty acids and 60
 - heart disease lipoprotein levels in 46 47
 - morphologic atherosclerosis and 44
 - serum cholesterol in 46 47
- sclerosis 26
- Cotton seed oil and cholesterol 65

D

- DESICCATED thyroid effect on cholesterol
 - levels 100
- Diabetes atherosclerosis in 45
 - control 100
- Diet curtailment of 108
 - in Guatemala 22 24 38
 - in Italy 24 26 38
 - in Japan 24 25 38
 - in Low Countries 26 30
 - in Scandinavia 26 30
 - in Spain 20 24
 - low cholesterol 48 95 96
 - low fat 95 96
 - effect on cholesterolemia 48
 - methionine deficiency 72
 - optimal 18
 - social differences and 20
- Dihydrocholesterol 69

E

- EMIGRANTS death rate 77
- Emigration effect of on Italians 26
 - on Japanese 26
 - on Yemenite Jews 26
- Essential familial xanthomatosis 45
 - fatty acids 64
- Estrogen induced reversal of atherosclerosis 86
- Estrogens 101
 - and atherogenesis 83
 - combined with other hormones 88
 - effect of on mammalian species 90
 - in man 91
 - serum effects of 103 104

Estrogens therapeutic experiments with 86

F

- FAT calories from 25
- intake 108
- Fats saturated 18 22
- Fatty acids and unsaturated oils 56
- Food consumption in U S A 33

H

- HORMONAL factors in atherosclerosis 76 83
- Hypercholesterolemia 107
- diet induced 61
- Hypertension 107

I

- INCOME and food consumption in U S A 38

L

- LIPOTROPIC factors 98
- Low cholesterol diets 95 96
- effect on cholesterolemia 48
- Low fat diets 95 96

M

- MARINE oils 50
- Methionine deficient diet 72
- Myocardial infarction anticoagulants and 99
- in Yemenite Jews 27

N

- NEPHROSIS atherosclerosis in 45
- Nutrition ecological background 16
- historical background 15

O

- OBESITY 41 42 107
- correction of 94
- mortality and 95
- Occupation and arteriosclerotic heart disease 37

Oils saturated vs unsaturated 57

unsaturated 97

P

- PHYSICAL activity and atherosclerosis 49
- Plasma lipids lipoproteins and atherogenesis 91
- Polyenoic fatty acids 69
- Prophylaxis 106

R

- RICE diet and serum cholesterol 50

S

- SATURATED vs unsaturated oils 57
- Serum lipids pre treatment with 107 103
- Sitosterols 69 97
- Smoking and atherosclerosis 78
- Social classes and serum cholesterol levels 39

T

- THYROID hormone 101
- and atherogenesis 83

U

- UNDERNUTRITION 17
- Unsaturated oils 97
- vs saturated oils 60

V

- VEGETABLE fats cholesterolemia and 53
- 54
- oils 50
- unsaturated 64
- Vitamin high protein supplement 73 14

W

- WEIGHT gain and loss and cholesterolemia 48 49

X

- XANTHOMATOSIS essential familial 45

